

# Kettenbruch Continued Fractions

<https://github.com/Nazgand/nazgandMathBook>

Mark Andrew Gerads: [Nazgand@Gmail.Com](mailto:Nazgand@Gmail.Com)

October 1, 2022

## Abstract

The goal of this paper is to have fun reviewing Kettenbruch notation for continued fractions.

## 1 Introduction

The 2 defining equations are:

$$\mathop{\mathrm{K}}\limits_{k=n}^n \frac{a_k}{b_k} = \frac{a_n}{b_n} \quad (1.1)$$

$$\mathop{\mathrm{K}}\limits_{k=n}^m \frac{a_k}{b_k} = \frac{a_n}{b_n + \mathop{\mathrm{K}}\limits_{k=n+1}^m \frac{a_k}{b_k}} \quad (1.2)$$

The most interesting continued fractions are infinite continued fractions, like so, yet careful consideration must be taken in this case, e.g. because the limit might not exist:

$$\mathop{\mathrm{K}}\limits_{k=1}^{\infty} \frac{a_k}{b_k} = \lim_{m \rightarrow \infty} \mathop{\mathrm{K}}\limits_{k=1}^m \frac{a_k}{b_k} = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \ddots}}} \quad (1.3)$$

## 2 Examples

One of the simplest cases to consider is  $a_k = a \in \mathbb{C}, b_k = b \in \mathbb{C}, b \neq 0$ .

$$x = \mathop{\mathrm{K}}\limits_{k=1}^{\infty} \frac{a}{b} = \frac{a}{b + \mathop{\mathrm{K}}\limits_{k=1}^{\infty} \frac{a}{b}} \Rightarrow x^2 + bx - a = 0 \quad (2.1)$$

