



Subject Areas:

behaviour, computer vision,
health and disease and
epidemiology

Keywords:

broiler chickens, animal welfare,
E. coli, deep learning, food safety

Author for correspondence:

David Alejandro Duchêne
e-mail: david.duchene@sund.ku.dk

Deep learning from videography as a tool for measuring *E. coli* infection in poultry

Neil Scheidwasser^{1†}, Louise Ladefoged Poulsen^{2†}, Prince Ravi Leow¹, Mark Poulsen Khurana¹, Maider Iglesias-Carrasco³, Daniel Joseph Laydon^{4,5}, Christl Ann Donnelly⁶, Anders Miki Bojesen^{2*}, Samir Bhatt^{1,4*}, David Alejandro Duchêne^{1*}

¹Section of Health Data Science and AI, Department of Public Health, University of Copenhagen, 1353 Copenhagen, Denmark

²Section of Veterinary Clinical Microbiology, Department of Veterinary and Animal Sciences, University of Copenhagen, 1870 Frederiksberg, Denmark

³Center for Evolutionary Hologenomics, Globe Institute, University of Copenhagen, 1353 Copenhagen, Denmark

⁴MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London, W12 0BZ, United Kingdom

⁵Centre for Health Economics & Policy Innovation, Department of Economics & Public Policy, Imperial College Business School, Imperial College London, London, SW7 2AZ, United Kingdom

⁶Department of Statistics, University of Oxford, Oxford, OX1 3LB, United Kingdom

Poultry farming is threatened by regular outbreaks of *Escherichia coli* (*E. coli*) that lead to significant economic losses and public health risks. However, traditional surveillance methods often lack sensitivity and scalability. Early detection of infected poultry using minimally invasive procedures is thus essential for preventing epidemics. To that end, we leverage recent advancements in computer vision, employing deep learning-based tracking to detect behavioural changes associated with *E. coli* infection in a case-control trial comprising two groups of 20 broiler chickens: (1) a healthy control group and (2) a group infected with a pathogenic *E. coli* field strain from the poultry industry. More specifically, kinematic features derived from deep learning-based tracking data revealed markedly reduced activity in the challenged group compared to the negative control. These findings were validated by lower mean optical flow in the infected flock, suggesting reduced movement and activity, and post-mortem physiological markers of inflammation which confirmed the severity of infection in the challenged group. Overall, this study demonstrates that deep learning-based tracking offers a promising solution for real-time monitoring and early infection detection in poultry farming, with the potential to help reduce economic losses and mitigate public health risks associated with infectious disease outbreaks in poultry.

1. Introduction

Poultry farming is a cornerstone of the agriculture industry and is essential to global nutritional security, accounting for nearly 35% of the world's meat production [1]. In many poultry-producing regions, *Escherichia coli* (*E. coli*) infection is a leading cause of morbidity and mortality, including China [2,3], the United States [4], Brazil [5], and several European countries such as the United Kingdom [6] and Denmark [7]. In addition, *E. coli* outbreaks are linked with intensive use of antibiotics, promoting the selection of resistant strains that have the potential to spill over into human populations [8]. Consequently, detecting early signs of *E. coli* infection in domesticated birds, such as broiler chickens, is crucial for safeguarding animal welfare, minimising economic losses, and protecting public health.

Over recent decades, a plethora of techniques have been proposed to leverage data from physiological markers (e.g., body temperature [9,10]) or behavioural outputs (e.g., locomotion, sound [11]) to assess welfare impairments and detect pathogen outbreaks. Measuring temperature requires precise thermal imaging cameras, which are expensive and sensitive to environmental disturbances. In contrast, acquiring sound and video recordings is logistically simpler and less expensive, but usually requires computer-intensive postprocessing to extract meaningful behavioural features. In addition, obtaining precise estimates of individual patterns remains challenging. For instance, commonly used video processing methods such as optical flow have been leveraged to provide valuable insights on behaviour [12,13], bacterial infection [14,15], footpad dermatitis and hockburn [16], and general animal welfare [17–19]. However, optical flow only allows for analysis at the level of a flock and not of the individual. Additionally, as it relies on the average displacement of pixels across consecutive frames, it is therefore prone to capturing irrelevant movements from the environment. To obtain individual-level measurements, gait-scoring systems have been developed for broiler chickens [20,21], but are impractical for real-time applications. Other frameworks based on markers or wearable sensors (e.g., accelerometers or radio-frequency identification (RFID) chips [22]) are promising for their high resolution, but are intrusive and not easily scalable to industrial settings.

Alternatively, recent progress in deep learning [23] for computer vision has catalysed the development of versatile frameworks for tracking and markerless pose estimation (i.e., without pre-placed physical markers) of individual [24–26] or multiple animals simultaneously [27–29]. These methods have garnered significant interest in behavioural neuroscience [30] and ecology [31]. As to animal behaviour and welfare monitoring, the potential advantages of deep learning-based videography analysis are multi-factorial. Indeed, state-of-the-art solutions are non-invasive, adaptable to various environments, user-friendly, and may allow for individual-level analysis with near-human-level precision. Although these systems initially require data-intensive training of deep neural networks on videos of interest, transfer learning allows for fast deployment of pre-trained models on new video data.

To this end, we develop a feasibility study that demonstrates the capacity of deep learning-based, markerless tracking from video recordings to detect phenotypical and physiological differences in broiler chickens following *E. coli* infection. We designed a trial in which 40 chickens were randomly assigned to two, equal-sized, separately-housed groups: (1) a healthy control group (control), and (2) a group infected with a pathogenic *E. coli* strain (Fig. 1). We used the multi-animal version of DeepLabCut [28] to identify behavioural changes associated with infection and found that *E. coli*-infected chickens moved and travelled less in their pens and spent less time near the food source (used as a proxy to measure feeding time). This finding was first validated with a more traditional optical flow-based analysis, which corroborated the lower activity in the infected group, and second with necropsy data which revealed distinct patterns of *E. coli*-caused internal lesions and inflammation in the infected group. Thus, our data provide evidence that deep learning on video is a useful and objective tool for detecting the behavioural and physiological changes associated with infection. The simplicity and objectivity of the statistics gleaned from videos, including distance travelled and changes in body area, indicate




Figure 1. Experimental setup. (a) Timeline of the experiment. (b) Pipeline for behavioural feature extraction using DLCRNet, the standard model of the multi-animal version of DeepLabCut (maDLC) [28]. (c) Diagram of the chicken coops. (d) Schematised top view of a broiler chicken with standardised body parts used for tracking and polygon used to approximate body area. (e) Example predictions from maDLC, based on one frame.

that scalability to industrial settings is feasible. Addressing sensitivity to other pathogens, high animal densities, and trade-offs between labelling individual animals and image resolution will be crucial for implementing reliable deep learning-based outbreak detection methods in broiler farms.

2. Methods

2.1 Animals and housing conditions

40 day-old, *E. coli*-unvaccinated, Ross 308 broiler chickens (mixed gender) were obtained from Blenta AB (Blentarp, Sweden) and followed for 38 days (Fig. 1a). The animals were accommodated in two groups of 20 (see [Experimental design](#)) in an 8.64 m² coop (3.6 m x 2.4 m; Fig. 1c) with a dust bath to a 16h light (06h-22h):1h dim (22.00-23.00):7h dark (23.00-06.00) schedule. Feed was provided *ad libitum* twice daily to accommodate the increased growth (from 15 to 200 grams/day) according to the feeding schedule of Aviagen [32]. The feed consisted of commercial whole-grain food for broiler chickens aged 0-8 weeks (Brogaarden, Lynge, Denmark). Water was available *ad libitum* at two different sources. Standard procedures were conducted to prevent cross-contamination from other pathogens.

2.2 Experimental design

On the day of arrival, each chicken was randomly assigned to one of two groups: uninfected control group, or infected with *E. coli*. For the infected group, a pathogenic strain of *E. coli* known for causing colibacillosis in broilers [33–35] (strain ST117 E44; accession number LXWV00000000.1) was administered intratracheally (9.9×10^5 colony-forming units (CFUs)/chicken in 0.5 ml) (week 5 of the experiment; Fig. 1a), while chickens from the control group were sham-challenged intratracheally (phosphate-buffered saline; (PBS)) following the same timeline. The strain used for inoculation was provided by the Section of Veterinary Clinical Microbiology at the University of Copenhagen and prepared as described in [35] (storage at -80°C , streaking on blood agar base supplemented with 5% bovine blood (BA) and incubation at 37°C overnight). A single colony was incubated in brain heart infusion (BHI) broth (Oxoid BO1230B (Thermo Fisher Scientific)) overnight at 37°C . Subsequently, 1 ml of the culture was transferred to 100 ml BHI and incubated until the optical density at 600 nm (OD_{600}) reached 1.56 (approximately two hours). At $OD_{600} = 1.56$ (exponential phase), 308 μL of culture was transferred to 199.69 ml PBS (Merck, 524650) and a volume of 0.5 ml per animal was used as inoculum and kept on ice until infection. The inoculum's CFU count was confirmed by 10-fold dilutions prepared in triplicates plated (100 μL) on BA, and incubated overnight before counting.

2.3 Biological data sampling

Blood samples were collected twice from the brachial vein: firstly a day before the challenge and secondly before euthanasia for analysis of serum amyloid A using an LZ-SAA assay (Eiken Chemical). Blood samples were centrifuged at 1,500 RPM for 10 min to isolate the serum, which was thereafter stored in a freezer (-20°C) until analysis. Euthanasia was performed by cervical dislocation directly following the induction of unconsciousness by blunt head trauma. Gross pathology was performed two days post-infection during the necropsy in a randomised and blinded manner to mitigate experimenter biases. An overall lesion score (see Table S1) derived from previous *E. coli* studies [35–37] was calculated by visually assessing and grading pathological lesions in the peritoneum, air sacs, lungs and spleen. To measure pulmonary *E. coli* CFU counts, each lung was aseptically removed and placed in a sterile stomacher bag, weighed, and then blended with a 1/1 (weight/vol) solution of 0.9% PBS. The resulting blend was serially diluted tenfold (10^{-1} – 10^{-7}) and $3 \times 30 \mu\text{L}$ per dilution was spotted separately for overnight incubation. CFU counts were normalised by lung weight.

2.4 Behavioural data sampling

For each group, five videos were recorded over three days: two in the first two days (morning and afternoon, after feeding) and one in the third morning (Fig. 1a). Each recording lasted between 20 and 30 minutes and was acquired at 30 frames per second (FPS) at a resolution of 1080×1920 pixels using GoPro HERO10 Black cameras (<https://gopro.com>). The cameras were placed 130 cm away from the floor. For each video, the first two minutes were removed to minimise any potential experimenter intervention.

2.5 DeepLabCut training and inference

Deep learning for animal tracking was implemented in markerless fashion using the multi-animal version of DeepLabCut [28], which allows tracking and pose estimation of multiple objects simultaneously. For each video, each frame from a single video is first clustered using *k*-means clustering, and twenty frames were sampled from the formed clusters (100 frames is considered sufficient to achieve reasonable tracking performance according to [24]). For each frame, annotations for five body parts (head, centre left, centre right, saddle, and tail) were

added for each animal. We used a pre-trained *DLCRNet_ms5* [28], which consists of a multi-scale convolutional neural network (CNN) stacked on top of another CNN (ResNet-50 [38]) for feature extraction and stacked with deconvolutional layers to predict where the body parts of interest were and which animals they belonged to. The multi-scale architecture fuses high and low-resolution maps, thus allowing for robust and accurate detection of body parts by leveraging fine-grained detail and broader contextual information. The model was fine-tuned in end-to-end fashion (i.e., all model weights were tunable) with recommended hyperparameters from [28] (Adam [39] optimisation with multi-stage learning rate (η) scheduling (stage 1: $\eta = 1 \times 10^{-4}$ for 7,500 iterations; stage 2: $\eta = \text{1} \times \text{10}^{\text{5e-5}}$ 5×10^{-5} for 12,000 iterations; stage 3: $\eta = 1 \times 10^{-5}$ until the end of the experiment), batch size of 8) using 95% of the labelled frames (split randomly) for 200,000 training iterations. Importantly, the test set was completely held out during model development. The model was trained only once on the training set, and the test frames were accessed solely after training was complete. The root mean squared errors (RMSEs) were: 3.03 pixels (train set) and 8.68 pixels (test set) for an image size of 1080×1920 pixels. The trained network was then used to extract the (x, y) positions of each body part for each animal in all videos. An example frame is shown in Figure 1e.

2.6 Post-processing

For each video, DeepLabCut outputs an array of (x-coordinate, y-coordinate, and likelihood) triplets of size $F \times (N \times B \times 3)$, where F is the number of frames, N the number of animals, and B the number of body parts. Predictions with a likelihood below 0.9 were dropped following standard DeepLabCut practice [24]. To recover some of the dropped values, forward linear interpolation was applied at a body part level, where at most 15 consecutive missing values could be filled (0.5 s). (x, y) positions were finally smoothed using Savitzky-Golay filtering with a polynomial order of 7 and a window length of 15 frames to strike a trade-off between preserving the sharpness of the original signal and reducing noise. Subsequently, two kinematic features could be extracted (Fig. 1b) for each chicken c in video v : the total distance travelled $L_{.,v,c}$ to quantify locomotor activity and the total rate of change in body area $\dot{A}_{.,v,c}$, for non-locomotor activity (e.g., stretching for improved ventilation).

$$L_{.,v,c} = \sum_{i=1}^N \frac{d(\bullet_{i,v,c} - \bullet_{i-1,v,c})}{d(\text{head}_{i,v,c}, \text{saddle}_{i,v,c})}$$

$$\dot{A}_{.,v,c} = \sum_{i=1}^N \frac{|\dot{A}_{i,v,c}|}{d(\text{head}_{i,v,c}, \text{saddle}_{i,v,c})}$$

$$\bullet_{i,v,c} = \left(\frac{1}{4}(x_{i,v,c,\text{head}} + x_{i,v,c,\text{centreleft}} + x_{i,v,c,\text{centreright}} + x_{i,v,c,\text{saddle}}), \right. \\ \left. \frac{1}{4}(y_{i,v,c,\text{head}} + y_{i,v,c,\text{centreleft}} + y_{i,v,c,\text{centreright}} + y_{i,v,c,\text{saddle}}) \right)$$

The distance travelled in each frame i is calculated as the Euclidean distance $\mathbf{d}$ between the chicken's centre of mass $\bullet$ in the current frame and the previous frame $i - 1$. The centre of mass is defined as the average (x, y) position of four key points: the head, centre-left, centre-right, and saddle. For each frame i , both $L_{i,v,c}$ and $\dot{A}_{i,v,c}$ are normalised by the distance between the head and saddle in each respective frame to account for potential radial and tangential distortions due to the camera setup. The frame-wise rate of change in body area was measured in a similar fashion to [40]. The shoelace formula was employed to calculate the area of a polygon defined by the four above-mentioned key points (Fig. 1d). The rate of change of that area $\dot{A}_i$ was measured as a discrete temporal derivative of the body area, obtained numerically using a Savitzky-Golay filter. Subsequently, the time spent near the food source (assumed as a proxy for time spent eating) was estimated for each animal by computing the number of frames in which the animal's

head was in the vicinity of the food source (Fig. 1c). The vicinity was defined as a rectangle whose scale was 1.05 times that of the food source in the video. Future work should investigate more direct measures of feeding behaviour, e.g., by detecting pecking or other related actions, to increase the accuracy of this measurement.

2.7 Statistical analysis

For each physiological covariate, individual differences were tested using non-parametric Mann-Whitney U tests. The *t*-test assumption of normality was assessed using quantile-quantile plots (Q-Q plots), while Levene's test was used to test for homoscedasticity. For all markers, both normality and homoscedasticity could not be reasonably assumed for lesion scores and SAA levels, leading us to opt for Mann-Whitney U tests to assess between-group disparities. Cohen's *d* was reported to reflect the effect size for all pairwise comparisons.

For each behavioural feature, a Bayesian mixed-effects model was employed to investigate between-group variations (with the control group as a reference) while accounting for random variations across the different recordings. Our mixed effects model had the following linear predictor

$$\mu_{ij} = \alpha + \beta X + \gamma_{0i}$$

where μ_{ij} is a behaviour feature of interest (e.g., distance travelled) for a day *i* and individual *j*, α is an intercept that corresponds to the control group, β is a scalar for the group *X* (infected or not), and γ is a random effect for the *i*th day. As the distance travelled and body area change were normalised by the body length (thus constituting rates distributed in (0, 1)), we assumed their values were generated from a Beta distribution. Whilst the time spent near the food source is also a rate, a zero-inflated version of the Beta distribution was used to account for the frequent zero-valued observations. All models were fitted with the default prior distributions suggested by **brms** [41]. To facilitate efficient sampling and minimise divergent transitions, a target acceptance rate of 0.99 was chosen and a maximum tree depth of 15 was selected for building during the trajectory phase of the Hamiltonian Monte Carlo sampler. For all variables, we compared the beta family models (with a logit link function) against models with the default family parameter (Gaussian with an identity link function) in **brms**. We selected the best family based on a Bayesian estimate of the expected log pointwise predictive density (ELPD) from leave-one-out cross-validation (LOOCV) (Table S4). Finally, convergence plots and posterior predictive checks were plotted to check model fit (see Fig. S1 to S4). For all variables, the models fitted using a beta distribution were preferable.

2.8 Optical flow

To complement deep learning-based animal tracking with more traditional computer vision methods, we measured optical flow, a commonly used technique to determine the average motion of chickens in a video as a means to assess animal welfare [12,14,42]. Optical flow works by estimating the motion of pixels in a video by comparing consecutive image frames. Here, the optical flow was computed using a state-of-the-art adaptation of the Recurrent All Pairs Field Transforms (RAFT) model [43] called SEA-RAFT [44]. This model consists of a deep neural network (DNN) trained to predict the optical flow between consecutive frames. It features a residual neural network (ResNet-34 [38]) to extract relevant features from pairs of images, a correlation block to compute similarities between pixels, and a recurrent neural network (RNN) to iteratively refine the flow field predictions from the correlation block outputs. Compared to traditional, handcrafted methods such as the Horn-Schunck [45] or Farnebäck methods [46], which are based on energy minimisation, deep learning-based approaches have been posited to be more robust to occlusions and rapid movement [47].

Here, we used a version of SEA-RAFT that was pre-trained on a collection of datasets including simulated moving objects, photorealistic scenes, and recorded scenes from autonomous

driving vehicles [44]. To reduce inference time, we downsampled the images by a factor of 2 ($h = 960 \times w = 540$ pixels) before feeding them into the model. For each pair of frames, the output consists of an array of size $2 \times h \times w$, which denotes the estimated (x, y) flow for each pixel. From this output, following previous optical flow-based animal welfare studies [12,14,42], the first four moments (mean, variance, skewness, and kurtosis) of the optical flow norms were measured for each frame. The statistics were finally averaged across frames for between-group comparisons, and the average mean optical flow was regressed using ordinary least squares (OLS) against the distance travelled measured from DeepLabCut-based tracking.

2.9 Implementation

Video pre-processing was performed using `ffmpeg` (<https://ffmpeg.org>) and tracking was implemented using the DLCRNet model [28] implemented in DeepLabCut [24] (version 2.3.5). Model training on our video dataset was performed on a single NVIDIA RTX A6000 graphical processing unit (GPU). Optical flow was computed frame by frame using an implementation of a pre-trained SEA-RAFT model [44] in PyTorch [48] (version 2.2.0). Feature extraction and statistical tests were performed using custom scripts in Python 3.10.10 using NumPy [49] (version 1.25.0), SciPy [50] (version 1.11.1), and Pingouin [51] (version 0.5.4). Bayesian mixed-effects modelling was implemented in R 4.4.0 using the `brms` package [41] (version 2.21.0). The code for reproducing feature extraction and statistical analyses is available at <https://github.com/Neclow/dlc4ecoli>.

3. Results

The goal of this study was to determine whether deep learning-based behavioural analysis from video can be used as a proxy for monitoring and detecting animal welfare impairments (here, caused by *Escherichia coli* (*E. coli*) infection). To this end, we recorded two groups of 20 animals: a negative control group and a group infected with *E. coli*. In particular, we compared standard physiological measurements obtained post-mortem (see Methods) against behavioural features extracted from multi-animal DeepLabCut [28].




Figure 2. Comparison of physiological biomarkers of inflammation in response to *E. coli* challenge. Significance levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Post-mortem examination of the chickens by necropsy revealed significant disparities between the groups in terms of lesion score, SAA levels, and number of colony-forming units (CFUs) (Fig. 2; Mann-Whitney U: $U_{\text{lesion}} = 39.5$, $U_{\text{SAA}} = 45.0$, $U_{\text{cfu}} = 31.0$; p -values < 0.001) with medium to large effect sizes (Cohen's d : $d_{\text{lesion}} = -2.121$, $d_{\text{SAA}} = -1.153$, $d_{\text{cfu}} = -0.607$). Chickens from the control group had near-zero presence of SAA and CFUs as well as very few lesions, while *E. coli*-infected groups showed significantly higher values for all markers (Table S2).




Figure 3. Comparison of behavioural features in response to *E. coli* challenge. Top: posterior distributions from Bayesian mixed-effects modelling to investigate between-group variations while accounting for random variations across the different recordings, predicting the group (infected or not). One point refers to one individual in one video. Points from the first recording (performed on challenge day) recording are marked in dark red. Middle: mean estimates of the same behaviour features for the *E. coli* group, with the negative control group serving as the reference. Bottom: average treatment effect for each behaviour feature, computed as a ratio between the *E. coli* and control groups. Error bars indicate 95% confidence intervals.

Behavioural differences between groups in the distance travelled, body area, and time near the food source (extracted from DeepLabCut-based animal tracking) were evaluated using Bayesian mixed-effects modelling, with random effects applied to the recording time (Fig. S1–S4). The infected group was found to be significantly less active than the uninfected control group as measured by the distance travelled (average treatment effect (ATE) ratio: 24% reduction (95% credible interval (CrI): 6–39%)) and the rate of change in body area (ATE ratio: 23% reduction (95% CrI: 4–39%)) (Fig. 3; Table S3). Additionally, we posit that the infected group fed less as its individuals spent less time near the food source on average (ATE ratio: 41% reduction (95% CrI: 24–56%); Fig. 3). Individual chicken identity was not recovered from one recording to another, such that a chicken labelled “chicken X” in a given recording could not be re-identified in subsequent recordings, preventing within-subject analyses.

The activity measured by deep learning-based individual tracking was validated against optical flow, a traditional method in computer vision used to measure object motion based on pixel intensity changes between consecutive frames (Fig. 4). Optical flow data from videos (Control: $n = 5$, Infected: $n = 5$) of the control group exhibited higher mean optical flow (Fig. 4a), indicating higher average activity [14]. Whilst not exhibiting higher optical flow kurtosis as in [14], control individuals also displayed higher optical flow variance, indicating that infected individuals had more uniform behaviour. The optical flow results were generally consistent with DeepLabCut tracking-based results, seen in the positive correlation between the mean optical flow and the mean distance travelled (Fig. 4b to 4d; group- and time-independent Pearson correlation: $r = 0.636$, $p = 0.048$; $BF_{10} = 2.17$). Overall, despite the limited validation




Figure 4. Validation of the travelled distance measured using the multi-animal DeepLabCut (maDLC) [28] against optical flow statistics extracted from a pre-trained SEA-RAFT model [43]. (a) Summary statistics (mean, variance, skewness, kurtosis) averaged over each frame and video for each group. Points from the first recording (performed on challenge day) recording are marked in dark red. (b) Comparison of the mean optical flow (across frames) against the mean distance travelled (across frames and animals) over time. (c) Normalised cross-correlation between the average distance travelled measured using maDLC and the average optical flow. (d) Group- and time-independent regression of the mean optical flow (across frames) against the mean distance travelled (across frames and animals). Pearson correlation (r): 0.636 ($p = 0.048$). In (b) and (d), the shaded areas represent bootstrapped confidence intervals (95%) around the estimates (number of bootstraps = 1,000). In each panel, one point refers to one 20 to 30-min video (see [Behavioural data sampling](#)).

provided by optical flow due to sample size limitations, deep learning-based video analysis could be deemed effective at discriminating behavioural patterns between the control and infected groups. Further, the impact of infection was confirmed by traditional post-mortem analysis of infection biomarkers.

4. Discussion

In this proof-of-concept study, we demonstrate the capacity of deep learning-based, markerless tracking from video recordings to detect phenotypical differences in broiler chickens following *E. coli* infection. This method holds significant promise for promptly identifying infection or compromised animal well-being at the individual level, without human intervention or intrusive sensors [22] - a capability previously limited to flock-level analyses by conventional techniques [14,17]. Moreover, the simplicity and objectivity of the statistical analyses derived from video data suggest promising avenues for future research in disease detection and management within the poultry industry. In our analysis, three intuitive features from multi-animal DeepLabCut tracking [28] revealed distinct behavioural patterns between *E. coli* infected and uninfected chickens, corresponding to observed differences in common physiological markers of infection, including serum amyloid A levels, *E. coli* concentrations and organ lesion scores.

These features were further validated against optical flow analysis, a traditional computer vision approach that is also promising for measuring animal welfare at industrial scales [12–19].

Both deep learning and optical flow pointed towards greater activity levels in control animals, reflecting a considerable impact of *E. coli* infection on the nervous system. Our data on raised SAA in infected chickens provides evidence that the root cause of behavioural change is likely induced by cytokine-mediated inflammation. Other forms of neurotoxicity that were not explored might also be at play, such as from shiga toxins [52], or gut-brain axis disruption [53]. Previous work has found lower mean optical flow in chickens infected by *Campylobacter* [14], while we observe slight changes in variance, but not in kurtosis after *E. coli* infection. Lower variance in infected individuals is consistent with reduced long-distance movement, while maintained kurtosis indicates stable core short-distance behaviours. Further investigation with larger samples and replicates across different centres will be crucial to confirm the robustness and generalisability of deep learning-based approaches. Additionally, these methods hold significant potential for farms and the poultry industry, particularly for real-time detection of welfare deterioration signs, such as lameness or infections caused by other pathogens.

Identifying infections using videography presents several challenges, some of which are inherent to the livestock industry, where individual animal identification is labour-intensive. Accordingly, we did not preserve bird identity across recordings and therefore did not perform within-subject analyses. Although mixed-effects modelling was used to compare trends between groups, analysing individual differences in behavioural and physiological data would facilitate high-resolution analyses. This warrants future work with more replicates and lifelong video monitoring, or the use of individual features for identification at maximal video resolution. Similarly, we focused on coarse behaviours that might not reflect the behavioural nuances of the immune state of individuals. Future research could investigate a broader spectrum of behavioural responses, e.g., pecking activity, to achieve more precise measurements of feeding behaviour.

Scaling up deep learning-based pose estimation methods to farm-level settings can be achieved via a series of technological and practical solutions. A critical aspect is the ability to perform long-term, continuous monitoring, with the goal of maintaining individual identity over time and potentially detecting early signs of decreased welfare on a per-animal basis. Current tracking of individual body parts becomes challenging when the number of individuals is large (e.g., > 50), but solutions are being quickly developed to mitigate the errors arising from crowded environments [54]. Other tracking frameworks, such as TRex [27], allow for tracking up to 256 individuals with reasonable precision, but do not track individual body parts and are still insufficient for poultry farm-level conditions where several thousand animals can coexist. To that end, we posit that a hybrid solution combining a coarse-grained technique, such as optical flow, to detect zones of lower activity with a high-resolution animal tracking framework to identify impaired individuals could constitute a viable solution for curbing outbreaks in farms and thus minimise potential economic losses.

5. Conclusion

In this study, we showcase a framework for deep learning-based, markerless tracking of broiler chickens for individual-level detection of behavioural changes associated with *E. coli* infection. The behavioural differences observed in an infected group vs. a control group were validated against physiological markers of infection. More generally, we believe our findings demonstrate the potential of deep learning-based video analysis as an objective tool for outbreak detection in industrial farming, which could be scaled jointly with current flock-level techniques such as optical flow. Livestock settings are complex, and yet advances in data science will continue to disentangle the links between infection, the nervous system, and behaviour for industrial applications. The simplicity and objectivity of the statistical analyses derived from video data, coupled with advancements in deep learning technology, suggest promising avenues for future research in disease outbreak detection and management in livestock.

Ethics. The study was approved by the Danish Animal Experiments Inspectorate under the Danish Ministry of Environment and Food, and all animal procedures were performed by this approval (license no. 2019-15-0201-01611) and with ARRIVE guidelines. The licence granted, and guidelines hereof are in agreement with the Danish law on animal experiments and the EU directive 2010/63. Predefined humane endpoints were determined and animals were observed every 30 min for the initial six hours after inoculation, subsequently at maximally eight-hour intervals for three days after inoculation with increased frequency in the event of clinical signs in any of the groups. If clinical signs were present, e.g., ruffled feathers, depression, anorexia, lethargy or dyspnoea, the bird was either treated with 0.1 mg/kg buprenorphine and observed with increased frequency or euthanised.

Data Accessibility. Data and relevant code for this research work are stored in GitHub: <https://github.com/Neclow/dlc4ecoli> and have been archived within the Zenodo repository: <https://doi.org/10.5281/zenodo.16795387> [55].

Authors' Contributions. N.S., L.L.P., A.M.B., S.B., D.A.D.: study conception and design. L.L.P., N.S.: practical execution. L.L.P., A.M.B.: necropsies and microbiology. N.S.: data analysis. M.P.K., M.I-C., P.R.L.: interpretation of results. L.L.P., A.M.B., D.A.D., S.B.: supervision. N.S., M.P.K., S.B., D.A.D.: writing of the original draft. N.S., L.L.P., M.P.K., M.I-C., P.R.L., D.J.L., C.A.D., A.M.B., S.B., D.A.D.: edition of the original draft.

Competing Interests. The authors declare no competing interests.

Funding. SB and DJL acknowledge support from the MRC Centre for Global Infectious Disease Analysis (MR/R015600/1), jointly funded by the UK Medical Research Council (MRC) and the UK Foreign, Commonwealth & Development Office (FCDO), under the MRC/FCDO Concordat agreement, and also part of the EDCTP2 programme supported by the European Union. SB is funded by the National Institute for Health Research (NIHR) Health Protection Research Unit in Modelling and Health Economics, a partnership between the UK Health Security Agency, Imperial College London and LSHTM (grant code NIHR200908). Disclaimer: "The views expressed are those of the author(s) and not necessarily those of the NIHR, UK Health Security Agency or the Department of Health and Social Care." SB acknowledges support from the Novo Nordisk Foundation via The Novo Nordisk Young Investigator Award (NNF20OC0059309). SB acknowledges support from the Danish National Research Foundation via a chair grant (DNRF160) which also supports NS and MPK. SB acknowledges support from The Eric and Wendy Schmidt Fund For Strategic Innovation via the Schmidt Polymath Award (G-22-63345) which also supports PRL. SB and NS acknowledge the Pioneer Centre for AI as affiliate researchers. DAD is funded by a Novo Nordisk Fonden Data Science Investigator Award (NNF23OC0084647). DJL acknowledges funding from the Wellcome Trust UK for the Vaccine Impact Modelling Consortium (VIMC) Climate Change Research Programme (grant ID: 226727_Z_22_Z). AMB acknowledges support from the Innovation Fund Denmark (Grant no. 071-00001) and the Danish Poultry Levy Foundation (Grant no. 122656).

Acknowledgements. The authors thank Alexandros Katsiferis for valuable insight on analysing the *brms* outputs.

References

1. OECD, FAO. 2023 *OECD-FAO Agricultural Outlook 2023-2032*. OECD. (doi:10.1787/08801ab7-en)
2. Liu C, Wang P, Dai Y, Liu Y, Song Y, Yu L, Feng C, Liu M, Xie Z, Shang Y et al. 2021 Longitudinal monitoring of multidrug resistance in *Escherichia coli* on broiler chicken fattening farms in Shandong, China. *Poult. Sci.* **100**, 100887. (doi:10.1016/j.psj.2020.11.064)
3. Zhou W, Lin R, Zhou Z, Ma J, Lin H, Zheng X, Wang J, Wu J, Dong Y, Jiang H et al. 2022 Antimicrobial resistance and genomic characterization of *Escherichia coli* from pigs and chickens in Zhejiang, China. *Front. Microbiol.* **13**, 1018682. (doi:10.3389/fmicb.2022.1018682)
4. Fancher CA, Zhang L, Kiess AS, Adhikari PA, Dinh TT, Sukumaran AT. 2020 Avian pathogenic *Escherichia coli* and *Clostridium perfringens*: Challenges in no

- antibiotics ever broiler production and potential solutions. *Microorganisms* **8**, 1533. (doi:10.3390/microorganisms8101533)
5. Braga JFV, Chanteloup NK, Trotereau A, Baucheron S, Guabiraba R, Ecco R, Schouler C. 2016 Diversity of *Escherichia coli* strains involved in vertebral osteomyelitis and arthritis in broilers in Brazil. *BMC Vet. Res.* **12**, 1–12. (doi:10.1186/s12917-016-0762-0)
 6. Kemmett K, Williams N, Chaloner G, Humphrey S, Wigley P, Humphrey T. 2014 The contribution of systemic *Escherichia coli* infection to the early mortalities of commercial broiler chickens. *Avian Pathol.* **43**, 37–42. (doi:10.1080/03079457.2013.866213)
 7. Kromann S, Baig S, Olsen RH, Edslev SM, Thøfner I, Bojesen AM, Jensen HE, Stegger M. 2023 Dramatic increase in slaughter condemnations due to *Escherichia coli* ST23 and ST101 within the Danish broiler production. *Vet. Microbiol.* **280**, 109696. (doi:10.1016/j.vetmic.2023.109696)
 8. Roth N, Käsbohrer A, Mayrhofer S, Zitz U, Hofacre C, Domig KJ. 2019 The application of antibiotics in broiler production and the resulting antibiotic resistance in *Escherichia coli*: A global overview. *Poult. Sci.* **98**, 1791–1804. (doi:10.3382/ps/pey539)
 9. Okada H, Itoh T, Suzuki K, Tsukamoto K. 2009 Wireless sensor system for detection of avian influenza outbreak farms at an early stage. In *Proc. SENSORS* pp. 1374–1377. IEEE. (doi:10.1109/ICSENS.2009.5398422)
 10. Noh JY, Kim KJ, Lee SH, Kim JB, Kim DH, Youk S, Song CS, Nahm SS. 2021 Thermal image scanning for the early detection of fever induced by highly pathogenic avian influenza virus infection in chickens and ducks and its application in farms. *Front. Vet. Sci.* **8**, 616755. (doi:10.3389/fvets.2021.616755)
 11. Rowe E, Dawkins MS, Gebhardt-Henrich SG. 2019 A systematic review of precision livestock farming in the poultry sector: Is technology focussed on improving bird welfare?. *Animals* **9**, 614. (doi:10.3390/ani9090614)
 12. Dawkins MS, Lee HJ, Waitt CD, Roberts SJ. 2009 Optical flow patterns in broiler chicken flocks as automated measures of behaviour and gait. *Appl. Anim. Behav. Sci.* **119**, 203–209. (doi:10.1016/j.applanim.2009.04.009)
 13. Gronskyte R, Clemmensen LH, Hviid MS, Kulahci M. 2016 Monitoring pig movement at the slaughterhouse using optical flow and modified angular histograms. *Biosystems Engineering* **141**, 19–30. (doi:10.1016/j.biosystemseng.2015.10.002)
 14. Colles FM, Cain RJ, Nickson T, Smith AL, Roberts SJ, Maiden MC, Lunn D, Dawkins MS. 2016 Monitoring chicken flock behaviour provides early warning of infection by human pathogen *Campylobacter*. *Proc. R. Soc. B* **283**, 20152323. (doi:10.1098/rspb.2015.2323)
 15. Fernández-Carrión E, Martínez-Avilés M, Ivorra B, Martínez-López B, Ramos ÁM, Sánchez-Vizcaíno JM. 2017 Motion-based video monitoring for early detection of livestock diseases: The case of African swine fever. *PLoS ONE* **12**, e0183793. (doi:10.1371/journal.pone.0183793)
 16. Dawkins M, Roberts S, Cain R, Nickson T, Donnelly C. 2017 Early warning of footpad dermatitis and hockburn in broiler chicken flocks using optical flow, bodyweight and water consumption. *Vet. Rec.* **180**, 499–499. (doi:10.1136/vr.104066)
 17. Stamp Dawkins M, Donnelly CA, Jones TA. 2004 Chicken welfare is influenced more by housing conditions than by stocking density. *Nature* **427**, 342–344. (doi:10.1038/nature02226)
 18. Roberts SJ, Cain R, Dawkins MS. 2012 Prediction of welfare outcomes for broiler chickens using Bayesian regression on continuous optical flow data. *J. R. Soc. Interface* **9**, 3436–3443. (doi:10.1098/rsif.2012.0594)
 19. Dawkins MS, Cain R, Merelie K, Roberts SJ. 2013 In search of the behavioural correlates of optical flow patterns in the automated assessment of broiler chicken welfare. *Appl. Anim. Behav. Sci.* **145**, 44–50. (doi:10.1016/j.applanim.2013.02.001)
 20. Kestin S, Knowles T, Tinch A, Gregory N. 1992 Prevalence of leg weakness in broiler chickens and its relationship with genotype. *Vet. Rec.* **131**, 190–194. (doi:10.1136/vr.131.9.190)
 21. Webster A, Fairchild B, Cummings T, Stayer P. 2008 Validation of a three-point gait-scoring system for field assessment of walking ability of commercial broilers. *J. Appl. Poult. Res.* **17**, 529–539. (doi:10.3382/japr.2008-00013)

22. Astill J, Dara RA, Fraser ED, Roberts B, Sharif S. 2020 Smart poultry management: Smart sensors, big data, and the internet of things. *Comput. Electron. Agric.* **170**, 105291. (doi:10.1016/j.compag.2020.105291)
23. LeCun Y, Bengio Y, Hinton G. 2015 Deep learning. *Nature* **521**, 436–444. (doi:10.1038/nature14539)
24. Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M. 2018 DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nat. Neurosci.* **21**, 1281–1289. (doi:10.1038/s41593-018-0209-y)
25. Graving JM, Chae D, Naik H, Li L, Koger B, Costelloe BR, Couzin ID. 2019 DeepPoseKit, a software toolkit for fast and robust animal pose estimation using deep learning. *eLife* **8**, e47994. (doi:10.7554/eLife.47994)
26. Pereira TD, Aldarondo DE, Willmore L, Kislin M, Wang SSH, Murthy M, Shaevitz JW. 2019 Fast animal pose estimation using deep neural networks. *Nat. Methods* **16**, 117–125.
27. Walter T, Couzin ID. 2021 TRex, a fast multi-animal tracking system with markerless identification, and 2D estimation of posture and visual fields. *eLife* **10**, e64000. (doi:10.1038/s41592-018-0234-5)
28. Lauer J, Zhou M, Ye S, Menegas W, Schneider S, Nath T, Rahman MM, Di Santo V, Soberanes D, Feng G et al. 2022 Multi-animal pose estimation, identification and tracking with DeepLabCut. *Nat. Methods* **19**, 496–504. (doi:10.1038/s41592-022-01443-0)
29. Pereira TD, Tabris N, Matsliah A, Turner DM, Li J, Ravindranath S, Papadoyannis ES, Normand E, Deutsch DS, Wang ZY et al. 2022 SLEAP: A deep learning system for multi-animal pose tracking. *Nat. Methods* **19**, 486–495. (doi:10.1038/s41592-022-01426-1)
30. Mathis MW, Mathis A. 2020 Deep learning tools for the measurement of animal behavior in neuroscience. *Curr. Opin. Neurobiol.* **60**, 1–11. (doi:10.1016/j.conb.2019.10.008)
31. Tuia D, Kellenberger B, Beery S, Costelloe BR, Zuffi S, Risse B, Mathis A, Mathis MW, van Langevelde F, Burghardt T et al. 2022 Perspectives in machine learning for wildlife conservation. *Nat. Commun.* **13**, 792. (doi:10.1038/s41467-022-27980-y)
32. Aviagen Group. 2022 ROSS 308/308 FF BROILER: Performance Objectives. https://aviagen.com/assets/Tech_Center/Ross_Broiler/RossxRoss308-BroilerPerformanceObjectives2022-EN.png.
33. Ronco T, Stegger M, Olsen RH, Sekse C, Nordstoga AB, Pohjanvirta T, Lilje B, Lyhs U, Andersen PS, Pedersen K. 2017 Spread of avian pathogenic *Escherichia coli* ST117 O78: H4 in Nordic broiler production. *BMC Genom.* **18**, 1–8. (doi:10.1186/s12864-016-3415-6)
34. Poulsen LL, Bisgaard M, Jørgensen SL, Dideriksen T, Pedersen JR, Christensen H. 2018 Characterization of *Escherichia coli* causing cellulitis in broilers. *Vet. Microbiol.* **225**, 72–78. (doi:10.1016/j.vetmic.2018.09.011)
35. Kromann S, Olsen RH, Bojesen AM, Jensen HE, Thøfner I. 2021 Development of an aerogenous *Escherichia coli* infection model in adult broiler breeders. *Sci. Rep.* **11**, 19556. (doi:10.1038/s41598-021-98270-8)
36. Pors SE, Olsen RH, Christensen JP. 2014 Variations in virulence of avian pathogenic *Escherichia coli* demonstrated by the use of a new in vivo infection model. *Vet. Microbiol.* **170**, 368–374. (doi:10.1016/j.vetmic.2014.02.043)
37. Kromann S, Olsen RH, Bojesen AM, Jensen HE, Thøfner I. 2021 Protective potential of an autogenous vaccine in an aerogenous model of *Escherichia coli* infection in broiler breeders. *Vaccines* **9**, 1233. (doi:10.3390/vaccines9111233)
38. He K, Zhang X, Ren S, Sun J. 2016 Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition* pp. 770–778. (doi:10.1109/CVPR.2016.90)
39. Kingma D, Ba J. 2015 Adam: A Method for Stochastic Optimization. In *Proc. ICLR* San Diego, CA, USA. (doi:10.48550/arXiv.1412.6980)
40. Sturman O, von Ziegler L, Schläppi C, Akyol F, Privitera M, Slominski D, Grimm C, Thieren L, Zerbi V, Grewe B et al. 2020 Deep learning-based behavioral analysis reaches human accuracy and is capable of outperforming commercial solutions. *Neuropsychopharmacology* **45**, 1942–1952. (doi:10.1038/s41386-020-0776-y)

41. Bürkner PC. 2017 brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.* **80**, 1–28. (doi:[10.18637/jss.v080.i01](https://doi.org/10.18637/jss.v080.i01))
42. Dawkins MS, Cain R, Roberts SJ. 2012 Optical flow, flock behaviour and chicken welfare. *Anim. Behav.* **84**, 219–223. (doi:[10.1016/j.anbehav.2012.04.036](https://doi.org/10.1016/j.anbehav.2012.04.036))
43. Teed Z, Deng J. 2020 RAFT: Recurrent all-pairs field transforms for optical flow. In *Proc. ECCV* pp. 402–419. Springer. (doi:[10.48550/arXiv.2003.12039](https://doi.org/10.48550/arXiv.2003.12039))
44. Wang Y, Lipson L, Deng J. 2025 SEA-RAFT: Simple, efficient, accurate RAFT for optical flow. In *Proc. ECCV* pp. 36–54. Springer. (doi:[10.48550/arXiv.2405.14793](https://doi.org/10.48550/arXiv.2405.14793))
45. Horn BK, Schunck BG. 1981 Determining optical flow. *Artif. Intell.* **17**, 185–203. (doi:[10.1016/0004-3702\(81\)90024-2](https://doi.org/10.1016/0004-3702(81)90024-2))
46. Farnebäck G. 2003 Two-frame motion estimation based on polynomial expansion. In *Proc. SCIA* pp. 363–370. Springer. (doi:[10.1007/3-540-45103-X_50](https://doi.org/10.1007/3-540-45103-X_50))
47. Hur J, Roth S. 2020 Optical flow estimation in the deep learning age. *Modelling human motion: from human perception to robot design* pp. 119–140. (doi:[10.1007/978-3-030-46732-6_7](https://doi.org/10.1007/978-3-030-46732-6_7))
48. Paszke A, Gross S, Massa F, Lerer A, Bradbury J, Chanan G, Killeen T, Lin Z, Gimelshein N, Antiga L et al. 2019 PyTorch: An imperative style, high-performance deep learning library. *Proc. NeurIPS* **32**. (doi:[10.48550/arXiv.1912.01703](https://doi.org/10.48550/arXiv.1912.01703))
49. Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, Wieser E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk MH, Brett M, Haldane A, del Río JF, Wiebe M, Peterson P, Gérard-Marchant P, Sheppard K, Reddy T, Weckesser W, Abbasi H, Gohlke C, Oliphant TE. 2020 Array programming with NumPy. *Nature* **585**, 357–362. (doi:[10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2))
50. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J et al. 2020 SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272. (doi:[10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2))
51. Vallat R. 2018 Pingouin: statistics in Python. *J. Open Source Softw.* **3**, 1026. (doi:<https://doi.org/10.21105/joss.01026>)
52. Obata F. 2010 Influence of Escherichia coli shiga toxin on the mammalian central nervous system. *Adv. Appl. Microbiol.* **71**, 1–19. (doi:[10.1016/s0065-2164\(10\)71001-7](https://doi.org/10.1016/s0065-2164(10)71001-7))
53. Hennessey C, Keogh CE, Barboza M, Brust-Mascher I, Knotts TA, Sladek JA, Pusceddu MM, Stokes P, Rabasa G, Honeycutt M et al. 2021 Neonatal enteropathogenic Escherichia coli infection disrupts microbiota-gut-brain axis signaling. *Infect. Immun.* **89**. (doi:[10.1128/iai.00059-21](https://doi.org/10.1128/iai.00059-21))
54. Zhou M, Stoffl L, Mathis MW, Mathis A. 2023 Rethinking pose estimation in crowds: overcoming the detection information bottleneck and ambiguity. In *Proc. ICCV* pp. 14689–14699. (doi:[10.1109/ICCV51070.2023.01350](https://doi.org/10.1109/ICCV51070.2023.01350))
55. Scheidwasser, N. 2025 Deep learning from videography as a tool for measuring E. coli infection in poultry. Zenodo. (doi:[10.5281/zenodo.16795387](https://doi.org/10.5281/zenodo.16795387))

A. Appendix

A.1 Supplementary Figures




Figure S1. brms output for distance travelled, showing density plots for each factor (left) and caterpillar plots (right). Indices: intercept ($b_Intercept$), animal group (between-subject factor; $b_camera_newE.coli$), day (within-subject factor; $sd_Day_Intercept$), dispersion parameter (ϕ).




Figure S2. brms output for the change in body area, showing density plots for each factor (left) and caterpillar plots (right). Indices: intercept (*b_Intercept*), animal group (between-subject factor; *b_camera_newE.coli*), day (within-subject factor; *sd_Day__Intercept*), dispersion parameter (*phi*).




Figure S3. brms output for the time spent near the food source, showing density plots for each factor (left) and caterpillar plots (right). Indices: intercept (*b_Intercept*), animal group (between-subject factor; *b_camera_newE.coli*), day (within-subject factor; *sd_Day__Intercept*), dispersion parameter (*phi*) and zero-inflation parameter (*zi*).




Figure S4. Posterior predictive checks for each model fit. y : observed outcome variable, y_{rep} : simulated datasets from the posterior predictive distribution.

A.2 Supplementary Tables

Table S1. Description of lesion score levels

Organ	Characteristic	Levels
Peritoneum	Inflammatory reaction	0: None 1: Local at the entrance 2: Around the ovary 3: In the omentum 4: All peritoneum
	Amount of exudate	0: None 1: Sparse 2: Some 3: Abundant
	Type of exudate	0: None 1: Aqueous 2: Lump of pus 3: Flakes 4: Confluent
	Transparency of peritoneum	0: Clear 1: Nuclear 2: Cloudy 3: Milky 4: Opaque
Liver	Inflammatory reaction	0: None 1: Fibrin covering < 25% 2: Fibrin covering > 25% 3: Fibrin covering all the liver
Airsac (left)	Transparency of airsac	0: Clear 1: Nuclear 2: Cloudy 3: Milky 4: Opaque
	Amount of exudate	0: None 1: Sparse 2: Some 3: Abundant
	Amount of exudate	0: None 1: Sparse 2: Some 3: Abundant
Lung (left)	Congestion	0: None 1: Sparse (< 25% of lung tissue) 2: Pronounced (> 25% of lung tissue)
	Lung lesions (oedema)	0: None 1: Slight oedema of the alveolar walls 2: Moderate oedematous thickening of alveolar walls with occasional alveoli containing coagulated oedema fluids 3: Extensive occurrence of alveolar and interstitial oedema
	Lung lesions (size)	0: No lesions 1: Small lesions between 1 cm ² and 2 cm ² 2: Moderate-size lesions between 3 cm ² and 4 cm ² 3: Extensive lesions with more than 4 cm ² and covering almost all the area of the lungs
	Granulomas	0: None 1: Few small white caseous/spherical nodules 2: Numerous small white caseous/spherical nodules 3: Many caseous lesions/nodules and green-grey mould due to sporulation
Spleen	Proliferation	0: None 1: Little proliferation 2: Distinct proliferation

Table S2. Mann-Whitney U tests to assess between-group differences in serum amyloid A levels, lesions scores, and number of colony-forming units.

A	B	U-val	<i>p</i> -value	Cohen's <i>d</i>
<i>Serum amyloid A (mg/L)</i>				
Control	<i>E. coli</i>	45.0	2.77e-05	-1.153
<i>Lesion score</i>				
Control	<i>E. coli</i>	39.5	1.064e-05	-2.121
<i>Colony-forming units (per gram of lung)</i>				
Control	<i>E. coli</i>	31.0	1.252e-06	-0.607

Table S3. *brms* summary tables for the behavioural features. Reference group: Control (uninfected). Significant differences (determined if the credible intervals exclude zero) with respect to the negative control are shown in bold. CI: credible interval (l: lower, u: upper). $\hat{R}$: Gelman-Rubin statistic for Markov chain Monte Carlo convergence; convergence is deemed suitable when $\hat{R}$ is close to 1. ESS: effective sample size.

	Estimate	Est.Error	l-95% CI	u-95% CI	$\hat{R}$	Bulk ESS	Tail ESS
<i>Distance travelled</i>							
Intercept	-3.32	0.116	-3.543	-3.079	1.003	2353.594	1771.796
<i>E. coli</i>	-0.286	0.112	-0.506	-0.069	1.002	3717.03	2639.991
<i>Rate of change in body area</i>							
Intercept	-2.912	0.125	-3.148	-2.658	1.001	1750.502	1543.872
<i>E. coli</i>	-0.283	0.117	-0.518	-0.049	1.001	3034.374	2758.279
<i>% Time near food source</i>							
Intercept	-2.281	0.245	-2.74	-1.761	1.002	1333.479	1306.324
<i>E. coli</i>	-0.585	0.148	-0.883	-0.302	1.001	3112.788	2167.955

Table S4. Comparison of different families for mixed-effects modelling using *brms*. ELPD: expected log pointwise predictive density. LOOCV: leave-one-out cross-validation. SE: standard error.

Family (link)	Δ ELPD		LOOCV ELPD	
	Estimate	SE	Estimate	SE
<i>Distance travelled</i>				
Beta (logit)			498.315	19.847
Gaussian (identity)	-153.134	8.685	345.181	19.389
<i>Rate of change in body area</i>				
Beta (logit)			418.771	22.354
Gaussian (identity)	-170.662	10.437	248.109	24.957
<i>% Time spent near food source</i>				
Zero-inflated Beta (logit)			292.026	31.327
Gaussian (identity)	-105.782	30.388	186.244	19.72