

Exploring models

Summary, explainability, and prediction

R.W. Oldford

Modelling

Recall how **J.W. Tukey and M.B. Wilk (1966)** likened analyzing data to conducting experiments.

¹Emphasis added in bold.

Recall how [J.W. Tukey and M.B. Wilk \(1966\)](#) likened analyzing data to conducting experiments.

They suggested that objectives in data analysis that are comparable to those of experimentation are¹

1. “to achieve more specific description of what is loosely known or suspected;
2. “to find unanticipated aspects in the data, and **to suggest unthought-of models** for the data’s **summarization and exposure**;
3. “to employ the data **to assess** the (always incomplete) **adequacy** of a **contemplated model**;
4. “to provide both incentives and guidance for further analysis of the data; and
5. “to keep the investigator usefully stimulated while he absorbs the feeling of his data and considers what to do next.”

¹Emphasis added in bold.

Modelling

Recall how [J.W. Tukey and M.B. Wilk \(1966\)](#) likened analyzing data to conducting experiments.

They suggested that objectives in data analysis that are comparable to those of experimentation are¹

1. “to achieve more specific description of what is loosely known or suspected;
2. “to find unanticipated aspects in the data, and **to suggest unthought-of models** for the data’s **summarization and exposure**;
3. “to employ the data **to assess** the (always incomplete) **adequacy** of a **contemplated model**;
4. “to provide both incentives and guidance for further analysis of the data; and
5. “to keep the investigator usefully stimulated while he absorbs the feeling of his data and considers what to do next.”

Here, model is generally (though not exclusively) to be understood in a more formal mathematical sense.

¹Emphasis added in bold.

Recall how [J.W. Tukey and M.B. Wilk \(1966\)](#) likened analyzing data to conducting experiments.

They suggested that objectives in data analysis that are comparable to those of experimentation are¹

1. “to achieve more specific description of what is loosely known or suspected;
2. “to find unanticipated aspects in the data, and **to suggest unthought-of models** for the data’s **summarization and exposure**;
3. “to employ the data **to assess** the (always incomplete) **adequacy** of a **contemplated model**;
4. “to provide both incentives and guidance for further analysis of the data; and
5. “to keep the investigator usefully stimulated while he absorbs the feeling of his data and considers what to do next.”

Here, model is generally (though not exclusively) to be understood in a more formal mathematical sense. Tukey and Wilk also cautioned against taking them too seriously. In their words, “**Models must be used but must never be believed.**”

¹Emphasis added in bold.

Models – what are they for?

Models serve several purposes.

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity
- ▶ to predict as yet unobserved values of some variates given the values observed on others

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity
- ▶ to predict as yet unobserved values of some variates given the values observed on others

George Box popularized the scientific sense of models with his oft-quoted phrase

“All models are wrong, but some are useful.”

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity
- ▶ to predict as yet unobserved values of some variates given the values observed on others

George Box popularized the scientific sense of models with his oft-quoted phrase

“All models are wrong, but some are useful.”

A less well known quote from Box is:

Since all models are wrong the scientist must be alert to what is importantly wrong.

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity
- ▶ to predict as yet unobserved values of some variates given the values observed on others

George Box popularized the scientific sense of models with his oft-quoted phrase

“All models are wrong, but some are useful.”

A less well known quote from Box is:

Since all models are wrong the scientist must be alert to what is importantly wrong. It is inappropriate to be concerned about mice when there are tigers abroad.

Models – what are they for?

Models serve several purposes. Here are some:

- ▶ to provide a summary pattern of the data, that is those models
 - ▶ contemplated to hold from past experience or prior knowledge
 - ▶ or those only just discovered empirically from the data
- ▶ to help explain how such patterns might have come about
 - ▶ competing models can equally well summarize known or discovered patterns
 - ▶ amongst these, we typically prefer models which are interpretable
 - ▶ amongst these, simple is preferred over complexity
- ▶ to predict as yet unobserved values of some variates given the values observed on others

George Box popularized the scientific sense of models with his oft-quoted phrase

“All models are wrong, but some are useful.”

A less well known quote from Box is:

Since all models are wrong the scientist must be alert to what is importantly wrong. It is inappropriate to be concerned about mice when there are tigers abroad.

G.E.P. Box (1976) “Science and Statistics”, *Journal of the American Statistical Association*, 71, pp. 791-799.

Response models

These are the most common statistical models and come in huge variety of forms.

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s).

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;
- ▶ the explanatory variate values are taken as given (either fixed by design, or conditioned on by choice)

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;
- ▶ the explanatory variate values are taken as given (either fixed by design, or conditioned on by choice) and
- ▶ we try to model the conditional expectation

$$E(Y \mid x_1, \dots, x_p) = \mu(x_1, \dots, x_p)$$

as a function $\mu(\cdot)$ of the explanatory variates x_1, \dots, x_p .

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;
- ▶ the explanatory variate values are taken as given (either fixed by design, or conditioned on by choice) and
- ▶ we try to model the conditional expectation

$$E(Y \mid x_1, \dots, x_p) = \mu(x_1, \dots, x_p)$$

as a function $\mu()$ of the explanatory variates x_1, \dots, x_p .

- ▶ we **fit** the model using observed values of all variates, giving the **estimate** $\hat{\mu}(x_1, \dots, x_p)$ of the **estimand** $\mu(x_1, \dots, x_p)$,

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;
- ▶ the explanatory variate values are taken as given (either fixed by design, or conditioned on by choice) and
- ▶ we try to model the conditional expectation

$$E(Y \mid x_1, \dots, x_p) = \mu(x_1, \dots, x_p)$$

as a function $\mu()$ of the explanatory variates x_1, \dots, x_p .

- ▶ we **fit** the model using observed values of all variates, giving the **estimate** $\hat{\mu}(x_1, \dots, x_p)$ of the **estimand** $\mu(x_1, \dots, x_p)$,
- ▶ we make **inferences** from the model about $\mu()$ using the **estimator** $\tilde{\mu}(x_1, \dots, x_p)$ and its distribution.

Response models

These are the most common statistical models and come in huge variety of forms.

Variates are distinguished by those which are response variates (the y s) and those which are explanatory (the x s). In general, each of x and y could be multivariate (e.g. p explanatory variates x_1, \dots, x_p ; m response variates y_1, \dots, y_m).

Most commonly:

- ▶ the response is univariate y (i.e. $m = 1$) and modelled as a univariate random variable Y ;
- ▶ the explanatory variate values are taken as given (either fixed by design, or conditioned on by choice) and
- ▶ we try to model the conditional expectation

$$E(Y \mid x_1, \dots, x_p) = \mu(x_1, \dots, x_p)$$

as a function $\mu()$ of the explanatory variates x_1, \dots, x_p .

- ▶ we **fit** the model using observed values of all variates, giving the **estimate** $\hat{\mu}(x_1, \dots, x_p)$ of the **estimand** $\mu(x_1, \dots, x_p)$,
- ▶ we make **inferences** from the model about $\mu()$ using the **estimator** $\tilde{\mu}(x_1, \dots, x_p)$ and its distribution.
- ▶ when $\mu()$ is expressed in terms of a finite number of unknown parameters, say $\theta_1, \dots, \theta_k$, we say that it is a **parametric model** with parameter estimates $\hat{\theta}_1, \dots, \hat{\theta}_k$ and corresponding estimators $\tilde{\theta}_1, \dots, \tilde{\theta}_k$.

Response models - examples

Regression models

$$\begin{aligned} Y &= \mu(x_1, \dots, x_q) + R \\ E(R) &= 0 \\ R &\sim F_R(r; \sigma) \end{aligned}$$

with normal regression models having F_R be a normal or Gaussian distribution ($R \sim G(0, \sigma)$).

Regression models

$$\begin{aligned} Y &= \mu(x_1, \dots, x_q) + R \\ E(R) &= 0 \\ R &\sim F_R(r; \sigma) \end{aligned}$$

with normal regression models having F_R be a normal or Gaussian distribution ($R \sim G(0, \sigma)$).

This is also rewritten as

$$\begin{aligned} Y \mid x_1, \dots, x_q &\sim F_Y(y; x_1, \dots, x_q) \\ E(Y \mid x_1, \dots, x_q) &= \mu(x_1, \dots, x_q). \end{aligned}$$

Regression models

$$\begin{aligned} Y &= \mu(x_1, \dots, x_q) + R \\ E(R) &= 0 \\ R &\sim F_R(r; \sigma) \end{aligned}$$

with normal regression models having F_R be a normal or Gaussian distribution ($R \sim G(0, \sigma)$).

This is also rewritten as

$$\begin{aligned} Y \mid x_1, \dots, x_q &\sim F_Y(y; x_1, \dots, x_q) \\ E(Y \mid x_1, \dots, x_q) &= \mu(x_1, \dots, x_q). \end{aligned}$$

The dependency of the mean on the explanatory variates is then usually modelled.

Regression models

$$\begin{aligned} Y &= \mu(x_1, \dots, x_q) + R \\ E(R) &= 0 \\ R &\sim F_R(r; \sigma) \end{aligned}$$

with normal regression models having F_R be a normal or Gaussian distribution ($R \sim G(0, \sigma)$).

This is also rewritten as

$$\begin{aligned} Y \mid x_1, \dots, x_q &\sim F_Y(y; x_1, \dots, x_q) \\ E(Y \mid x_1, \dots, x_q) &= \mu(x_1, \dots, x_q). \end{aligned}$$

The dependency of the mean on the explanatory variates is then usually modelled. Note that this model is **generative** in that it describes how the response values might have been **generated**.

Response models - examples

Regression models

$$\begin{aligned} Y &= \mu(x_1, \dots, x_q) + R \\ E(R) &= 0 \\ R &\sim F_R(r; \sigma) \end{aligned}$$

with normal regression models having F_R be a normal or Gaussian distribution ($R \sim G(0, \sigma)$).

This is also rewritten as

$$\begin{aligned} Y \mid x_1, \dots, x_q &\sim F_Y(y; x_1, \dots, x_q) \\ E(Y \mid x_1, \dots, x_q) &= \mu(x_1, \dots, x_q). \end{aligned}$$

The dependency of the mean on the explanatory variates is then usually modelled. Note that this model is **generative** in that it describes how the response values might have been **generated**.

Such models include the **linear model** whereby

$$\mu(x_1, \dots, x_q) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p.$$

Here linear refers to the mean model being linear in the unknown parameters θ_j .

(There are non-linear regression models as well.)

Generalizing the linear model

A slight generalization is to instead model a function of the conditional mean, as in the so-called **generalized linear model** where now there is a known function $g(\mu)$ called the **link** function and we model

$$g(\mu(x_1, \dots, x_q)) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

with everything else as before.

Generalizing the linear model

A slight generalization is to instead model a function of the conditional mean, as in the so-called **generalized linear model** where now there is a known function $g(\mu)$ called the **link** function and we model

$$g(\mu(x_1, \dots, x_q)) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

with everything else as before.

Another way we might generalize the linear model is to model the mean as

$$\mu(x_1, \dots, x_q) = \theta + h_1(x_1) + \dots + h_p(x_p)$$

where $h_i(x_i)$ are arbitrary functions, each of only a single explanatory variate (x_i). This is called an **additive model** (being additive in functions of the explanatory variates).

Generalizing the linear model

A slight generalization is to instead model a function of the conditional mean, as in the so-called **generalized linear model** where now there is a known function $g(\mu)$ called the **link** function and we model

$$g(\mu(x_1, \dots, x_q)) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

with everything else as before.

Another way we might generalize the linear model is to model the mean as

$$\mu(x_1, \dots, x_q) = \theta + h_1(x_1) + \dots + h_p(x_p)$$

where $h_i(x_i)$ are arbitrary functions, each of only a single explanatory variate (x_i). This is called an **additive model** (being additive in functions of the explanatory variates).

And, if additionally, it is $g(\mu)$ that is modelled additively, the model is called a **generalized additive model**

Generalizing the linear model

A slight generalization is to instead model a function of the conditional mean, as in the so-called **generalized linear model** where now there is a known function $g(\mu)$ called the **link** function and we model

$$g(\mu(x_1, \dots, x_q)) = \theta_0 + \theta_1 x_1 + \dots + \theta_p x_p$$

with everything else as before.

Another way we might generalize the linear model is to model the mean as

$$\mu(x_1, \dots, x_q) = \theta + h_1(x_1) + \dots + h_p(x_p)$$

where $h_i(x_i)$ are arbitrary functions, each of only a single explanatory variate (x_i). This is called an **additive model** (being additive in functions of the explanatory variates).

And, if additionally, it is $g(\mu)$ that is modelled additively, the model is called a **generalized additive model**

These are only a few of the many models that are possible.

Response models - in R a consistent interface

Providers of response models in R try to have a consistent modelling interface.

²Formulas in R generalize [Wilkinson-Rogers](#) notation developed much earlier for specifying linear and generalized linear models.

Response models - in R a consistent interface

Providers of response models in R try to have a consistent modelling interface.

Formulas

Models are typically specified using a standard **formula** representation.²

²Formulas in R generalize [Wilkinson-Rogers](#) notation developed much earlier for specifying linear and generalized linear models.

Response models - in R a consistent interface

Providers of response models in R try to have a consistent modelling interface.

Formulas

Models are typically specified using a standard **formula** representation.² E.g.

$$y \sim x1 + x2 + x3$$

specifies a linear model with y as the response and the variates named x1, x2, and x3 as the explanatory variates (or **predictors**). The variates x1, x2, and x3 are sometimes called the **terms** of the model.

²Formulas in R generalize [Wilkinson-Rogers](#) notation developed much earlier for specifying linear and generalized linear models.

Response models - in R a consistent interface

Providers of response models in R try to have a consistent modelling interface.

Formulas

Models are typically specified using a standard **formula** representation.² E.g.

$$y \sim x1 + x2 + x3$$

specifies a linear model with y as the response and the variates named $x1$, $x2$, and $x3$ as the explanatory variates (or **predictors**). The variates $x1$, $x2$, and $x3$ are sometimes called the **terms** of the model.

On the parameters:

- ▶ linear parameters θ_1 , θ_2 , and θ_3 multiplying $x1$, $x2$, and $x3$ are implicitly associated with each explanatory variate named
- ▶ the intercept term θ_0 is always implicitly assumed to be part of the model; it can be removed by adding a -1 term to the model.
- ▶ that is
 - ▶ $y \sim x1 + x2 + x3$ fits the linear model having conditional mean of Y being

$$\mu = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3$$

²Formulas in R generalize [Wilkinson-Rogers](#) notation developed much earlier for specifying linear and generalized linear models.

Response models - in R a consistent interface

Providers of response models in R try to have a consistent modelling interface.

Formulas

Models are typically specified using a standard **formula** representation.² E.g.

$$y \sim x1 + x2 + x3$$

specifies a linear model with y as the response and the variates named $x1$, $x2$, and $x3$ as the explanatory variates (or **predictors**). The variates $x1$, $x2$, and $x3$ are sometimes called the **terms** of the model.

On the parameters:

- ▶ linear parameters θ_1 , θ_2 , and θ_3 multiplying $x1$, $x2$, and $x3$ are implicitly associated with each explanatory variate named
- ▶ the intercept term θ_0 is always implicitly assumed to be part of the model; it can be removed by adding a -1 term to the model.
- ▶ that is

- ▶ $y \sim x1 + x2 + x3$ fits the linear model having conditional mean of Y being

$$\mu = \theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3$$

and

- ▶ $y \sim x1 + x2 + x3 - 1$ fits the linear model having conditional mean

$$\mu = \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_3.$$

²Formulas in R generalize [Wilkinson-Rogers](#) notation developed much earlier for specifying linear and generalized linear models.

Response models - in R a consistent interface

Formulas

- ▶ terms in the formula can be any specified function of one (or more) explanatory variates named in the data set
- ▶ in some cases (e.g. for generalized additive models), the function expression `s()` is reserved to indicate a **nonparametric smooth** function to be fitted as the additive term.

Formulas

- ▶ terms in the formula can be any specified function of one (or more) explanatory variates named in the data set
- ▶ in some cases (e.g. for generalized additive models), the function expression `s()` is reserved to indicate a **nonparametric smooth** function to be fitted as the additive term. E.g.

$$y \sim x1 + s(x2) + s(x3)$$

specifies that the additive term `x1` enters the model as a usual linear model term, while `s(x2)` and `s(x3)` indicate that the model terms for `x2` and `x3` are to be separate smooth additive functions for each of `x2` and `x3` respectively.

Response models - in R a consistent interface

Formulas

- ▶ terms in the formula can be any specified function of one (or more) explanatory variates named in the data set
- ▶ in some cases (e.g. for generalized additive models), the function expression `s()` is reserved to indicate a **nonparametric smooth** function to be fitted as the additive term. E.g.

$$y \sim x1 + s(x2) + s(x3)$$

specifies that the additive term `x1` enters the model as a usual linear model term, while `s(x2)` and `s(x3)` indicate that the model terms for `x2` and `x3` are to be separate smooth additive functions for each of `x2` and `x3` respectively.

- ▶ terms are joined together with binary operators `+`, `-`, `:`, `*`, and `/`, where for terms `a` and `b` we understand that
 - ▶ `+` `b` indicates adding a separate term `b` to the model,
 - ▶ `-` `b` indicates removing the term `b` from the model,
 - ▶ `a:b` indicates an *interaction* term between `a` and `b` be added,
 - ▶ `a*b` is a short-hand equivalent to `a + b + a:b`,
 - ▶ `a/b` indicates `b` *nested* within `a` and is equivalent to `a + a:b`

Response models - in R a consistent interface

Formulas

- ▶ terms in the formula can be any specified function of one (or more) explanatory variates named in the data set
- ▶ in some cases (e.g. for generalized additive models), the function expression `s()` is reserved to indicate a **nonparametric smooth** function to be fitted as the additive term. E.g.

$$y \sim x1 + s(x2) + s(x3)$$

specifies that the additive term `x1` enters the model as a usual linear model term, while `s(x2)` and `s(x3)` indicate that the model terms for `x2` and `x3` are to be separate smooth additive functions for each of `x2` and `x3` respectively.

- ▶ terms are joined together with binary operators `+`, `-`, `:`, `*`, and `/`, where for terms `a` and `b` we understand that
 - ▶ `a + b` indicates adding a separate term `b` to the model,
 - ▶ `a - b` indicates removing the term `b` from the model,
 - ▶ `a:b` indicates an *interaction* term between `a` and `b` be added,
 - ▶ `a*b` is a short-hand equivalent to `a + b + a:b`,
 - ▶ `a/b` indicates `b` *nested* within `a` and is equivalent to `a + a:b`
- ▶ `poly(x, p)` specifies a polynomial in `x` of degree `p` (uses orthogonal polynomials)

Response models - in R a consistent interface

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model.

Response models - in R a consistent interface

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)`

Response models - in R a consistent interface

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as

Response models - in R a consistent interface

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)`

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality
- ▶ `predict(fit , ...)`

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality
- ▶ `predict(fit , ...)` provide predictions of the response (typically its estimated conditional mean) at any collection of variate values

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality
- ▶ `predict(fit , ...)` provide predictions of the response (typically its estimated conditional mean) at any collection of variate values
 - ▶ requires a data set of new values for every variate named in the model formula

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality
- ▶ `predict(fit , ...)` provide predictions of the response (typically its estimated conditional mean) at any collection of variate values
 - ▶ requires a data set of new values for every variate named in the model formula
 - ▶ often also produces prediction intervals for a new observation and confidence intervals for the conditional mean

Fitted models

Once estimated from the available data, there are common interfaces we expect to have with the fitted model. Suppose the fitted model has been assigned to the variable `myfit`, then common interactions we might expect include

- ▶ `summary(myfit)` should return (and print) a **statistical summary** of the data such as
 - ▶ an overall measure of the quality of the fit
 - ▶ an indication of the statistical significance of each term in the model
- ▶ `plot(fit)` should produce one or more plots that summarize the fit and provide some diagnostic tools for assessing its quality
- ▶ `predict(fit , ...)` provide predictions of the response (typically its estimated conditional mean) at any collection of variate values
 - ▶ requires a data set of new values for every variate named in the model formula
 - ▶ often also produces prediction intervals for a new observation and confidence intervals for the conditional mean
- ▶ `str(fit)` reveals the structure of the fitted model. Here we expect to also find `myfit$residuals` containing the residuals, or deviations, of the observed responses from their fitted conditional mean

Facebook data - fitting linear models

Linear models are fitted in R using the `lm()` function.

```
fit1 <- lm(log10(Impressions) ~ Paid, data = facebook)
summary(fit1)
```

```
##
## Call:
## lm(formula = log10(Impressions) ~ Paid, data = facebook)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.25955 -0.32022 -0.09619  0.28444  2.03001
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)  4.01543     0.02655 151.236 < 2e-16 ***
## Paid         0.21142     0.05031   4.203 3.13e-05 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.5038 on 497 degrees of freedom
## (1 observation deleted due to missingness)
## Multiple R-squared:  0.03432,    Adjusted R-squared:  0.03238
## F-statistic: 17.66 on 1 and 497 DF,  p-value: 3.128e-05
```

Facebook data - contents of linear fits

Extracting contents

```
fit1$coefficients
```

```
## (Intercept)      Paid  
##  4.0154262  0.2114186
```

```
head(model.matrix(fit1))
```

```
## (Intercept) Paid  
## 1          1    0  
## 2          1    0  
## 3          1    0  
## 4          1    1  
## 5          1    0  
## 6          1    0
```

```
head(fit1$residuals)
```

```
##          1          2          3          4          5          6  
## -0.3086231  0.2646283 -0.3746467  0.7175934  0.1179211  0.3036590
```

And prediction (based on the estimated mean don't forget)

```
predict(fit1, newdata = data.frame(Paid = c(0,1)))
```

```
##          1          2  
## 4.015426 4.226845
```

The predicted mean increase in Impressions for paid advertising

```
diff(10^predict(fit1, newdata = data.frame(Paid = c(0,1))))
```

```
##          2  
## 6497.921
```

Facebook data - linear model with a factor

Recall that Category took values Product, Inspiration, Action

```
fit2 <- lm(log10(Impressions) ~ Category, data = facebook)
summary(fit2)
```

```
##
## Call:
## lm(formula = log10(Impressions) ~ Category, data = facebook)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.3727 -0.3074 -0.1079  0.2854  1.9168
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      4.12860    0.03482  118.583  <2e-16 ***
## CategoryInspiration -0.09723    0.05379   -1.807   0.0713 .
## CategoryProduct    -0.09449    0.05672   -1.666   0.0963 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.5105 on 497 degrees of freedom
## Multiple R-squared:  0.008645,    Adjusted R-squared:  0.004655
## F-statistic: 2.167 on 2 and 497 DF,  p-value: 0.1156
```

Facebook data - contents of linear model with a factor

Extracting contents

```
fit2$coefficients
```

```
##          (Intercept) CategoryInspiration    CategoryProduct  
##          4.12860385          -0.09722799          -0.09449137
```

```
head(model.matrix(fit2))
```

```
##   (Intercept) CategoryInspiration CategoryProduct  
## 1           1                0                1  
## 2           1                0                1  
## 3           1                1                0  
## 4           1                0                1  
## 5           1                0                1  
## 6           1                0                1
```

```
head(fit2$residuals)
```

```
##           1           2           3           4           5           6  
## -0.32730938  0.24594205 -0.39059638  0.91032577  0.09923478  0.28497275
```

And prediction on original scale of Impressions

```
10~predict(fit2, newdata = data.frame(Category = factor(levels(facebook$Category))))
```


```
##           1           2           3  
## 13446.33 10749.19 10817.14
```

Conclusions?

Facebook data - other uses of formula

Formulas are also used by other functions (e.g. `boxplot()`)

```
boxplot(log10(Impressions) ~ Category, data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

Formulas are also used by other functions (e.g. `boxplot()`)

```
boxplot(log10(Impressions) ~ Category, data = facebook, col = "lightgrey")
```




Comments? How is this “model” different from the one constructed by `lm()`’?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Paid?

```
boxplot(log10(Impressions) ~ Paid, data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Paid?

```
boxplot(log10(Impressions) ~ Paid, data = facebook, col = "lightgrey")
```




Comments? How is this model formula interpreted?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Type?

```
boxplot(log10(Impressions) ~ Type, data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Type?

```
boxplot(log10(Impressions) ~ Type, data = facebook, col = "lightgrey")
```




Comments? How is this model formula interpreted?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Type?

```
boxplot(log10(like + 1) ~ Type, data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

How about $\log_{10}(\text{Impressions})$ as a function of Type?

```
boxplot(log10(like + 1) ~ Type, data = facebook, col = "lightgrey")
```




Comments? How is this model formula interpreted?

Facebook data - other uses of formula

This works well when explanatory variates are categorical.

```
boxplot(log10(Impressions) ~ Category + Paid, data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

This works well when explanatory variates are categorical.

```
boxplot(log10(Impressions) ~ Category + Paid, data = facebook, col = "lightgrey")
```




Comments?

Note the labels on the horizontal axis.

Facebook data - other uses of formula

This works well when explanatory variates are categorical.

```
boxplot(log10(Impressions) ~ Category + Paid, data = facebook, col = "lightgrey")
```




Comments?

Note the labels on the horizontal axis. How is this model formula interpreted?

Facebook data - other uses of formula

What has changed here?

```
boxplot(log10(Impressions) ~ Paid + Category, data = facebook, col = "lightgrey")
```




Note the labels on the horizontal axis.

Facebook data - other uses of formula

What has changed here?

```
boxplot(log10(Impressions) ~ Paid + Category, data = facebook,  
        col = rep(c("lightgrey", "firebrick"), 3))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Note the labels on the horizontal axis.

Comments?

Facebook data - other uses of formula

By month

```
boxplot(log10(Impressions) ~ Post.Month,  
        xlab = "Season in which post was made",  
        data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

A numeric variate could be made categorical using cut()

```
boxplot(log10(Impressions) ~ cut(Post.Month, 4,  
                                labels = c("Winter", "Spring", "Summer", "Fall")),  
        xlab = "Season in which post was made",  
        data = facebook, col = "lightgrey")
```




Comments?

Facebook data - other uses of formula

How about?

```
boxplot(log10(Impressions) ~ Paid + cut(Post.Month, 4,  
                                         labels = c("Winter", "Spring",  
                                                    "Summer", "Fall")),  
        xlab = "Season in which post was made",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), 4))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

How about?

```
boxplot(log10(Impressions) ~ Paid + cut(Post.Month, 4,  
                                         labels = c("Winter", "Spring",  
                                                    "Summer", "Fall")),  
        xlab = "Season in which post was made",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), 4))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments? Huh? What happened here?

Facebook data - other uses of formula

How about?

```
boxplot(log10(Impressions) ~ Paid + cut(Post.Month, 4,  
                                         labels = c("Winter", "Spring",  
                                                    "Summer", "Fall")),  
        xlab = "Season in which post was made",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), 4))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments? Huh? What happened here? A bug?

Facebook data - other uses of formula

How about?

```
facebook2 <- data.frame(facebook,
                        season = cut(facebook$Post.Month, 4,
                                   labels = c("Winter", "Spring",
                                               "Summer", "Fall")))

boxplot(log10(Impressions) ~ Paid + season,
        xlab = "Season in which post was made",
        data = facebook2, col = rep(c("lightgrey", "firebrick"), 4))
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

How about?

```
facebook2 <- data.frame(facebook,  
                        season = cut(facebook$Post.Month, 4,  
                                   labels = c("Winter", "Spring",  
                                              "Summer", "Fall")))  
  
boxplot(log10(Impressions) ~ Paid + season,  
        xlab = "Season in which post was made",  
        data = facebook2, col = rep(c("lightgrey", "firebrick"), 4))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

Or how about?

```
boxplot(log10(Impressions) ~ Paid + cut(Post.Month, 12,  
                                         labels = month.abb),  
        xlab = "Season in which post was made",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), 6))  
legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

Change response.

```
boxplot(log10(All.interactions + 1) ~ Paid + Category,  
        xlab = "Paid within Category",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), length(facebook$Category)),  
        legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```



Comments?

Facebook data - other uses of formula

Change response. Fitted model

```
fit3 <- lm(log10(All.interactions + 1) ~ Paid + Category, data = facebook)
summary(fit3)
```

```
##
## Call:
## lm(formula = log10(All.interactions + 1) ~ Paid + Category, data = facebook)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.95563 -0.23538  0.01645  0.26075  1.49121
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    1.80436    0.03615  49.907 < 2e-16 ***
## Paid           0.15127    0.04857   3.115  0.00195 **
## CategoryInspiration 0.39403    0.05121   7.695 7.74e-14 ***
## CategoryProduct  0.36251    0.05417   6.692 5.96e-11 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.4859 on 495 degrees of freedom
## (1 observation deleted due to missingness)
## Multiple R-squared:  0.1433, Adjusted R-squared:  0.1382
## F-statistic: 27.61 on 3 and 495 DF,  p-value: < 2.2e-16
```

Comments?

Facebook data - other uses of formula

Change response. A slightly different fitted model

```
fit4 <- lm(log10(All.interactions + 1) ~ Paid * Category, data = facebook)
summary(fit4)
```

```
##
## Call:
## lm(formula = log10(All.interactions + 1) ~ Paid * Category, data = facebook)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -2.01793 -0.23100  0.02586  0.27246  1.51761
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)    1.77795    0.03941  45.111 < 2e-16 ***
## Paid           0.23998    0.07224   3.322  0.00096 ***
## CategoryInspiration 0.46570    0.06040   7.711 6.96e-14 ***
## CategoryProduct  0.37776    0.06302   5.994 3.95e-09 ***
## Paid:CategoryInspiration -0.25185    0.11299  -2.229  0.02627 *
## Paid:CategoryProduct  -0.04374    0.12234  -0.358  0.72083
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.4843 on 493 degrees of freedom
## (1 observation deleted due to missingness)
## Multiple R-squared:  0.1524, Adjusted R-squared:  0.1438
## F-statistic: 17.72 on 5 and 493 DF,  p-value: 3.639e-16
```

Comments?

Facebook data - other uses of formula

Change response. A slightly different fitted model

```
fit4$coefficients
```

##	(Intercept)	Paid	CategoryInspiration
##	1.77795481	0.23997989	0.46569543
##	CategoryProduct	Paid:CategoryInspiration	Paid:CategoryProduct
##	0.37776394	-0.25185230	-0.04374152

```
head(model.matrix(fit4))
```

##	(Intercept)	Paid	CategoryInspiration	CategoryProduct	Paid:CategoryInspiration
## 1	1	0	0	1	0
## 2	1	0	0	1	0
## 3	1	0	1	0	0
## 4	1	1	0	1	0
## 5	1	0	0	1	0
## 6	1	0	0	1	0
##	Paid:CategoryProduct				
## 1	0				
## 2	0				
## 3	0				
## 4	1				
## 5	0				
## 6	0				

Comments?

Facebook data - other uses of formula

Change explanatory variates to two factors

```
fit5 <- lm(log10(All.interactions + 1) ~ Type * Category, data = facebook)
summary(fit5)
```

```
##
## Call:
## lm(formula = log10(All.interactions + 1) ~ Type * Category, data = facebook)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -1.83783 -0.23716  0.01508  0.25688  1.60199
##
## Coefficients: (2 not defined because of singularities)
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept)      1.73278    0.10895   15.904 <2e-16 ***
## TypePhoto         0.10504    0.11469    0.916  0.3602
## TypeStatus        0.36213    0.30167    1.200  0.2306
## TypeVideo         0.65020    0.21397    3.039  0.0025 **
## CategoryInspiration  0.20172    0.49927    0.404  0.6864
## CategoryProduct   -0.03381    0.49927   -0.068  0.9460
## TypePhoto:CategoryInspiration  0.20383    0.50213    0.406  0.6850
## TypeStatus:CategoryInspiration -0.09280    0.62270   -0.149  0.8816
## TypeVideo:CategoryInspiration      NA         NA      NA      NA
## TypePhoto:CategoryProduct    0.39575    0.50315    0.787  0.4319
## TypeStatus:CategoryProduct    0.16441    0.57849    0.284  0.7764
## TypeVideo:CategoryProduct      NA         NA      NA      NA
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 0.4872 on 490 degrees of freedom
## Multiple R-squared:  0.1473, Adjusted R-squared:  0.1316
## F-statistic: 9.405 on 9 and 490 DF, p-value: 3.024e-13
```

Comments?

Facebook data - other uses of formula

Just the coefficients.

```
fit5$coefficients
```

```
##              (Intercept)              TypePhoto
##              1.73278345              0.10504178
##              TypeStatus              TypeVideo
##              0.36213204              0.65020444
##              CategoryInspiration          CategoryProduct
##              0.20171500              -0.03381344
## TypePhoto:CategoryInspiration TypeStatus:CategoryInspiration
##              0.20382942              -0.09279854
## TypeVideo:CategoryInspiration          TypePhoto:CategoryProduct
##              NA              0.39574802
##              TypeStatus:CategoryProduct          TypeVideo:CategoryProduct
##              0.16440892              NA
```


Facebook data - other uses of formula

And now the corresponding model matrix.

```
head(model.matrix(fit5))
```


```
## (Intercept) TypePhoto TypeStatus TypeVideo CategoryInspiration
## 1          1          1          0          0                0
## 2          1          0          1          0                0
## 3          1          1          0          0                1
## 4          1          1          0          0                0
## 5          1          1          0          0                0
## 6          1          0          1          0                0
## CategoryProduct TypePhoto:CategoryInspiration TypeStatus:CategoryInspiration
## 1              1                          0                          0
## 2              1                          0                          0
## 3              0                          1                          0
## 4              1                          0                          0
## 5              1                          0                          0
## 6              1                          0                          0
## TypeVideo:CategoryInspiration TypePhoto:CategoryProduct
## 1                          0                          1
## 2                          0                          0
## 3                          0                          0
## 4                          0                          1
## 5                          0                          1
## 6                          0                          0
## TypeStatus:CategoryProduct TypeVideo:CategoryProduct
## 1              0              0
## 2              1              0
## 3              0              0
## 4              0              0
## 5              0              0
## 6              1              0
```

What does the intercept term represent?

Facebook data - other uses of formula

Or how about?

```
boxplot(log10(All.interactions + 1) ~ Paid + Type + Category,  
        xlab = "Combination",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), length(facebook$Type) * length(facebook$Paid)),  
        legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

Or

```
boxplot(log10(All.interactions + 1) ~ Paid + Category + cut(Post.Month, 12, labels = month.abb),  
        xlab = "Paid within Category within Month",  
        data = facebook, col = rep(c("lightgrey", "firebrick"), 12 * length(facebook$Category)),  
        legend("topright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

Facebook data - other uses of formula

Focus only on Fall

```
Fall <- facebook$Post.Month %in% 9:12
with(facebook[Fall,], {
  boxplot(log10(All.interactions + 1) ~ Paid + cut(Post.Month, 4, labels = month.abb[9:12]) + Category,
    xlab = "Paid within Month within Category", main = "Fall only",
    col = rep(c("lightgrey", "firebrick"), 4 * length(Category)))
  legend("bottomright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
})
```




Comments?

What about a continuous explanatory variate?

Could use `cut()` on the continuous (ratio-scaled) variate to turn it into a categorical and proceed as before.

For example equal width intervals:

```
boxplot(log10(like + 1) ~ Paid + cut(log10(Impressions), 4),  
        xlab = "Paid within log10(Impressions)",  
        data = facebook, main = "Equal width intervals",  
        col = rep(c("lightgrey", "firebrick"), 2))  
legend("bottomright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```




Comments?

What about a continuous explanatory variate?

Or perhaps four intervals of equal numbers

```
boxplot(log10(like + 1) ~ Paid + cut(log10(Impressions),
                                     breaks = quantile(log10(Impressions))),
        xlab = "Paid within log10(Impressions)",
        data = facebook, main = "Equal count intervals",
        col = rep(c("lightgrey", "firebrick"), 2))
legend("bottomright", legend = c("free", "paid"), fill = c("lightgrey", "firebrick"))
```



Comments?

What about a continuous explanatory variate?

Alternatively, we could build a (perhaps complicated) linear model, say modelling the mean of response Y as a polynomial of explanatory variate x :

$$\mu(x) = \beta_0 + \beta_1 x + \beta_2 x^2 + \dots + \beta_p x^p$$

(or as any other linear (in the coefficients) model).

What about a continuous explanatory variate?

Alternatively, we could build a (perhaps complicated) linear model, say modelling the mean of response Y as a polynomial of explanatory variate x :

$$\mu(x) = \beta_0 + \beta_1 x + \beta_2 x^2 + \dots + \beta_p x^p$$

(or as any other linear (in the coefficients) model).

Such models can be fitted by least-squares.

What about a continuous explanatory variate?

Alternatively, we could build a (perhaps complicated) linear model, say modelling the mean of response Y as a polynomial of explanatory variate x :

$$\mu(x) = \beta_0 + \beta_1 x + \beta_2 x^2 + \dots + \beta_p x^p$$

(or as any other linear (in the coefficients) model).

Such models can be fitted by least-squares.

Unfortunately, these models require a parametric form (e.g. a polynomial) be specified that will fit the data **everywhere** (i.e. globally for all x).

Alternatively, we could try fitting many simple functions of x **locally**, different at every value of x . Connecting the fitted values together produces an estimated $\mu(x)$

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x .

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x_i))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$.

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x_i))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$. We fit $\mu(x)$ for every x on in the range of the data.

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x_i))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$. We fit $\mu(x)$ for every x on in the range of the data.

We could also make the weight function $w_i(x)$ to be 1 for those x_i near x and 0 for those far away. In this way, the weights determine the x_i values that contribute to fitting $\mu(x)$ and those which do not.

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$. We fit $\mu(x)$ for every x on in the range of the data.

We could also make the weight function $w_i(x)$ to be 1 for those x_i near x and 0 for those far away. In this way, the weights determine the x_i values that contribute to fitting $\mu(x)$ and those which do not.

For example, for the i th observation x_i when fitting at any point x we could have

$$w_i(x) = K \left(\frac{x_i - x}{a} \right)$$

for some $a > 0$ and function $K(t) \geq 0$ that is maximal at ($t = 0$) and decreasing with $|t|$.

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$. We fit $\mu(x)$ for every x on in the range of the data.

We could also make the weight function $w_i(x)$ to be 1 for those x_i near x and 0 for those far away. In this way, the weights determine the x_i values that contribute to fitting $\mu(x)$ and those which do not.

For example, for the i th observation x_i when fitting at any point x we could have

$$w_i(x) = K \left(\frac{x_i - x}{a} \right)$$

for some $a > 0$ and function $K(t) \geq 0$ that is maximal at $(t = 0)$ and decreasing with $|t|$.

The constant a controls the size of the region (i.e. which x_i s) that will be used to fit at x .

What about a continuous explanatory variate?

For example, while we might not be willing to have one line fit all points, we might be willing to have *different lines* fitted in separate (and contiguous) regions of x . That is we could fit lines **locally** within each region of x .

We can fit locally by using **weighted least squares** which minimizes

$$\sum_{i=1}^n w_i(x) (y_i - \mu(x))^2$$

where $w_i(x)$ depends on the location x where we are fitting $\mu(x)$. We fit $\mu(x)$ for every x on in the range of the data.

We could also make the weight function $w_i(x)$ to be 1 for those x_i near x and 0 for those far away. In this way, the weights determine the x_i values that contribute to fitting $\mu(x)$ and those which do not.

For example, for the i th observation x_i when fitting at any point x we could have

$$w_i(x) = K \left(\frac{x_i - x}{a} \right)$$

for some $a > 0$ and function $K(t) \geq 0$ that is maximal at $(t = 0)$ and decreasing with $|t|$.

The constant a controls the size of the region (i.e. which x_i s) that will be used to fit at x . The smaller these regions were, the less structure we would be imposing on the underlying function $\mu(x)$.

Locally weighted sum of squares fitting – loess()

In R there is a function called `loess` that fits a locally weighted sum of squares estimate that pays a little more attention to some of these problems.

```
loess(formula, data, ..., span = 0.75, ...)
```


Locally weighted sum of squares fitting – *loess()*

In R there is a function called `loess` that fits a locally weighted sum of squares estimate that pays a little more attention to some of these problems.

```
loess(formula, data, ..., span = 0.75, ...)
```

As its default weight function `loess` uses Tukey's tri-cube weight

$$K(t) = \begin{cases} (1 - |t|^3)^3 & t \in [-1, 1] \\ 0 & \text{otherwise.} \end{cases}$$




Locally weighted sum of squares fitting – *loess()*

In R there is a function called `loess` that fits a locally weighted sum of squares estimate that pays a little more attention to some of these problems.

```
loess(formula, data, ..., span = 0.75, ...)
```

As its default weight function `loess` uses Tukey's tri-cube weight

$$K(t) = \begin{cases} (1 - |t|^3)^3 & t \in [-1, 1] \\ 0 & \text{otherwise.} \end{cases}$$




This is generally more resistant to outlying x_i s than, say, a Gaussian weight function (i.e. $K(W) = \phi(w)$ the $N(0,1)$ density).

N.B. `loess` is not restricted to fitting local lines. It can fit any degree polynomial locally (though typically only degree 1 or 2 is used in practice).

Locally weighted sum of squares fitting – loess()

```
fit <- loess(log10(like + 1) ~ log10(Impressions), data=facebook)
plot(log10(facebook$Impressions), log10(facebook$like + 1),
     col="grey50", pch=19, main = "Loess smooth")
# Get the values predicted by the loess
xmin <- min(facebook$Impressions)
xmax <- max(facebook$Impressions)
x <- seq(xmin, xmax, length.out = 200)
pred <- predict(fit, newdata = data.frame(Impressions = x))
lines(log10(x),pred, lwd=3, col="steelblue")
```



Locally weighted sum of squares fitting – loess()

```
fit2 <- loess(log10(like + 1) ~ log10(Impressions), data=facebook , span = 0.3)
plot(log10(facebook$Impressions), log10(facebook$like + 1),
     col="grey50", pch=19, main = "Loess smooth")
# Get the values predicted by the loess
xmin <- min(facebook$Impressions)
xmax <- max(facebook$Impressions)
x <- seq(xmin, xmax, length.out = 200)
pred <- predict(fit2, newdata = data.frame(Impressions = x))
lines(log10(x),pred, lwd=3, col="steelblue")
```

