

Binomial random variables

56 marks

Suppose $X \sim \text{Binomial}(n, p)$, then

$$Pr(X = x) = \binom{n}{x} p^x (1-p)^{n-x} \quad x = 0, 1, \dots, n.$$

from which it follows that $E(X) = np$ and $Var(X) = np(1-p)$.

Once x is observed, the unknown proportion can be estimated (e.g. via maximum likelihood) as the numerical value of $\hat{p} = x/n$; the corresponding random estimator would be $\tilde{p} = X/n$.

In this question, you are going to develop your intuition about simple proportions as random variables through a bit of mathematics, some introductory R programming, and a little simulation and visualization.

- a. The **odds** (in favour of the event) are defined by the ratio $p/(1-p)$ (e.g. even odds are 1, or 1:1, when $p = 1/2$; odds are 9:1 in favour when $p = 9/10$; odds less than one are often inverted to be described as the odds *against* the event).

Suppose we are interested in comparing the binomial probabilities $Pr(X = a)$ to $Pr(X = b)$. For example, the ratio, $Pr(X = a)/Pr(X = b)$, tells us how many times more (or less) likely it is to observe $x = a$ than $x = b$.

- i. (1 mark) Express the ratio $Pr(X = a)/Pr(X = b)$ as a function of the odds.

$$\frac{Pr(X = a)}{Pr(X = b)} = \frac{\binom{n}{a} p^a (1-p)^{n-a}}{\binom{n}{b} p^b (1-p)^{n-b}} = \frac{\binom{n}{a}}{\binom{n}{b}} \frac{p^{a-b}}{(1-p)^{-n+a+n-b}} = \frac{\binom{n}{a}}{\binom{n}{b}} \frac{p^{a-b}}{(1-p)^{a-b}} = \frac{\binom{n}{a}}{\binom{n}{b}} \text{odds}^{a-b}$$

- ii. (4 marks) Here, in two different ways, you will write a function of the odds which calculates $Pr(X = a)/Pr(X = b)$ for a given n .

For the first way, write `prob_ratio1()` to do the calculation using the function `choose()`:

```
prob_ratio1 <- function (n, a, b, odds = 1) {  
  choose_n_a <- choose(n,a)  
  choose_n_b <- choose(n,b)  
  ans <- choose_n_a/choose_n_b*odds^(a-b)  
  return(ans)  
}  
# print(prob_ratio1(5,3,4,5))
```

For the second way, write ``prob_ratio1()`` to do the calculation using the function ``dbinom()``:

```
prob_ratio2 <- function (n, a, b, odds = 1) {  
  p <- odds/(1+odds)  
  p_a <- dbinom(a,size=n,prob=p)  
  p_b <- dbinom(b,size=n,prob=p)  
  return(p_a/p_b)  
}  
# print(prob_ratio2(5,3,4,5))
```

Both are calculating the same values.

iii. **(2 marks)** Report the following values for ``prob_ratio1()`` and ``prob_ratio2()``

```
# using choose()
prob_ratio1(50, a = 5, b = 45)

## [1] 1

prob_ratio1(50, a = 5, b = 45, odds = 9)

## [1] 6.765496e-39

# and using dbinom()
prob_ratio2(50, a = 5, b = 45)

## [1] 1

prob_ratio2(50, a = 5, b = 45, odds = 9)

## [1] 6.765496e-39
```

b. Extreme proportions like $\hat{p} \approx 0$ or $\hat{p} \approx 1$ often generate a great deal of interest in an analysis.

For example, p might be the proportion of people in some population who perhaps die from some exposure to some toxin, or are cured of a disease by some treatment, or maybe just say they would vote for a particular party or candidate. In any of these cases it can be surprising (even alarming) to see either $\hat{p} \approx 0$ or $\hat{p} \approx 1$, so much so that some explanation seems in order.

Suppose we have observed x from $Binomial(n, p)$ and y from $Binomial(m, p)$ – that is the same probability of occurrence but different sample sizes. Denote the observed proportions as $\hat{p}_x = x/n$ and $\hat{p}_y = y/m$, respectively.

i. *(2 marks)* Give the mathematical expression for the ratio

$$\frac{Pr(\tilde{p}_x = 0)}{Pr(\tilde{p}_y = 0)}$$

and for the ratio

$$\frac{Pr(\tilde{p}_x = 1)}{Pr(\tilde{p}_y = 1)}$$

$$\frac{Pr(\tilde{p}_x = 0)}{Pr(\tilde{p}_y = 0)} = \frac{\binom{n}{0}p^0(1-p)^n}{\binom{m}{0}p^0(1-p)^m} = \frac{\binom{n}{0}}{\binom{m}{0}}(1-p)^{n-m} = (1-p)^{n-m}$$

$$\frac{Pr(\tilde{p}_x = 1)}{Pr(\tilde{p}_y = 1)} = \frac{\binom{n}{1}p^1(1-p)^0}{\binom{m}{1}p^1(1-p)^0} = \frac{\binom{n}{1}}{\binom{m}{1}}p^{n-m} = p^{n-m}$$


ii. *(2 marks)* Describe what happens to each of these ratios for $n < m$ and m increases. As m increases, the first ratio will increase. The base $1-p$ is between 0 and 1 and the exponent decreases from 0. The power will increase if m increases. As m increases, the second ratio will increase. The base p is between 0 and 1 and the exponent decreases from 0. The power will increase if m increases.

iii. *(3 marks)* Suppose $p = 1/2$, $n = 5$ and $m \in \{5, 6, 7, \dots, 14, 15\}$. Using the `plot()` function (with appropriate title and axis labels), plot the curve of the pair (n, m) for all values.

```

ms <- seq(from=5, to=15)
get_ratio <- function(p,n,ms){
  ans = c()
  for (i in 1:length(ms)){
    ans <- c(ans, p^(n-ms[i]))
  }
  return(ans)
}
ratios <- get_ratio(1/2,5,ms)
plot(ms,ratios,xlab="value of m",ylab="ratio",col="blue")

```



iv. **(2 marks)** In plain English, express when \widehat{p} is most likely to be 1 and when you should

c. For the binomial definition, as given above:

i. *(2 marks)* Mathematically derive $E(\tilde{p})$ and the **standard deviation** $SD(\tilde{p})$.

ii. *(2 marks)* Write a function that calculates the standard deviation of \tilde{p} for any pair of values of n and p as follows:

```

sd_p_wig <- function (n, p) {
  # Your code here
}

```

d. Chebyshev's inequality relates the nearness of a random variable Y to its expectation μ as a function of its standard deviation σ (provided both exist and are finite) as follows:

$$Pr(|Y - \mu| \geq k\sigma) \leq \frac{1}{k^2}$$

for any constant $k > 1$.

i. *(2 marks)* Write this inequality when $Y = \tilde{p}$, the binomial proportion estimator.

- ii. (4 marks) Suppose that $k = 5$. As a function of p , mathematically express the sample size n needed to ensure by Chebyshev's inequality that for our estimator \tilde{p} we have

$$Pr\left(|\tilde{p} - p| \geq \frac{1}{50}\right) \leq \frac{1}{25}.$$

Using the function `plot()`, with appropriate title and x and y axis labels, plot the curve of n as a function of p for `p = seq(0,1, 0.01)`.

- iii. (4 marks) Suppose that $k = 5$, and $n = 2500$. As a function of p , mathematically express the bound B given by Chebyshev's inequality, so that for our estimator \tilde{p} we have

$$Pr(|\tilde{p} - p| \geq B) \leq \frac{1}{25}.$$

Using the function `plot()`, with appropriate title and x and y axis labels, plot the curve of B as a function of p for `p = seq(0,1, 0.01)`.

- iv (2 marks) In simple English, summarize what the largest B says about how well p is likely to be estimated when $n = 2500$ according to Chebyshev's inequality.
- e. The functions `plot()`, `lines()`, and `abline()` can be used to plot some data, to add curves, and to add a straight line to a plot. To learn more, type `help("plot.default")`, `help("lines")`, and `help("abline")`.

Use these plotting functions, and appropriate arguments (including meaningful titles, and axis labels), to produce the following plots of the function `sd_p_wig()`:

- i. (3 marks) As a function of `p = seq(0, 1, 0.01)` for a fixed `n = 10`. Add a vertical **dashed** line in **red** at the value of p which maximizes the standard deviation.

Hand in your code and plot.

- ii. (4 marks) As a function of `n ∈ {5, 10, 15, ..., 50}` for the different values of `p ∈ {0.1, 0.3, 0.5, 0.8}`.

- Use a different colour and line type for each curve (i.e. value of `p`).
- Use `lwd = 2` for all curves.
- Use `legend()` to add a legend to the “topright” corner of the plot, identify each curve by its colour and line type.
- Hand in your code and plot.

- iii. (2 marks) Comment on your findings about the dependency of the standard deviation of the binomial proportion estimator \tilde{p} as a function of n and p .

- f. The R function `rbinom()` can be used to generate pseudo-random values `x` from a binomial distribution. In this question, you will examine the samples of the sample **proportions**, x/n , drawn from binomials with the same value of p but different values of n .

- i. (3 marks) Using `rbinom()`, for every $n \in \{5, 10, 15, \dots, 45, 50\}$ generate 100 proportions, each one based on an independent pseudo-random value from $\text{Binomial}(n, 0.5)$.

You might find the `rep()` function useful in constructing `n`.

Show your code.

- ii. (3 marks) Plot the pairs (n, \hat{p}) as points (use plot arguments `ylim = c(0,1)`, `pch = 19`, `col = adjustcolor("steelblue", 0.2)`, `cex = 0.5`).

Note that there will be 100 proportions for every n .

Add a red dashed horizontal line at $p = 0.5$.

- iii. (2 marks) Repeat the production of the above plot, complete with horizontal line, but instead of using `n` as the x variable in the plot, use `jitter(n, 2)`.
Show your code and resulting plot.
Comment on the effect of `jitter()`
- iv. (1 mark) Based on the either of the above plots, what do you conclude about the distribution of binomial proportions as n increases?
- g. (6 marks) With supporting reference to any/all suitable discoveries you have made in the above questions, comment on each of the following:
 - i. Which values for the true binomial probability p are hardest/easiest to estimate from a sample. Why?
 - ii. Law of large numbers? What can you say about the effect of increasing sample size n on the quality and/or interpretation of your estimate \hat{p} of p ?
 - iii. Law of small numbers? What can you say about the effect of decreasing sample size n on the quality and/or interpretation of your estimate \hat{p} of p ?