

ՀՀ ԿԳՄՍ ՆԱԽԱՐԱՐՈՒԹՅՈՒՆ
ՀԱՅԱՍՏԱՆԻ ԵՎՐՈՊԱԿԱՆ ՀԱՄԱԼՍԱՐԱՆ
ՏԵՂԵԿԱՏՎԱԿԱՆ ՀԱՄԱԿԱՐԳԵՐ

ԱՆՀԱՏԱԿԱՆ ԱՇԽԱՏԱՆՔ

Գծային հանրահաշիվ՝ տարբերակ 19

ԱՌԱՐԿԱ՝ ԳԾԱՅԻՆ ՀԱՆՐԱՅԱՇԻՎ
ԿՈՒՐՍ՝ ԱՌԱՋԻՆ
ՈՒՍԱՆՈՂ՝ ՍԵՐՈՔ ՏԻԳՐԱՆՅԱՆ
ԴԱՍԱԽՈՍ՝ Հ. ՂՈՒՄԱՇՅԱՆ

ԵՐԵՎԱՆ 2024

Linear Algebra

Serob Tigranyan

May 22, 2025

Contents

1 Task 1	2
1.1 Linear Independence	2
1.2 Vector expressed in given basis	2
2 Task 2	4
2.1 Orthogonality	4
2.1.1 Orthogonal Basis Vectors	4
3 Task 3	7
4 Task 4	9

1 Task 1

Given the following basis vectors:

$$e_1 = (1, 1, 3)$$

$$e_2 = (1, 1, 0)$$

$$e_3 = (1, 2, 0)$$

And vector x :

$$x = (5, 4, 6)$$

1.1 Linear Independence

We need to check if the given basis vectors are valid:

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 2 \\ 3 & 0 & 0 \end{bmatrix}$$

Determine the Rank by first finding the determinant:

$$D(A) = 0 + 6 + 0 - 3 - 0 - 0 = 3$$

Since our determinant is non-zero, then we have a full rank which is 3 which further means that the basis vectors are linearly independent.

$$\text{rank}(A) = 3$$

1.2 Vector expressed in given basis

We need to express vector x in our basis e_1, e_2, e_3 .

$$x = \lambda_1 e_1 + \lambda_2 e_2 + \lambda_3 e_3$$

Therefore we have:

$$(5, 4, 6) = \lambda(1, 1, 3) + \lambda(1, 1, 0) + \lambda(1, 2, 0)$$

$$(5, 4, 6) = (\lambda_1 + \lambda_2 + \lambda_3; \lambda_1 + \lambda_2 + 2\lambda_3; 3\lambda_1)$$

We form a linear system of equations:

$$\begin{cases} \lambda_1 + \lambda_2 + \lambda_3 = 5 \\ \lambda_1 + \lambda_2 + 2\lambda_3 = 4 \\ 3\lambda_1 = 6 \end{cases}$$

Now we solve the system:

$$\begin{cases} \lambda_1 + \lambda_2 + \lambda_3 = 5 \\ \lambda_1 + \lambda_2 + 2\lambda_3 = 4 \\ \lambda_1 = 2 \end{cases}$$

$$\begin{cases} \lambda_2 = 3 - \lambda_3 \\ 2 + (3 - \lambda_3) + 2\lambda_3 = 4 \\ \lambda_1 = 2 \end{cases}$$

$$\begin{cases} \lambda_2 = 3 - \lambda_3 \\ \lambda_3 = -1 \\ \lambda_1 = 2 \end{cases}$$

$$\begin{cases} \lambda_2 = 4 \\ \lambda_3 = -1 \\ \lambda_1 = 2 \end{cases}$$

Therefore our vector x expressed in our basis vectors e_1, e_2, e_3 is:

$$x_e = (2, 4, -1)$$

2 Task 2

Given the vectors a_1 and a_2 we need to find if they're orthogonal and if so, then find their orthogonal basis vectors:

$$a_1 = (4, 4, 1, -1)$$

$$a_2 = (1, -1, 1, 1)$$

2.1 Orthogonality

To check if these two are orthogonal we perform scalar multiplication on both of them:

$$(a_1 \cdot a_2) = 4 - 4 + 1 - 1 = 0 \Rightarrow a_1 \perp a_2$$

Our vectors are indeed orthogonal, now we need to find their orthogonal basis vectors.

2.1.1 Orthogonal Basis Vectors

To find the orthogonal basis vectors we pick arbitrary vector:

$$\gamma = (x, y, z, w)$$

So that:

$$\gamma \perp a_1$$

$$\gamma \perp a_2$$

We form a linear system of equations from this like so:

$$\begin{cases} (\gamma \cdot a_1) = 0 \\ (\gamma \cdot a_2) = 0 \end{cases} \Rightarrow \begin{cases} 4x + 4y + z - w = 0 \\ x - y + z + w = 0 \end{cases}$$

Now we try and solve the system:

$$\begin{cases} 4x + 4y + z - w = 0 \\ x - y + z + w = 0 \end{cases}$$

To solve the system we'll need to find the rank:

$$A = \begin{bmatrix} 4 & 4 & 1 & -1 \\ 1 & -1 & 1 & 1 \end{bmatrix}$$

Find the deltas of matrix A :

$$\Delta_1 = 4 \neq 0$$

$$\Delta_2 = \begin{bmatrix} 4 & 4 \\ 1 & -1 \end{bmatrix} = -4 - 4 = -8 \neq 0 \Rightarrow \text{rank}(A) = 2$$

Now we need the identity matrix based on our rank:

$$I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Now we solve our system based on the identity matrix.

Because our identity matrix is 2×2 , we move two elements from the left to right:

$$\begin{cases} 4x + 4y = -z + w \\ x - y = -z - w \end{cases}$$

Now solve by placing the first row elements to be on the right side of the equations:

$$\begin{cases} 4x + 4y = -1 \\ x - y = -1 \end{cases}$$

$$\begin{cases} 4x + 4y = -1 \\ x = -1 + y \end{cases}$$

$$\begin{cases} y = \frac{3}{8} \\ x = -1 + y \end{cases}$$

$$\begin{cases} y = \frac{3}{8} \\ x = -\frac{5}{8} \end{cases}$$

Now the second row of elements to be on the right side of the equations:

$$\begin{cases} 4x + 4y = 1 \\ x - y = -1 \end{cases} \Rightarrow \begin{cases} y = \frac{5}{8} \\ x = -\frac{3}{8} \end{cases}$$

Now we have to basis vectors γ_1 and γ_2 :

$$\gamma_1 = \left(-\frac{5}{8}; \frac{3}{8}; 1; 0 \right)$$

$$\gamma_2 = \left(-\frac{3}{8}; \frac{5}{8}; 0; 1\right)$$

Now we check if they're orthogonal:

$$(\gamma_1 \cdot \gamma_2) = \left(\frac{5}{8} \cdot \frac{3}{8} + \frac{3}{8} \cdot \frac{5}{8} + 0 + 0\right) = \frac{15}{32} \neq 0$$

Unfortunately our vectors are not orthogonal therefore we must find the fourth β .

We do this by picking any of the previous vectors to serve as a_3 , in this case $a_3 = \gamma_1$:

$$\beta = (x, y, z, w) \neq 0$$

$$\beta \perp a_1, \beta \perp a_2, \beta \perp a_3$$

Form a linear system of equations:

$$\begin{cases} (\beta \cdot a_1) = 0 \\ (\beta \cdot a_2) = 0 \\ (\beta \cdot a_3) = 0 \end{cases} \Rightarrow \begin{cases} 4x + 4y + z - w = 0 \\ x - y + z + w = 0 \\ -\frac{5x}{8} + \frac{3y}{8} + z = 0 \end{cases}$$

To solve we'll need to turn this into a matrix and find the rank again:

$$A = \begin{bmatrix} 4 & 4 & 1 & -1 \\ 1 & -1 & 1 & 1 \\ -\frac{5}{8} & \frac{3}{8} & 1 & 0 \end{bmatrix}$$

Find the rank:

$$\Delta_3 = \begin{bmatrix} 4 & 4 & 1 \\ 1 & -1 & 1 \\ -\frac{5}{8} & \frac{3}{8} & 1 \end{bmatrix} = -\frac{49}{4} \neq 0 \Rightarrow \text{rank}(A) = 3$$

Now form a linear system of equations again:

$$\begin{cases} 4x + 4y + z = w \\ x - y + z = -w \\ -\frac{5}{8}x + \frac{3}{8}y + z = 0 \end{cases}$$

We can solve this system of equations using Cramer's Rule:

$$\begin{bmatrix} 4 & 4 & 1 \\ 1 & -1 & 1 \\ -\frac{5}{8} & \frac{3}{8} & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$D(A) = -\frac{49}{4}$$

$$x = \left| \begin{bmatrix} 1 & 4 & 1 \\ -1 & -1 & 1 \\ 0 & \frac{3}{8} & 1 \end{bmatrix} \right| = \frac{9}{4}$$

$$y = \left| \begin{bmatrix} 4 & 1 & 1 \\ 1 & -1 & 1 \\ -\frac{5}{8} & 0 & 1 \end{bmatrix} \right| = -\frac{25}{4}$$

$$z = \left| \begin{bmatrix} 4 & 4 & 1 \\ 1 & -1 & -1 \\ -\frac{5}{8} & \frac{3}{8} & 0 \end{bmatrix} \right| = \frac{15}{4}$$

Therefore we get our vector β which is our fourth basis vector a_4 :

$$a_4 = \beta = \left(-\frac{9}{49}; \frac{25}{49}; -\frac{15}{49}; 1 \right)$$

We conclude with our orthogonal basis vectors:

$$a_1 = (4; 4; 1; -1)$$

$$a_2 = (1; -1; 1; 1)$$

$$a_3 = \left(-\frac{5}{8}; \frac{3}{8}; 1; 0 \right)$$

$$a_4 = \left(-\frac{9}{49}; \frac{25}{49}; -\frac{15}{49}; 1 \right)$$

3 Task 3

Given the basis vectors:

$$\{a_1 = (2, 3, 5); a_2 = (0, 1, 2); a_3 = (1, 0, 0)\}$$

We need to find the matrix that transforms these basis vectors to:

$$\{e_1, e_2, e_3\} \Rightarrow \{(1, 1, 3); (1, 1, 0); (1, 2, 0)\}$$

First we will form a linear system of equations:

$$\begin{cases} e_1 = x_1 a_1 + x_2 a_2 + x_3 a_3 \\ e_2 = y_1 a_1 + y_2 a_2 + y_3 a_3 \\ e_3 = z_1 a_1 + z_2 a_2 + z_3 a_3 \end{cases}$$

Now we solve each equations separately by forming a yet another system of equations for each entry of e_1 and solve them with Cramer's rule.
I will skip Cramer's parts because we already know it from the first semester.

Solve for e_1 :

$$(1, 1, 3) = x_1(2, 3, 5) + x_2(0, 1, 2) + x_3(1, 0, 0) \Rightarrow (1, 1, 3) = (2x_1 + x_3; 3x_1 + x_2; 5x_1 + 2x_2)$$

$$\begin{cases} 2x_1 + x_3 = 1 \\ 3x_1 + x_2 = 1 \\ 5x_1 + 2x_2 = 3 \end{cases} \Rightarrow \begin{cases} x_1 = -1 \\ x_2 = 4 \\ x_3 = 3 \end{cases}$$

Solve for e_2 :

$$(1, 1, 0) = (2x_1 + x_3; 3x_1 + x_2; 5x_1 + 2x_2)$$

$$\begin{cases} 2x_1 + x_3 = 1 \\ 3x_1 + x_2 = 1 \\ 5x_1 + 2x_2 = 0 \end{cases} \Rightarrow \begin{cases} x_1 = 2 \\ x_2 = -5 \\ x_3 = -3 \end{cases}$$

Solve for e_3 :

$$(1, 2, 0) = (2x_1 + x_3; 3x_1 + x_2; 5x_1 + 2x_2)$$

$$\begin{cases} 2x_1 + x_3 = 1 \\ 3x_1 + x_2 = 2 \\ 5x_1 + 2x_2 = 0 \end{cases} \Rightarrow \begin{cases} x_1 = 4 \\ x_2 = -10 \\ x_3 = -7 \end{cases}$$

From this we get our final system of equations:

$$\begin{cases} e_1 = -a_1 + 4a_2 + 3a_3 \\ e_2 = 2a_1 - 5a_2 - 3a_3 \\ e_3 = 4a_1 - 10a_2 - 7a_3 \end{cases}$$

$$A = \begin{bmatrix} -1 & 4 & 3 \\ 2 & -5 & -3 \\ 4 & -10 & -7 \end{bmatrix}$$

4 Task 4

We are given matrix A to find the Transition matrix α : Find the Transition matrix α :

$$a \rightarrow A = \begin{bmatrix} 1 & -3 & 3 \\ -2 & -6 & 13 \\ -1 & -4 & 8 \end{bmatrix}$$

$$\gamma_\alpha(x) = |x \cdot I - A| \Rightarrow \left| \begin{bmatrix} x & 0 & 0 \\ 0 & x & 0 \\ 0 & 0 & x \end{bmatrix} - \begin{bmatrix} 1 & -3 & 3 \\ -2 & -6 & 13 \\ -1 & -4 & 8 \end{bmatrix} \right| = x^3 - 3x^2 + 3x - 1 = (x-1)^3$$

$$(x-1)^3 = 0$$

$$x = 1$$

$$\alpha(a) = \lambda a$$

$$A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 & -3 & 3 \\ -2 & -6 & 13 \\ -1 & -4 & 8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 - 3x_2 + 3x_3 \\ -2x_1 - 6x_2 + 13x_3 \\ -x_3 - 4x_2 + 8x_3 \end{bmatrix}$$

Now we form a linear system of equations to solve:

$$\begin{cases} -3x_2 + 3x_3 = 0 \\ -2x_1 - 7x_2 + 13x_3 = 0 \\ -x_1 - 4x_2 + 7x_3 = 0 \end{cases}$$

Now we form a matrix to solve the system:

$$\begin{bmatrix} 0 & -3 & 3 \\ -2 & -7 & 13 \\ -1 & -4 & 7 \end{bmatrix}$$

$$\Delta_1 = -3 \neq 0$$

$$\Delta_2 = \begin{bmatrix} 0 & -3 \\ -2 & -7 \end{bmatrix} = -6 \neq 0$$

$$\begin{bmatrix} 0 & -3 & 3 \\ -2 & -7 & 13 \\ -1 & -4 & 7 \end{bmatrix} = 0 \Rightarrow \text{rank}(A) = 0$$

Because the rank is 3, we ignore the third row in the system and solve for the first two:

$$\begin{cases} -3x_2 = -3x_3 \\ -2x_1 - 7x_2 = -13x_3 \end{cases} = \begin{cases} x_1 = 3 \\ x_2 = 1 \end{cases}$$

From this we get our vector:

$$(3, 1, 1)$$

ԳՐԱԿԱՆՈՒԹՅՈՒՆ

1. Կուրոշ <<բարձրագույն հանրահաշվի դասընթաց>>
2. Գելֆանդ <<Դասախոսություններ գծային հանրահաշվից>>
3. Մովսիսյան <<բարձրագույն հանրահաշիվ և թվերի տեսություն>>



Редактировать в WPS Office