

ՀՀ ԿԳՄՍ ՆԱԽԱՐԱՐՈՒԹՅՈՒՆ
ՀԱՅԱՍՏԱՆԻ ԵՎՐՈՊԱԿԱՆ ՀԱՄԱԼՍԱՐԱՆ
ՏԵՂԵԿԱՏՎԱԿԱՆ ՀԱՄԱԿԱՐԳԵՐ

ԱՆՀԱՏԱԿԱՆ ԱՇԽԱՏԱՆՔ

Դիսկրետ Մաթեմատիկա՝ տարբերակ 19

ԱՌԱՐԿԱ՝ ԴԻՍԿՐԵՏ ՄԱԹԵՄԱՏԻԿԱ
ԿՈՒՐՍ՝ ԱՌԱՋԻՆ
ՈՒՍԱՆՈՂ՝ ՍԵՐՈՔ ՏԻԳՐԱՆՅԱՆ
ԴԱՍԱԽՈՍ՝ Ն. ՀԱԿՈԲՅԱՆ

ԵՐԵՎԱՆ 2025

Discrete Mathematics

Serob Tigranyan

May 19, 2025

Contents

1 Task 1	2
1.1 Reflexivity	2
1.1.1 Reflexive Closure	2
1.2 Symmetry	2
1.2.1 Symmetric Closure	3
1.3 Transitivity	3
1.3.1 Transitive Closure	3
2 Task 2	4
2.1 Calculate the numbers that 1000 is divisible by each number . .	4
2.2 Subtract overlap	4
2.3 Add back numbers counted 3 times	5
2.4 Putting it all together using Inclusion-Exclusion Formula	5
3 Task 3	6

1 Task 1

We are given:

$$A = \{1, 2, 3, 4, 5\}$$

And a relation matrix a where $R \subseteq A \times A$:

$$a = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

We need to find whether relation a is **reflexive**, **symmetric** and **transitive**, if not then find their **closures** respectively.

1.1 Reflexivity

A relation is **reflexive** if every element is related to itself, e.g., $a_{ii} = 1$ for all i .

This means that the diagonal elements are all 1's, but it's not in our case therefore relation a is **not reflexive**.

1.1.1 Reflexive Closure

Reflexive closure for relation a would be a_{ref} where all the diagonal elements are 1's:

$$a_{ref} = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

1.2 Symmetry

A relation is **symmetric** if $a_{ij} = a_{ji}$.

We can check this by transposing our relation matrix and if $T(a) = a$ then the relation is **symmetric**:

$$T(a) = \begin{bmatrix} 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 \end{bmatrix}$$

We can clearly observe that the transposed matrix differs from the original relation matrix therefore:

$$T(a) \neq a$$

So the relation a is **not symmetric**.

1.2.1 Symmetric Closure

The **symmetric closure** for relation a would be a_{sym} where every $a_{ij} = a_{ji} = 1$:

$$a_{sym} = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

1.3 Transitivity

A relation is **transitive** if whenever $(i, j) \in R$ and $(j, k) \in R$, then $(i, k) \in R$.

We can easily check by taking the **boolean** square of the relation matrix a :

$$a^2 = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{bmatrix}$$

If all the 1's in a^2 exists in a then the relation is **transitive**, but this is not the case for us therefore the relation a is **not transitive**.

1.3.1 Transitive Closure

We can use Warshall's algorithm to find the transitive closure for relation a . After applying Warshall's algorithm to our initial relation matrix we get:

$$a_{trans} = \begin{bmatrix} 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

2 Task 2

We are asked to find how many integers below or equal to 1000 are divisible by 8, 22 or 32.

Let:

- A : Numbers divisible by 8
- B : Numbers divisible by 22
- C : Numbers divisible by 32

2.1 Calculate the numbers that 1000 is divisible by each number

We start by finding how many numbers up to 1000 are divisible by:

8: Divide 1000 by 8:

$$\left\lfloor \frac{1000}{8} \right\rfloor = 125$$

22: Divide 1000 by 22:

$$\left\lfloor \frac{1000}{22} \right\rfloor = 45$$

32: Divide 1000 by 32:

$$\left\lfloor \frac{1000}{32} \right\rfloor = 31$$

2.2 Subtract overlap

Then we will need to subtract the overlap numbers counted twice.
Some numbers are divisible by both 8 and 22, or 8 and 32, etc.
We subtract them so we don't count them twice.

To do this, we find how many numbers are divisible by both pairs:

- 8 and 22: Find LCM (Least Common Multiple) of 8 and 22

$$LCM(8, 22) = 88$$

$$\left\lfloor \frac{1000}{88} \right\rfloor = 11$$

- 8 and 32: Find LCM of 8 and 32

$$LCM(8, 32) = 32$$

$$\left\lfloor \frac{1000}{32} \right\rfloor = 31$$

- 22 and 32: Find LCM of 22 and 32

$$LCM(22, 32) = 352$$

$$\left\lfloor \frac{1000}{352} \right\rfloor = 2$$

2.3 Add back numbers counted 3 times

Some numbers are divisible by all three so we add these back because we subtracted them earlier.

- $LCM(8, 22, 32) = 352$

$$\left\lfloor \frac{1000}{352} \right\rfloor = 2$$

2.4 Putting it all together using Inclusion-Exclusion Formula

The **Inclusion-Exclusion** Formula states the following:

$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

Plug in our values:


- $A = 125$
- $B = 45$
- $C = 31$
- $AB = 11$
- $AC = 31$
- $BC = 2$
- $ABC = 2$

Therefore:

$$125 + 45 + 31 - 11 - 31 - 2 + 2 = 159$$

3 Task 3

We are given graphs H and G :



We must determine if the given graphs are **isomorphic**. First we recall the definition of two isomorphic graphs:

Let $G_1 = (V_1, E_1)$ & $G_2 = (V_2, E_2)$ be two simple graphs where:

- V_1, V_2 are the sets of vertices
- E_1, E_2 are the sets of edges

Then G_1 & G_2 are **isomorphic** if there exists a bijective function $\varphi: V_1 \rightarrow V_2$ such that:

$$\{u, v\} \in E_1, \text{ if and only if } \{\varphi(u), \varphi(v)\} \in E_2$$

In simpler terms G & H are **isomorphic** if there exists a bijection function φ that directly maps G 's vertices to H 's vertices (and vice-versa) and that each degree of both vertices match.

In our case it is visually apparent that these two graphs are not **isomorphic**. More specifically, the degrees of u_3, v_3 do not match, u_3 has degree of two while v_3 has degree of just one.

Therefore these two graphs are not **isomorphic**.

ԳՐԱԿԱՆՈՒԹՅՈՒՆ

1. Տոնոյան Ռ. Ն., Դիսկրետ մաթեմատիկայի դասընթաց: Երևան, ԵՊՀ, 1999, 130 էջ:
2. Ն. Ա. Սահակյան, Դիսկրետ մաթեմատիկա 1, 2, ՀՊՃՀ. – Եր.: Ճարտարագետ, 2014, 68 էջ:
3. Rosen, Kenneth H, Discrete mathematics and its applications-8th edition, 2021, 1118 pages.
4. A. Ross, Kenneth, Discrete mathematics, Edition, 2, illustrated ; Publisher, Prentice Hall, 1988 ; Original from, the University of Michigan ; Digitized, Nov 20, 2007 ; ISBN, 0132154277, 9780132154277.