

Table 5. Association between Gender and Disease Pattern among Study Participants (N=600)

Endocrine Diseases & Comorbidities	Overall N = 600¹	Female N = 414¹	Male N = 186¹	p-value²
Type 2 Diabetes	318 (53%)	192 (46%)	126 (68%)	<0.001
Hypothyroidism (Congenital)	12 (2.0%)	6 (1.4%)	6 (3.2%)	0.2
Hypothyroidism (Adult)	174 (29%)	146 (35%)	28 (15%)	<0.001
Hyperthyroidism	24 (4.0%)	17 (4.1%)	7 (3.8%)	0.8
Hypoparathyroidism	24 (4.0%)	19 (4.6%)	5 (2.7%)	0.3
Hyperprolactinemia	12 (2.0%)	10 (2.4%)	2 (1.1%)	0.4
PCOS & Menstrual Abnormality	24 (4.0%)	24 (5.8%)	0 (0%)	<0.001
Male Hypogonadism	12 (2.0%)	0 (0%)	12 (6.5%)	<0.001
Hypertension	60 (10%)	44 (11%)	16 (8.6%)	0.4
Dyslipidemia	48 (8.0%)	34 (8.2%)	14 (7.5%)	0.8
Coronary/Cerebrovascular disease	12 (2.0%)	7 (1.7%)	5 (2.7%)	0.5
Neuropathy	12 (2.0%)	10 (2.4%)	2 (1.1%)	0.4

¹n (%)

²Pearson's Chi-squared test; Fisher's exact test