

Table 6. Association between Age Group and Disease Pattern among Study Participants (N=600)

Endocrine Diseases & Comorbidities	Overall N = 600¹	<50 Years N = 364¹	>60 Years N = 109¹	50-59 Years N = 127¹	p-value²
Type 2 Diabetes	318 (53%)	107 (29%)	108 (99%)	103 (81%)	<0.001
Hypothyroidism (Congenital)	12 (2.0%)	12 (3.3%)	0 (0%)	0 (0%)	0.016
Hypothyroidism (Adult)	174 (29%)	174 (48%)	0 (0%)	0 (0%)	<0.001
Hyperthyroidism	24 (4.0%)	1 (0.3%)	1 (0.9%)	22 (17%)	<0.001
Hypoparathyroidism	24 (4.0%)	24 (6.6%)	0 (0%)	0 (0%)	<0.001
Hyperprolactinemia	12 (2.0%)	12 (3.3%)	0 (0%)	0 (0%)	0.016
PCOS & Menstrual Abnormality	24 (4.0%)	23 (6.3%)	0 (0%)	1 (0.8%)	<0.001
Male Hypogonadism	12 (2.0%)	11 (3.0%)	0 (0%)	1 (0.8%)	0.10
Hypertension	60 (10%)	36 (9.9%)	18 (17%)	6 (4.7%)	0.011
Dyslipidemia	48 (8.0%)	28 (7.7%)	16 (15%)	4 (3.1%)	0.005
Coronary/Cerebrovascular disease	12 (2.0%)	6 (1.6%)	6 (5.5%)	0 (0%)	0.009
Neuropathy	12 (2.0%)	8 (2.2%)	3 (2.8%)	1 (0.8%)	0.5

¹n (%)

²Pearson's Chi-squared test; Fisher's exact test