

Table 7. Association between Gender and Endocrine Diseases & other comorbidities Medications (N=600)

Endocrine Diseases & Comorbidities Medications	Overall N = 600 ¹	Female N = 414 ¹	Male N = 186 ¹	p-value ²
Metformin	342 (57%)	216 (52%)	126 (68%)	<0.001
Gliclazide	60 (10%)	36 (8.7%)	24 (13%)	0.11
Glimeperide	144 (24%)	79 (19%)	65 (35%)	<0.001
Linagliptin	168 (28%)	118 (29%)	50 (27%)	0.7
Sitagliptin	144 (24%)	86 (21%)	58 (31%)	0.006
Vindagliptin	12 (2.0%)	8 (1.9%)	4 (2.2%)	>0.9
Empaglifozin	144 (24%)	81 (20%)	63 (34%)	<0.001
Short-acting Insulin	48 (8.0%)	32 (7.7%)	16 (8.6%)	0.7
Long-acting Insulin	96 (16%)	64 (15%)	32 (17%)	0.6
Mixed Insulin	120 (20%)	79 (19%)	41 (22%)	0.4
Levothyroxine	186 (31%)	152 (37%)	34 (18%)	<0.001
Carbimazole	24 (4.0%)	17 (4.1%)	7 (3.8%)	0.8
Estrogen + Progesteron	24 (4.0%)	24 (5.8%)	0 (0%)	<0.001
Cabergolin	12 (2.0%)	10 (2.4%)	2 (1.1%)	0.4
Testosterone	12 (2.0%)	0 (0%)	12 (6.5%)	<0.001
Vit-D + Calcium	24 (4.0%)	19 (4.6%)	5 (2.7%)	0.3
Losartan	30 (5.0%)	18 (4.3%)	12 (6.5%)	0.3
Amlodipin	30 (5.0%)	21 (5.1%)	9 (4.8%)	>0.9
Frusemide	20 (3.3%)	14 (3.4%)	6 (3.2%)	>0.9
Atorvastatin	48 (8.0%)	31 (7.5%)	17 (9.1%)	0.5
Clopidogrel	12 (2.0%)	11 (2.7%)	1 (0.5%)	0.12
PPI, Antibiotics, Vitamins, Sedatives	112 (19%)	82 (20%)	30 (16%)	0.3

¹n (%)

²Pearson's Chi-squared test; Fisher's exact test