

MACHINE LEARNING

In Q1 to Q11, only one option is correct, choose the correct option:

1. Which of the following methods do we use to find the best fit line for data in Linear Regression?

- A) Least Square Error B) Maximum Likelihood
- C) Logarithmic Loss D) Both A and B

Ans. A) Least Square Error

2. Which of the following statement is true about outliers in linear regression?

- A) Linear regression is sensitive to outliers B) linear regression is not sensitive to outliers
- C) Can't say D) none of these

Ans. A) Linear regression is sensitive to outliers.

3. A line falls from left to right if a slope is _____?

- A) Positive B) Negative
- C) Zero D) Undefined

Ans. B) Negative

4. Which of the following will have symmetric relation between dependent variable and independent variable?

- A) Regression B) Correlation
- C) Both of them D) None of these

Ans. B) Correlation

5. Which of the following is the reason for over fitting condition?

- A) High bias and high variance B) Low bias and low variance
- C) Low bias and high variance D) none of these

Ans. A) High bias and high variance

6. If output involves label then that model is called as:

- A) Descriptive model B) Predictive modal
- C) Reinforcement learning D) All of the above

Ans. B) Predictive modal

7. Lasso and Ridge regression techniques belong to _____?

- A) Cross validation B) Removing outliers
- C) SMOTE D) Regularization

Ans. D) Regularization

8. To overcome with imbalance dataset which technique can be used?

- A) Cross validation B) Regularization
- C) Kernel D) SMOTE

Ans. D) SMOTE

9. The AUC Receiver Operator Characteristic (AUCROC) curve is an evaluation metric for binary classification problems. It uses _____ to make graph?

- A) TPR and FPR B) Sensitivity and precision
- C) Sensitivity and Specificity D) Recall and precision

Ans. C) Sensitivity and Specificity

10. In AUC Receiver Operator Characteristic (AUCROC) curve for the better model area under the curve should be less.

- A) True B) False

Ans. B) False

11. Pick the feature extraction from below:

- A) Construction bag of words from a email
- B) Apply PCA to project high dimensional data
- C) Removing stop words
- D) Forward selection

Ans. B) Apply PCA to project high dimensional data

In Q12, more than one options are correct, choose all the correct options:

12. Which of the following is true about Normal Equation used to compute the coefficient of the Linear Regression?

- A) We don't have to choose the learning rate.
- B) It becomes slow when number of features is very large.
- C) We need to iterate.
- D) It does not make use of dependent variable.

Ans. A) We don't have to choose the learning rate.

MACHINE LEARNING

Q13 and Q15 are subjective answer type questions, Answer them briefly.

13. Explain the term regularization?

Ans. The word regularize means to make things regular or acceptable. This is exactly why we use it for. Regularizations are techniques used to reduce the error by fitting a function appropriately on the given training set and avoid overfitting.

14. Which particular algorithms are used for regularization?

Ans. There are three main regularization techniques, namely:

1. Ridge Regression (L2 Norm)

Ridge regression is also called L2 norm or regularization. When using this technique, we add the sum of weight's square to a loss function and thus create a new loss function.

2. Lasso (L1 Norm)

In ridge regression, loss function along with the optimization algorithm brings parameters near to zero but not actually zero, while lasso eliminates less important features and sets respective weight values to zero. Thus, lasso also performs feature selection along with regularization.

3. Dropout

Dropout is a regularization technique used in neural networks. It prevents complex co-adaptations from other neurons.

15. Explain the term error present in linear regression equation?

Ans. Linear regression most often uses mean-square error (MSE) to calculate the error of the model. MSE is calculated by:

1. measuring the distance of the observed y-values from the predicted y-values at each value of x;
2. squaring each of these distances;
3. Calculating the mean of each of the squared distances.

Linear regression fits a line to the data by finding the regression coefficient that results in the smallest MSE.