

5. DISCRETE RANDOM VARIABLES

Introduction

- We introduced random events and probability in the previous chapter.
- Now we will formalize the random events mathematically.
- This chapter focuses on discrete random variables, while the next chapter will focus on continuous random variables.

Random variables

Random variables

- A **random variable** is a variable whose values are determined by chance.
 - ▣ In other words, random variable is the random outcome of an experiment.
- Discrete random variable – random variable that assumes countable values.
 - ▣ Example:
 - The number of students getting an A
 - The number of fish caught on a fishing trip
- We usually denote a random variable by X (uppercase letter), and x (lowercase letter) for a specific value of the random variable.

Probability distribution

- The **probability distribution** is a table or function, that lists all the possible values for a random variable and their corresponding probabilities.
- Also called **probability mass function** (pmf) for discrete random variables.
- Notation: $P(x)$ or $P(X = x)$ is the probability that the random variable X takes the value x .
- Two properties of probability distribution of discrete random variable:
 - ▣ $0 \leq P(X = x) \leq 1$
 - ▣ $\sum_x P(X = x) = 1$

Example

- Consider rolling a die.
 - ▣ Let X be the observed outcome when rolling a die.
 - ▣ The following table is the probability distribution of X :

x	1	2	3	4	5	6
$P(X = x)$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$	$1/6$

- ▣ $P(X = 1) = 1/6$, means that the probability of X takes value 1 is $1/6$
- ▣ Also note that

$$\sum P(X = x) = P(1) + P(2) + \cdots + P(6) = 1$$

Exercise

- Determine whether the following represents a probability distribution. If it does not, state why.

7.

X	15	16	20	25
$P(X)$	0.2	0.5	0.7	-0.8

8.

X	5	7	9
$P(X)$	0.6	0.8	-0.4

9.

X	-5	-3	0	2	4
$P(X)$	0.1	0.3	0.2	0.3	0.1

10.

X	20	30	40	50
$P(X)$	0.05	0.35	0.4	0.2

11.

X	3	6	9	1
$P(X)$	0.3	0.4	0.3	0.1

12.

X	3	7	9	12	14
$P(X)$	$\frac{4}{13}$	$\frac{1}{13}$	$\frac{3}{13}$	$\frac{1}{13}$	$\frac{2}{13}$

Two properties

1) $0 \leq P(X=x) \leq 1$

2) $\sum P(X=x) = 1$

7) $P(X=25) = -0.8 < 0$
Not a probability distribution

8) $P(X=9) = -0.4 < 0$

Not a probability distribution

$$9) 0 \leq P(X=x) \leq 1$$

$$\begin{aligned} \sum P(X=x) &= 0.1 + 0.3 \\ &\quad + 0.2 + 0.3 \\ &\quad + 0.1 \\ &= 1 \end{aligned}$$

Yes, it is a probability distribution

$$10) 0 \leq P(X=x) \leq 1$$

$$\begin{aligned} \sum P(X=x) &= 0.65 + 0.35 \\ &\quad + 0.4 + 0.2 \\ &= 1 \end{aligned}$$

Yes, it is a probability distribution

$$11) 0 \leq P(X=x) \leq 1$$

$$\begin{aligned} \sum P(X=x) &= 0.3 + 0.4 + 0.3 \\ &\quad + 0.1 \\ &= 1.1 \neq 1 \end{aligned}$$

This is not a probability distribution

$$12) 0 \leq P(X=x) \leq 1$$

$$\begin{aligned} \sum P(X=x) &= \frac{4}{13} + \frac{1}{13} + \frac{3}{13} \\ &\quad + \frac{1}{13} + \frac{2}{13} \\ &= \frac{11}{13} \neq 1 \end{aligned}$$

Not a probability distribution

Exercise

- An industrial psychologist administered a personality inventory test for passive-aggressive traits to 150 employees. Individuals were given a score from 1 to 5, where 1 was extremely passive and 5 extremely aggressive. A score of 3 indicated neither trait. Construct a probability distribution for the random variable X .

Score, x	Frequency, f
1	24
2	33
3	42
4	30
5	21

$$P(X=n)$$

$$24/150 = 0.16$$

$$33/150 = 0.22$$

$$0.28$$

$$0.2$$

$$0.14$$

Exercise

- At a drop-in mathematics tutoring center, each teacher sees 4 to 8 students per hour. The probability that a tutor sees 4 students in an hour is 0.117; 5 students, 0.123; 6 students, 0.295; and 7 students, 0.328.
- a) Find the probability that a tutor sees 8 students in an hour
 - b) Construct the probability distribution.
 - c) Find the probability that a tutor sees 6 or less students in an hour.

$$a) \sum P(X=n) = 1$$

$$P(4) + P(5) + P(6) + P(7) + P(8) = 1$$

$$P(8) = 1 - P(4) - P(5) - P(6) - P(7)$$

$$= 1 - 0.117 - 0.123 - 0.295 - 0.328$$

$$= 0.137$$

b)

x	4	5	6	7	8
$P(X=x)$	0.117	0.123	0.295	0.328	0.137

c)

$$\begin{aligned}
 P(X \leq 6) &= P(4) + P(5) + P(6) \\
 &= 0.117 + 0.123 + 0.295 \\
 &= 0.535
 \end{aligned}$$

Exercise

- The number 1, 2, 3 and 4 are printed one each on one side of card. The cards are placed face down and mixed. Choose two cards at random; and let X be the sum of the two numbers. Construct the probability distribution for this random variable X .

Possible outcomes : $1\frac{1}{2}2$ $1\frac{1}{2}3$ $1\frac{1}{2}4$ $2\frac{1}{2}3$ $2\frac{1}{2}4$ $3\frac{1}{2}4$

X : 3 4 5 5 6 7

x	3	4	5	6	7
$P(X=x)$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{2}{6} = \frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{6}$

eg: $P(X < 5) = P(3) + P(4)$

$$= \frac{1}{6} + \frac{1}{6}$$

$$= \frac{1}{3}$$

Cumulative distribution function

- Another important concept in the distribution of random variable is the **cumulative distribution function** (also called **distribution function**).
- If X is a discrete random variable, the cumulative distribution function is given by

$$F(x) = P(X \leq x) = \sum_{t \leq x} P(X = t)$$

for $-\infty < x < \infty$.

- $F(x)$ is the probability of X taking value less than equal to x .

Cumulative distribution function

- Properties of a discrete distribution function:
 - ▣ $0 \leq F(x) \leq 1$ for all x
 - ▣ If $a < b$, then $F(a) \leq F(b)$
 - ▣ $F(x)$ is a step function
 - ▣ $F(-\infty) = 0$ and $F(\infty) = 1$

- Additionally:
 - ▣ $P(a < X \leq b) = F(b) - F(a)$
 - ▣ $P(X > a) = 1 - P(X \leq a) = 1 - F(a)$
 - ▣ $P(X < a) = P(X \leq a - 1) = F(a - 1)$
 - ▣ $P(X = a) = P(X \leq a) - P(X \leq a - 1) = F(a) - F(a - 1)$

Example – CDF

- Using the following probability distribution, find its cumulative distribution function $F(x)$.

x	0	1	2	3
$P(X = x)$	$1/8$	$3/8$	$3/8$	$1/8$

- From the table we get:

$$\begin{aligned}
 & \text{Handwritten: } P(X \leq 0) \\
 F(0) &= \frac{1}{8} \\
 & \text{Handwritten: } P(X \leq 1) \\
 F(1) &= \frac{1}{8} + \frac{3}{8} = \frac{4}{8} = \frac{1}{2} \\
 & \text{Handwritten: } P(X \leq 2) \\
 F(2) &= \frac{4}{8} + \frac{3}{8} = \frac{7}{8} \\
 F(3) &= \frac{7}{8} + \frac{1}{8} = 1
 \end{aligned}$$

Example – CDF

- Additionally, $F(x) = 0$ for $x < 0$, and $F(x) = 1$ for $x > 3$.
- Therefore:

$$F(x) = \begin{cases} 0, & x < 0 \\ \underline{1/8}, & \underline{0 \leq x} < 1 \\ \underline{1/2}, & \underline{1 \leq x} < 2 \\ \underline{7/8}, & 2 \leq x < 3 \\ 1, & x \geq 3 \end{cases}$$

Exercise

- You are given the following table of probability distribution of X . Find the distribution function $F(x)$.

x	1	2	3	4	5
$P(X = x)$	0.16	0.22	0.28	0.20	0.14

$p(X \leq x)$ 0.16 0.38 0.66 0.86 1

$\swarrow +0.22$ $\swarrow +0.28$

$$F(x) = \begin{cases} 0, & x < 1 \\ 0.16, & 1 \leq x < 2 \\ 0.38, & 2 \leq x < 3 \\ 0.66, & 3 \leq x < 4 \\ 0.86, & 4 \leq x < 5 \\ 1, & x \geq 5 \end{cases}$$

Exercise

- Given the following cumulative distribution function for a discrete random variable X :

$$F(x) = \begin{cases} 0, & x < 1 \\ 1/6, & 1 \leq x < 2 \\ 1/2, & 2 \leq x < 3 \\ 5/6, & 3 \leq x < 4 \\ 1, & x \geq 4 \end{cases}$$

Handwritten calculations for probabilities:

$$P(X=1) = \frac{1}{6} - 0 = \frac{1}{6}$$

$$P(X=2) = \frac{1}{2} - \frac{1}{6} = \frac{1}{3}$$

$$P(X=3) = \frac{5}{6} - \frac{1}{2} = \frac{1}{3}$$

$$P(X=4) = 1 - \frac{5}{6} = \frac{1}{6}$$

- Find
- $P(1 < X < 3)$
 - $P(X > 2)$
 - $P(2 \leq X \leq 4)$
 - The probability distribution of X

$$\begin{aligned}
 \text{a) } P(1 < X < 3) &= P(X \leq 2) - P(X \leq 1) = F(2) - F(1) \\
 &= P(1 < X \leq 2) \\
 &= P(X=2) = \frac{1}{2} - \frac{1}{6} = \frac{1}{3}
 \end{aligned}$$

$$\text{b) } P(X > 2) = 1 - P(X \leq 2) = 1 - F(2) = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\begin{aligned}
 \text{c) } P(2 \leq X \leq 4) &= P(X \leq 4) - P(X \leq 1) = F(4) - F(1) \\
 &= P(1 < X \leq 4) = 1 - \frac{1}{6} = \frac{5}{6}
 \end{aligned}$$





d)

x	1	2	3	4
$P(X=x)$	$\frac{1}{6}$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{6}$

Graphing probability distribution


- The probability distribution can be shown visually using for example:
 - ▣ Histogram
 - ▣ Line plot
- The distribution function can be shown visually using:
 - ▣ Line plot
- Presenting the probability distribution visually helps with describing the distribution.

Example – graphing probability distribution



Example – graphing distribution function

$$F(x) = \begin{cases} 0, & x < 1 \\ 1/6, & 1 \leq x < 2 \\ 1/2, & 2 \leq x < 3 \\ 5/6, & 3 \leq x < 4 \\ 1, & x \geq 4 \end{cases}$$



Mean, variance, standard deviation of discrete random variables

Mean of discrete random variables

- The mean, variance and standard deviation for random variable are calculated in a similar way to grouped data.
- We use the probability $P(X = x)$ to replace f/N (frequency over total population).
- The **mean of random discrete variable** is defined as:

$$\mu = \sum xP(X = x)$$

Variance and standard deviation of discrete random variables

- Similarly, the variance and standard deviation can be calculated using:

$$\sigma^2 = \sum (x - \mu)^2 P(X = x), \quad \sigma = \sqrt{\sum (x - \mu)^2 P(X = x)}$$

- As before, for easier computation:

$$\sigma^2 = \sum x^2 P(X = x) - \mu^2,$$

$$\sigma = \sqrt{\sum x^2 P(X = x) - \mu^2}$$

Example

- Three coins are tossed and let X be the number of heads that occur.

x	0	1	2	3
$P(X = x)$	$1/8$	$3/8$	$3/8$	$1/8$

$$\begin{aligned}\mu &= \sum xP(X = x) \\ &= \left(0 \times \frac{1}{8}\right) + \left(1 \times \frac{3}{8}\right) + \left(2 \times \frac{3}{8}\right) + \left(3 \times \frac{1}{8}\right) \\ &= 1.5\end{aligned}$$

$$\begin{aligned}\sigma^2 &= \sum x^2P(X = x) - \mu^2 \\ &= \left(0^2 \times \frac{1}{8}\right) + \left(1^2 \times \frac{3}{8}\right) + \left(2^2 \times \frac{3}{8}\right) + \left(3^2 \times \frac{1}{8}\right) - 1.5^2 \\ &= 3 - 2.25 = 0.75\end{aligned}$$

$$\sigma = \sqrt{0.75} = 0.8660$$

Another example

- The probability distribution for the number of batteries sold over the weekend at a convenience store is given below:

x	2	4	6	8
$P(x)$	0.20	0.40	0.32	0.08

$$\begin{aligned}\mu &= \sum xP(X = x) \\ &= 2(0.2) + 4(0.4) + 6(0.32) + 8(0.08) \\ &= 4.56\end{aligned}$$

$$\begin{aligned}\sigma^2 &= \sum x^2P(X = x) - \mu^2 \\ &= 2^2(0.2) + 4^2(0.4) + 6^2(0.32) + 8^2(0.08) - 4.56^2 \\ &= 23.84 - 20.7936 = 3.046\end{aligned}$$

$$\sigma = \sqrt{3.046} = 1.745$$

Exercise

5.29 Let x be the number of heads obtained in two tosses of a coin. The following table lists the probability distribution of x .

x	0	1	2
$P(x)$.25	.50	.25

Calculate the mean and standard deviation of x . Give a brief interpretation of the value of the mean.

$$\begin{aligned}\mu &= \sum n P(X=n) \\ &= (0 \times 0.25) + (1 \times 0.5) + (2 \times 0.25) \\ &= 1\end{aligned}$$

$$\begin{aligned}
 \sigma^2 &= \sum x^2 P(x=x) - \mu^2 \\
 &= (0^2 \times 0.25) + (1^2 \times 0.5) + (2^2 \times 0.25) \\
 &\quad - 1^2 \\
 &= 0.5
 \end{aligned}$$

$$\sigma = \sqrt{0.5} = 0.7071$$

If the experiment is repeated many times,
the average number of heads in two tosses
is 1

Expectation

- Another concept related to the mean for a probability distribution is that of expected value or expectation
- The **expected value** of a random variable is the theoretical average of the variable.

- For discrete random variable, the formula is

$$E[X] = \sum xP(X = x) = \mu$$

- Also note that the variance

$$\sigma^2 = E[(X - \mu)^2] = E[X^2] - E[X]^2$$

$$\sum \downarrow x^2 P(X=x) \quad \uparrow \mu^2$$

Exercise

- A person pays \$2 to play a certain game by rolling a single die once. If a 1 or a 2 comes up, the person wins nothing. If, however, the player rolls a 3, 4, 5, or 6, he or she wins the difference between the number rolled and \$2. Find the expectation and variance for this game. Is the game fair?

Die outcome	1	2	3	4	5	6
Amount win	0	0	1	2	3	4

Let $X = \text{amount win}$

x	0	1	2	3	4
$P(X=x)$	$\frac{2}{6} = \frac{1}{3}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

$$\begin{aligned}
 E[X] = \mu &= \sum x P(X=x) = \left(0 \times \frac{1}{3}\right) + \left(1 \times \frac{1}{6}\right) + \left(2 \times \frac{1}{6}\right) \\
 &\quad + \left(3 \times \frac{1}{6}\right) + \left(4 \times \frac{1}{6}\right) \\
 &= \frac{5}{3} = 1.66\bar{7}
 \end{aligned}$$

$$\begin{aligned}
 \sigma^2 &= \sum x^2 P(X=x) - \mu^2 \\
 &= \left(0^2 \times \frac{1}{3}\right) + \left(1^2 \times \frac{1}{6}\right) + \left(2^2 \times \frac{1}{6}\right) + \left(3^2 \times \frac{1}{6}\right) \\
 &\quad + \left(4^2 \times \frac{1}{6}\right) - \left(\frac{5}{3}\right)^2 \\
 &= \frac{20}{9} = 2.22\bar{2}
 \end{aligned}$$

The game is not fair because the expected amount of winnings (\$1.67) is less than the cost (\$2)

Exercise

- The probability distribution for the number of batteries sold over the weekend at a convenience store is given below:

x	2	4	6	8
$P(X = x)$	0.20	0.40	0.32	0.08

- Calculate

- a) $E[X]$
- b) $E[2X]$
- c) $E[X^2]$

$$\begin{aligned} \text{a) } E[\underline{x}] &= \sum \underline{x} P(X=x) \\ &= (2 \times 0.2) + (4 \times 0.4) + (6 \times 0.32) \\ &\quad + (8 \times 0.08) \\ &= 4.56 \end{aligned}$$

$$b) E[\underline{2x}] = \sum \underline{2x} P(X=x)$$

$$= (2 \times 2 \times 0.2) + (2 \times 4 \times 0.4) + (2 \times 6 \times 0.32) \\ + (2 \times 8 \times 0.08)$$

$$= 9.12 \quad \left(\begin{array}{l} E[2x] = 2E[x] \\ E[ax+b] = aE[x] + b \end{array} \right)$$

$$c) E[\underline{x^2}] = \sum \underline{x^2} P(X=x)$$

$$= (2^2 \times 0.2) + (4^2 \times 0.4) + (6^2 \times 0.32) \\ + (8^2 \times 0.08)$$

$$= 23.84 \quad \left(E[x^2] \neq E[x]^2 \right)$$

Binomial distribution

Binomial experiment

- Consider a probability experiment that only has two outcomes – “success” or “failure”.
 - ▣ Eg: Toss a coin. If head is observed, then it is a success.
- This experiment is called a **Bernoulli** trial.
- On the other hand, suppose we repeat the trial a few times and count **how many success we observed**.
- These are called binomial experiments.

Binomial experiment

- A **binomial experiment** is a probability experiment that satisfies the following four requirements:
 - ▣ There must be a fixed number of trials.
 - ▣ Each trial can have only two outcomes. These outcomes can be considered as either “success” or “failure”.
 - ▣ The outcomes of each trial must be independent of one another.
 - ▣ The probability of a success must remain the same for each trial.
- “Success” does not imply that something good or positive has occurred.

Binomial distribution

- Let X be the number of successes observed in a binomial experiment with n number of trials.
- Then the probability distribution of X is called the **binomial distribution**.
- Example:
 - ▣ Toss a coin 10 times, and let X be the number of times head is observed.
 - ▣ Roll a die 5 times, and let X be the number of times odd numbers are observed.

Notation for binomial distribution

- We usually use these notations:
 - ▣ n : Total number of trials.
 - ▣ p : The probability of success.
 - ▣ q : The probability of failure. $q = 1 - p$.
 - ▣ X : Number of successes in n trials.

- Note that $0 \leq X \leq n$. X can take the values 0, 1, 2, ..., n

Probability distribution for binomial


- The probability distribution of X :

$$P(X = x) = {}^nC_x p^x q^{n-x}$$

for $x = 0, 1, \dots, n$

- How do we get this probability?
 - ▣ Out of the n trials, x number of them are successes.
 - Total possible way to get x successes out of n trials is nC_x
 - ▣ There are x number of successes, each with probability p .
 - The probability for this is p^x
 - ▣ There are $n - x$ number of failures, each with probability q .
 - The probability for this is q^{n-x}
 - ▣ Since they are independent, we can multiply their probabilities.

Probability distribution for binomial



Mean and variance for binomial distribution

□ Mean:

$$\mu = np$$

□ Variance:

$$\sigma^2 = npq = np(1 - p)$$

□ Standard deviation:

$$\sigma = \sqrt{npq} = \sqrt{np(1 - p)}$$

Example

- A coin is tossed 3 times.
 - ▣ Find the probability that we observe 2 heads.
 - ▣ Find the mean, variance, and standard deviation of the number of heads that will be obtained.
- Let X be the number of times we observe heads. X has a binomial distribution.
- In this case, “success” is defined as “observing head”.

$$p = P(\text{head}) = 0.5, \quad q = P(\text{tail}) = 0.5$$

- The number of trials, n is 3.
- Probability of observing 2 heads:

$$P(X = 2) = {}^3C_2(0.5)^2(0.5)^1 = 0.375$$

$${}^nC_r p^r q^{n-r}$$

Example

- Mean:

$$\mu = np = 3(0.5) = 1.5$$

- Variance:

$$\sigma^2 = npq = 3(0.5)(0.5) = 0.75$$

- Standard deviation:

$$\sigma = \sqrt{npq} = \sqrt{0.75} = 0.8660$$

Another example

- A die is rolled 3 times.
 - ▣ Find the probability that we observe numbers 1 or 2 two times.
 - ▣ Find the mean and variance of the number of times 1 or 2 are observed.
- Let X be the number of times 1 or 2 are observed. X has a binomial distribution.
- Number of trials: $n = 3$.
- “Success” is observing 1 or 2.
- Probability of success and failure:

$$p = \frac{2}{6} = \frac{1}{3}, \quad q = 1 - p = \frac{2}{3}$$

- Probability $X = 2$:

$$P(2) = {}^3C_2 \left(\frac{1}{3}\right)^2 \left(\frac{2}{3}\right)^1 = 0.2222$$

Another example

□ Mean:

$$\mu = np = 3 \left(\frac{1}{3} \right) = 1$$

□ Variance:

$$\sigma^2 = npq = 3 \left(\frac{1}{3} \right) \left(\frac{2}{3} \right) = \frac{2}{3}$$

Exercise

- Forty percent of prison inmates were unemployed when they entered prison. If 5 inmates are randomly selected, find these probabilities:
- a) Exactly 3 were unemployed.
 - b) At most 4 were unemployed.
 - c) At least 3 were unemployed.
 - d) Fewer than 2 were unemployed.

Let X be the number of inmates that were unemployed out of the five inmates.

X has a binomial distribution with

$$n = 5, \quad p = 0.4, \quad q = 0.6$$

X can take values 0, 1, 2, 3, 4, 5

$$a) P(X=3) = {}^n C_x p^x q^{n-x} = {}^5 C_3 (0.4)^3 (0.6)^2$$

$$= 0.2304$$

$$b) P(X \leq 4) = 1 - P(5) = 1 - {}^5 C_5 (0.4)^5 (0.6)^0$$

$$= 0.9898 \quad (\text{or } P(0) + P(1) + P(2) + P(3) + P(4))$$

$$c) P(X \geq 3) = P(3) + P(4) + P(5)$$

$$= {}^5 C_3 (0.4)^3 (0.6)^2 + {}^5 C_4 (0.4)^4 (0.6)^1$$

$$+ {}^5 C_5 (0.4)^5 (0.6)^0$$

$$= 0.3174$$

$$d) P(X < 2) = P(0) + P(1) = {}^5 C_0 (0.4)^0 (0.6)^5 + {}^5 C_1 (0.4)^1 (0.6)^4$$

$$= 0.3370$$

Exercise

- Thirty-two percent of adult Internet users have purchased groceries online. For a random sample of 200 adult Internet users, find the mean, variance, and standard deviation for the number who have purchased groceries online.

X = number of adults who have purchased groceries online, out of 200 internet users.

X has a binomial distribution with
 $n = 200$, $p = 0.32$, $q = 0.68$

$$\mu = np = 250 \times 0.32 = 64$$

$$\sigma^2 = npq = 250 \times 0.32 \times 0.68 = 43.52$$

$$\sigma = \sqrt{43.52} = 6.597$$

Poisson distribution

Poisson experiment

- A **Poisson experiment** is a probability experiment that satisfies the following requirements:
 - ▣ The random variable X is the number of occurrences of an event over some interval (i.e., length, area, volume, period of time, etc.).
 - ▣ The occurrences occur randomly.
 - ▣ The occurrences are independent of one another.
 - ▣ The average number of occurrences over an interval is known.

- Note that $X \geq 0$ and can take the values 0, 1, 2, ...

Poisson distribution

- Let X be the outcome of a Poisson experiment.
- Then the distribution of X is a **Poisson distribution**.
- Example:
 - ▣ Number of patients admitted in a hospital in a day.
 - ▣ Number of customers in 5 hours interval.
 - ▣ Number of typographical error in a page.

Probability distribution for Poisson

- Denote λ be the mean number of occurrences in that interval.

- Then probability distribution of X :

$$P(X = x) = \frac{\lambda^x e^{-\lambda}}{x!}$$

for $x = 0, 1, 2, \dots$

- Mean:


$$\mu = \lambda$$

- Variance and standard deviation:

$$\sigma^2 = \lambda,$$

$$\sigma = \sqrt{\lambda}$$

Probability distribution for Poisson



Example

- A sales firm receives, on average, 3 calls per hour on its toll-free number. For any given hour, find the probability that it receives exactly 5 calls.
- Let X be the number calls received in an hour. X has a Poisson distribution.
- Mean number of calls per hour, $\lambda = 3$.
- Probability receiving exactly 5 calls:

$$P(5) = \frac{3^5 e^{-3}}{5!} = 0.1008$$

$$\frac{\lambda^x e^{-\lambda}}{x!}$$

Another example

- A sales firm receives, on average, 3 calls per hour on its toll-free number. Find the probability that it receives exactly 5 calls in a two-hour period.
- Let X be the number calls received in a two-hour period. X has a Poisson distribution.
- Mean number of calls in a two-hour period:

$$\lambda = \text{mean number per hour} \times 2 = 3 \times 2 = 6$$

- Probability receiving exactly 5 calls:

$$P(5) = \frac{6^5 e^{-6}}{5!} = 0.1606$$

Exercise

- A recent study of robberies for a certain geographic region showed an average of 1 robbery per 20,000 people. In a city of 80,000 people, find the probability of the following.
- a) 0 robberies
 - b) 1 robbery
 - c) 2 robberies

X = number of robberies in a city of 80,000 people

X has a Poisson distribution

$$\lambda = \frac{1}{20,000 \text{ people}} \times 80,000 \text{ people} = 1 \times 4 = 4$$

$$a) P(0) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{4^0 e^{-4}}{0!} = 0.01832$$

$$b) P(1) = \frac{4^1 e^{-4}}{1!} = 0.07326$$

$$c) P(2) = \frac{4^2 e^{-4}}{2!} = 0.1465$$

Exercise

- The mean number of accidents per month at a certain intersection is three. Find the probability that in any given month,
- a) No accidents will occur.
 - b) At least one accident will occur.
 - c) Less than three accident will occur.

X = number of accidents in a month

X has a Poisson distribution with

$$\lambda = 3$$

$$a) P(X=0) = \frac{\lambda^x e^{-\lambda}}{x!} = \frac{3^0 e^{-3}}{0!} = 0.04979$$

$$b) P(X \geq 1) = 1 - P(X=0) = 1 - 0.04979 \\ = 0.9502$$

$$c) P(X < 3) = P(0) + P(1) + P(2) \\ = \frac{3^0 e^{-3}}{0!} + \frac{3^1 e^{-3}}{1!} + \frac{3^2 e^{-3}}{2!} \\ = 0.4232$$

Geometric distribution

Geometric distribution

- A **geometric experiment** is a probability experiment such that:
 - ▣ Each trial has two outcomes, either success or failure
 - ▣ The outcomes are independent of each other
 - ▣ The probability of a success is the same for each trial
 - ▣ The experiment continues until a success is obtained.

- Geometric distribution is related to the binomial.
 - ▣ Binomial random variable counts number of successes in n trials.
 - ▣ Geometric random variable counts number of trials until success.

- Note: in some books, the geometric r.v. may refer to the number of failures until success

Probability distribution for geometric

- Probability distribution:

$$P(X = x) = (1 - p)^{x-1}p$$

for $x = 1, 2, 3, \dots$


- Mean:

$$\mu = \frac{1}{p}$$

- Variance:

$$\sigma^2 = \frac{1 - p}{p^2}$$

Probability distribution for geometric



Example

- Suppose that the probability that an applicant for a driver's license will pass the road test on any given try is 0.75, and assume the tests are independent.
- Let X be the number of road test taken by an applicant until they pass the test.
- In this case, X can take values 1, 2, 3, ... and X follows the geometric distribution with $p = 0.75$.
- For example, the probability that it will take an applicant passing the test for the first time at the 4th attempt is

$$P(X = 4) = (0.25)^3 0.75 = 0.01171$$

- Additionally, the mean number of attempts it will take for an applicant to pass the test is $\frac{1}{p} = 1.333$

Exercise

- The probability that you will make a sale on any given telephone call is 0.19. Find the probability that you:
- a) make your first sale on the fifth call.
 - b) make your first sale on the first, second, or third call.
 - c) do not make a sale on the first three calls.

X = number of call until the first sale

X has a geometric distribution with $p = 0.19$

$$\begin{aligned} \text{a) } P(X=5) &= (1-p)^{n-1} p = (1-0.19)^4 (0.19) \\ &= (0.81)^4 (0.19) \\ &= 0.08179 \end{aligned}$$

$$\begin{aligned} b) \quad P(1 \leq X \leq 3) &= P(1) + P(2) + P(3) \\ &= (0.81)^0(0.19) + (0.81)^1(0.19) \\ &\quad + (0.81)^2(0.19) \\ &= 0.4686 \end{aligned}$$

$$\begin{aligned} c) \quad P(X > 3) &= 1 - P(X \leq 3) \\ &= 1 - P(1) - P(2) - P(3) \\ &= 1 - 0.4686 \\ &= 0.5314 \end{aligned}$$

Summary

- This chapter introduces discrete random variables.
- We first introduced what random variables are.
- Then we focused on discrete random variables including
 - ▣ Probability distribution
 - ▣ Distribution function
 - ▣ Mean
 - ▣ Variance
- We looked at three special case of discrete random variables:
 - ▣ Binomial distribution – number of successes in n trials.
 - ▣ Poisson distribution – number of events in an interval.
 - ▣ Geometric distribution – number of trials until success