

Literature review and thesis proposal  
MRes. Neurotechnology

---

**Investigating plasticity in  
Cortico-Basal Ganglia-Thalamus models  
to improve stimulation-based treatments**

---

Aaron Panaiteanu  
(*CID*: 02054726)

**Supervisor:**  
Dr. Hayriye Cagnan

# Contents

<b>1</b>	<b>Introduction / Abstract?</b>	<b>2</b>
<b>2</b>	<b>Background and literature review</b>	<b>2</b>
2.1	Parkinson's Disease . . . . .	2
2.2	Deep brain stimulation: theory and practice . . . . .	2
2.3	Stimulating at the right time . . . . .	2
2.4	Plasticity to recover network states . . . . .	3
2.5	Neuron-level vs. Mean-field models . . . . .	3
2.5.1	Integrate-And-Fire . . . . .	3
2.5.2	Hodgkin-Huxley . . . . .	4
2.5.3	Rubin-Terman . . . . .	4
2.5.4	Jansen . . . . .	4
2.6	**Other ways of improving stimulation-based treatments** . . . . .	5
<b>3</b>	<b>Project Plan</b>	<b>5</b>
3.1	Aims . . . . .	5
3.2	Methodolgy . . . . .	5
3.3	Timeline . . . . .	5
	<b>References</b>	<b>6</b>
	<b>Appendices</b>	<b>8</b>
A	First appendix . . . . .	8
B	Second appendix . . . . .	8

# 1 Introduction / Abstract?

\*\*Stylistically, I would prefer to do an abstract here and breakdown the concepts in the following section\*\*

## 2 Background and literature review

\*\*start this off nicely with a diagram of the research gap\*\*

### 2.1 Parkinson's Disease

Parkinson's Disease (PD) is characterized by the loss of dopaminergic neurons and the presence of *Lewy Bodies* (LB) in the *Substantia Nigra pars Compacta* (SNc) [Del Rey et al., 2018]. This dopaminergic depletion in the SNc leads to striatal dopamine deficiency, which in turn leads to excessive synchronization in the basal ganglia. Hyper-synchrony is a hallmark of the PD [Hammond et al., 2007, Helmich et al., 2012], and thus, can be a promising therapeutic target.

### 2.2 Deep brain stimulation: theory and practice

Tremor is primarily generated in the Cortical-Basal Ganglia-Thalamic (CBGT) network. Thus, deep brain stimulation (DBS) in this region emerged as an effective tool in the treatment of advanced PD patients [Del Rey et al., 2018]. DBS works through the implantation of electrodes in particular regions of the brain. The intuition behind this method is that it enables stimulation driven decorrelation of the different basal ganglia nuclei, pushing it to a non-pathological state. There is significant evidence that performing DBS in the STN or the GPi, can have major quality of life improvements in PD patients, reducing tremor and restoring motor control [Rodriguez-Oroz et al., 2005]. DBS does, however have significant limitations.

Firstly, like any highly-invasive procedure, the implantation procedure has many risks. Advances in electrode technology and surgical techniques have reduced some of these risks. \*cite\*

Secondly, patient symptoms are only ameliorated while the stimulation is on. This means that the device needs to be on continuously, which can induce numerous side effects in patients \*\*citeee\*\* and also creates the need for a continuous power supply.

Finally, the mechanism through which DBS reduces tremors is not fully understood yet, there are many questions left to answer. For instance, why does DBS in PD get effective results around 130 Hz, when tremor frequency is on average around 20 Hz \*\*check this\*\*.

Therefore, several areas of research emerge in improving stimulation-based treatments of PD: finding non-invasive alternatives [Saturnino et al., 2017, Schwab et al., 2020], eliciting lasting, plastic changes in CBGT circuit, closing the loop and coupling stimulation to the presence of patient symptoms [Beudel et al., 2018], and improving stimulation patterns to harness the exploit the rhythmic patterns activity in the basal ganglia [Cagnan et al., 2017, West et al., 2022].

### 2.3 Stimulating at the right time

The Parkinsonian state is modulated by the interplay between four pathways in the CBGT: the **hyperdirect** excitatory cortico-subthalamic pathway, the **direct** inhibitory striato-pallidal pathway, the **indirect** inhibitory subthalamic-pallido pathway and the inhibitory **pallido-subthalamic** (PS) pathway. In the Parkinsonian state the hyperdirect pathway is down-regulated, while pallido-subthalamic pathway is upregulated, leading to a shift in beta band oscillatory activity from sub-band  $\beta_1$  to sub-band  $\beta_2$ , which correlates to hyper-synchrony in the basal ganglia [West et al., 2022]. \*\* add diagram \*\*

**\*\*important\*\*** [Caglan et al., 2017] [Beudel et al., 2018] [West et al., 2022]

## 2.4 Plasticity to recover network states

Neuroplasticity can be defined as "the ability of the nervous system to change its activity in response to intrinsic or extrinsic stimuli by reorganizing its structure, functions, or connections" [Mateos-Aparicio and Rodríguez-Moreno, 2019]. Plasticity occurs at different scales in the brain, ranging from molecular to neural networks. The focus of this project is synaptic plasticity, which occurs between a pre-synaptic and a pos-synaptic neuron. While, the mechanisms of plasticity have been thoroughly studied **\*\*cite\*\***, this understanding is just starting to find its way into clinical applications in neurorehabilitation (e.g. stroke, trauma, spinal cord injury [Cramer et al., 2011]) and neuroprostheses [Lebedev and Nicolelis, 2017].

**\*\*this should be relevant as well\*\*** [Schwab et al., 2020]

**\*\*at this point i should introduce STDP\*\***

## 2.5 Neuron-level vs. Mean-field models

Neuronal dynamics models are defined by defined by rules (ODEs) of how voltage in an unit changes over time in response to voltage in the network and external current (e.g. noise, stimulation current). The question then becomes what is an unit? There are two broad categories branching out of this question: An unit can be an individual neuron or a cluster of neurons. This section introduces some common types of models, justifying the direction chosen for this project.

**Neuron models** (or spiking models) work by combining networks of units that individually simulate neuronal action potentials (APs). These networks can then be organized into different nuclei (e.g. STN) and coerced to respect excitatory or inhibitory connections between nuclei. Within this class of models, different schemes for modeling neuronal dynamics exist, such as Integrate-And-Fire (IF) [Gerstner et al., 2014a] and Hodgkin-Huxley [Hodgkin et al., 1952, Gerstner et al., 2014b].

**Mean-field models** simulate the local field potentials (LFPs) of clusters of neurons and connect these clusters with population-level connectivity rules. (e.g. [Jansen and Rit, 1995])

### 2.5.1 Integrate-And-Fire

IF models treat action potentials as events. This reduction is justified by the fact that the shape of APs is always *approximately* the same, meaning they *cannot* convey information. [Gerstner et al., 2014a] describe a leaky IF model is represented by an electrical circuit with a resistor( $R$ ) and a capacitor( $C$ ) in parallel ( $I(t) = I_R + I_C$ ), combined with a reset condition when the potential exceeds threshold  $\theta$ . This can be modeled by,

$$\tau_m \dot{u} = -[u(t) - u_{rest}] + RI(t), \quad (1)$$

Together with the reset condition,

$$\lim_{\delta \rightarrow 0; \delta > 0} u(t^f + \delta) = u_r, \quad (2)$$

$$t^f = \{t | u(t) = \theta\}. \quad (3)$$

Where  $u$  is the membrane voltage,  $u_{rest}$  is the resting potential,  $u_r$  is the reset potential,  $\tau_m = RC$  is the time constant, and  $I$  is external current.

These equations describe the update rules in a single unit. Connectivity rules between units can be added to form neural networks. These connections can then be strengthened or weakened depending on spike timings between units, modeling STDP. [Shupe and Fetz, 2021] simulate plasticity in cortical columns; [Kromer et al., 2023] model plasticity between basal ganglia nuclei both using a IF model.

### 2.5.2 Hodgkin-Huxley

The Hodgkin-Huxley (HH) model [Hodgkin et al., 1952] simulates membrane potential by considering the dynamics of different ion conductances. Each ion has a reversal potential at which the force of the electrical and chemical gradients across the membrane even out. These can be modeled in an electrical circuit as a battery and the total channels of a particular ion can be modeled as a resistor. Each modelled ion has a parallel branch in the corresponding electrical circuit. The following equations adapted from [Gerstner et al., 2014b] describe the HH model:

$$I(t) = I_C(t) + \sum_k I_k(t), \quad (4)$$

Where the current of each ion  $k$  can be described in terms of the open probability of a  $k$  channel,  $p_k$  and the maximum conductance when all  $k$  channels are open,  $g_k$ :

$$I_k = g_k p_k (u - E_k). \quad (5)$$

In particular Hodgkin and Huxley modeled  $Na^+$ ,  $K^+$  and a leak  $L$  currents:

$$C\dot{u} = g_{Na}m^3h(E_{Na} - u) + g_Kn^4(E_K - u) + g_L(E_L - u), \quad (6)$$

Where  $m$ ,  $h$  and  $n$  are open probabilities for subunits of  $Na$  or  $K$  channels governed by:

$$\dot{x} = -\frac{1}{\tau_m(u)}[x - x_0(u)], \quad x \in \{m, h, n\}. \quad (7)$$

As in the IF model these rules describe the dynamics of a single unit, which can subsequently be connected to model a networks of neurons (e.g. [Terman et al., 2002]) and augmented to include plasticity rules [Borges et al., 2016].

### 2.5.3 Rubin-Terman

[Terman et al., 2002] expanded on the HH model to create a subthalamopallido network of the basal ganglia, where GPe and STN neurons are modeled according following equation,

$$C\dot{u} = -I_L - I_K - I_{Na} - I_T - I_{Ca} - I_{AHP} - I_{pre \rightarrow post} \quad (8)$$

Where leak,  $K^+$ ,  $Na^+$  and high-threshold  $Ca^+$  current [Song et al., 2000], are described in HH-like equations, and low threshold calcium current  $I_T$  is different for GPe and STN neurons.  $I_{AHP}$  represents calcium activated, voltage independent afterhyperpolarization (AHP)  $K^+$  current.  $I_{pre \rightarrow post}$  is synaptic current dependent on the nuclei the pre and post neurons belong to (either GPe or STN).

**\*\*BG\*\*** [Terman et al., 2002, Rubin and Terman, 2004, Rubin et al., 2012]

**\*\*plasticity\*\*** [Madadi Asl et al., 2022]

### 2.5.4 Jansen

**\*\*plasticity\*\*** [Duchet et al., 2023]

[Jansen and Rit, 1995] [West et al., 2022]

## 2.6 **\*\*Other ways of improving stimulation-based treatments\*\***

**\*\*stimulation parameter optimizations, closing the loop (e.g. aDBS [Beudel et al., 2018])\*\***  
*Maybe this can be folded into the stimulating at the right time part since they are pretty closely related. Also should mention the idea that these methods are not mutually exclusive, meaning that in principle they could be combined, adaptive stim closing the loop, with inducing plastic changes as target*

## 3 Project Plan

### 3.1 Aims

1. Model neuroplasticity in a Parkinsonian CGBT network
2. Investigate the viability of harnessing plasticity to remove the system from the pathological state and analyze the dynamics that follow  
**\*\*here i care about things like for how long and how does the network change. To what degree can we induce changes etc.\*\***  
**\*\*should look into viable timescales + noise analyze how long does it take to rebound to pathological state\*\***
3. Try to link potential results to potential stimulation protocols?

### 3.2 Methodolgy

**\*\*How indeed?\*\*** HH/IF Parkinsoni model + plasticity rules, trying different stimulation-based protocols (link with experimental data?)  
**\*\*look into how to tune IF model parameters from experimental data\*\***

### 3.3 Timeline

**\*\*Gant chart thingy\*\***

## References

- [Beudel et al., 2018] Beudel, M., Cagnan, H., and Little, S. (2018). Adaptive brain stimulation for movement disorders. *Current Concepts in Movement Disorder Management*, 33:230–242.
- [Borges et al., 2016] Borges, R., Borges, F. S., Lameu, E. L., Batista, A. M., Iarosz, K. C., Caldas, I. L., Viana, R. L., and Sanjuán, M. A. (2016). Effects of the spike timing-dependent plasticity on the synchronisation in a random hodgkin–huxley neuronal network. *Communications in Nonlinear Science and Numerical Simulation*, 34:12–22.
- [Cagnan et al., 2017] Cagnan, H., Pedrosa, D., Little, S., Pogosyan, A., Cheeran, B., Aziz, T., Green, A., Fitzgerald, J., Foltynie, T., Limousin, P., et al. (2017). Stimulating at the right time: phase-specific deep brain stimulation. *Brain*, 140(1):132–145.
- [Cramer et al., 2011] Cramer, S. C., Sur, M., Dobkin, B. H., O’Brien, C., Sanger, T. D., Trojanowski, J. Q., Rumsey, J. M., Hicks, R., Cameron, J., Chen, D., et al. (2011). Harnessing neuroplasticity for clinical applications. *Brain*, 134(6):1591–1609.
- [Del Rey et al., 2018] Del Rey, N. L.-G., Quiroga-Varela, A., Garbayo, E., Carballo-Carbajal, I., Fernández-Santiago, R., Monje, M. H., Trigo-Damas, I., Blanco-Prieto, M. J., and Blesa, J. (2018). Advances in parkinson’s disease: 200 years later. *Frontiers in neuroanatomy*, 12:113.
- [Duchet et al., 2023] Duchet, B., Bick, C., and Byrne, Å. (2023). Mean-field approximations with adaptive coupling for networks with spike-timing-dependent plasticity. *Neural Computation*, 35(9):1481–1528.
- [Gerstner et al., 2014a] Gerstner, W., Kistler, W. M., Naud, R., and Paninski, L. (2014a). *Neuronal dynamics: From single neurons to networks and models of cognition*, chapter 1.3, pages 10–19. Cambridge University Press.
- [Gerstner et al., 2014b] Gerstner, W., Kistler, W. M., Naud, R., and Paninski, L. (2014b). *Neuronal dynamics: From single neurons to networks and models of cognition*, chapter 2.2, pages 31–41. Cambridge University Press.
- [Hammond et al., 2007] Hammond, C., Bergman, H., and Brown, P. (2007). Pathological synchronization in parkinson’s disease: networks, models and treatments. *Trends in neurosciences*, 30(7):357–364.
- [Helmich et al., 2012] Helmich, R. C., Hallett, M., Deuschl, G., Toni, I., and Bloem, B. R. (2012). Cerebral causes and consequences of parkinsonian resting tremor: a tale of two circuits? *Brain*, 135(11):3206–3226.
- [Hodgkin et al., 1952] Hodgkin, A. L., Huxley, A. F., and Katz, B. (1952). Measurement of current-voltage relations in the membrane of the giant axon of loligo. *The Journal of physiology*, 116(4):424.
- [Jansen and Rit, 1995] Jansen, B. H. and Rit, V. G. (1995). Electroencephalogram and visual evoked potential generation in a mathematical model of coupled cortical columns. *Biological cybernetics*, 73(4):357–366.
- [Kromer et al., 2023] Kromer, J. A., Bokil, H., and Tass, P. A. (2023). Synaptic network structure shapes cortically evoked spatio-temporal responses of stn and gpe neurons in a computational model. *Frontiers in Neuroinformatics*, 17:1217786.

- [Lebedev and Nicolelis, 2017] Lebedev, M. A. and Nicolelis, M. A. (2017). Brain-machine interfaces: from basic science to neuroprostheses and neurorehabilitation. *Physiological reviews*, 97(2):767–837.
- [Madadi Asl et al., 2022] Madadi Asl, M., Asadi, A., Enayati, J., and Valizadeh, A. (2022). Inhibitory spike-timing-dependent plasticity can account for pathological strengthening of pallido-subthalamic synapses in parkinson’s disease. *Frontiers in Physiology*, 13:915626.
- [Mateos-Aparicio and Rodríguez-Moreno, 2019] Mateos-Aparicio, P. and Rodríguez-Moreno, A. (2019). The impact of studying brain plasticity. *Frontiers in cellular neuroscience*, 13:66.
- [Rodriguez-Oroz et al., 2005] Rodriguez-Oroz, M. C., Obeso, J., Lang, A., Houeto, J.-L., Pollak, P., Rehncrona, S., Kulisevsky, J., Albanese, A., Volkmann, J., Hariz, M., et al. (2005). Bilateral deep brain stimulation in parkinson’s disease: a multicentre study with 4 years follow-up. *Brain*, 128(10):2240–2249.
- [Rubin et al., 2012] Rubin, J. E., McIntyre, C. C., Turner, R. S., and Wichmann, T. (2012). Basal ganglia activity patterns in parkinsonism and computational modeling of their downstream effects. *European Journal of Neuroscience*, 36(2):2213–2228.
- [Rubin and Terman, 2004] Rubin, J. E. and Terman, D. (2004). High frequency stimulation of the subthalamic nucleus eliminates pathological thalamic rhythmicity in a computational model. *Journal of computational neuroscience*, 16:211–235.
- [Saturnino et al., 2017] Saturnino, G. B., Madsen, K. H., Siebner, H. R., and Thielscher, A. (2017). How to target inter-regional phase synchronization with dual-site transcranial alternating current stimulation. *Neuroimage*, 163:68–80.
- [Schwab et al., 2020] Schwab, B. C., König, P., and Engel, A. K. (2020). Spike-timing-dependent plasticity can account for aftereffects of dual-site transcranial alternating current stimulation. *bioRxiv*.
- [Shupe and Fetz, 2021] Shupe, L. and Fetz, E. (2021). An integrate-and-fire spiking neural network model simulating artificially induced cortical plasticity. *Eneuro*, 8(2).
- [Song et al., 2000] Song, W.-J., Baba, Y., Otsuka, T., and Murakami, F. (2000). Characterization of  $Ca^{2+}$  channels in rat subthalamic nucleus neurons. *Journal of Neurophysiology*, 84(5):2630–2637.
- [Terman et al., 2002] Terman, D., Rubin, J. E., Yew, A., and Wilson, C. (2002). Activity patterns in a model for the subthalamopallidal network of the basal ganglia. *Journal of Neuroscience*, 22(7):2963–2976.
- [West et al., 2022] West, T. O., Magill, P. J., Sharott, A., Litvak, V., Farmer, S. F., and Cagnan, H. (2022). Stimulating at the right time to recover network states in a model of the cortico-basal ganglia-thalamic circuit. *PLOS Computational Biology*, 18(3):1–32.



## Appendices

A First appendix

B Second appendix