

ALGEBRA

Arithmetic Operations

$$a(b+c) = ab+ac$$
$$\frac{a+c}{b} = \frac{a}{b} + \frac{c}{b}$$

$$\frac{a}{b} + \frac{c}{d} = \frac{ad+bc}{db}$$
$$\frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$$

Exponents and Radicals

$$x^m x^n = x^{m+n}$$
$$(x^m)^n = x^{mn}$$
$$(xy)^n = x^n + y^n$$
$$x^{\frac{1}{n}} = \sqrt[n]{x}$$
$$\sqrt[n]{xy} = \sqrt[n]{x} \sqrt[n]{y}$$

$$\frac{x^m}{x^n} = x^{m-n}$$
$$x^{-n} = \frac{1}{x^n}$$
$$\left(\frac{x}{y}\right)^n = \frac{x^n}{y^n}$$
$$x^{\frac{m}{n}} = \sqrt[n]{x^m} = (\sqrt[n]{x})^m$$
$$\sqrt[n]{\frac{x}{y}} = \frac{\sqrt[n]{x}}{\sqrt[n]{y}}$$

Factoring Special Polynomials

$$x^2 - y^2 = (x + y)(x - y)$$

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2)$$
$$x^3 - y^3 = (x - y)(x^2 + xy + y^2)$$

Binomial Theorem

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k$$

$$(x + y)^2 = x^2 + 2xy + y^2$$
$$(x - y)^2 = x^2 - 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$
$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

Quadratic Formula

If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Inequalities and Absolute Value

If $a < b$ and $b < c$, then $a < c$

If $a < b$, then $a + c < b + c$

If $a < b$ and $c > 0$, then $ca < cb$

If $a < b$ and $c < 0$, then $ca > cb$

If $a > 0$, then

$|x| = a$ means $x = a$ or $x = -a$

$|x| < a$ means $-a < x < a$


$|x| > a$ means $x > a$ or $x < -a$

GEOMETRY


Geometric Formulas

- Formulas for:
- Area A
 - Circumference C
 - Volume V


Triangle

$$A = \frac{1}{2}bh$$
$$A = \frac{1}{2}ab \sin(\theta)$$


Circle

$$A = \pi r^2$$
$$C = 2\pi r$$


Sector of Circle

$$A = \frac{1}{2}r^2\theta$$
$$s = r\theta \ (\theta \text{ in radians})$$


0.1. Trigonometric Functions

Function	Definition	Reciprocal
$\sin(x)$	$\frac{\text{opposite}}{\text{hypotenuse}}$	$\frac{1}{\sin(x)} = \csc(x)$
$\cos(x)$	$\frac{\text{adjacent}}{\text{hypotenuse}}$	$\frac{1}{\cos(x)} = \sec(x)$
$\tan(x)$	$\frac{\sin(x)}{\cos(x)} = \frac{\text{opposite}}{\text{adjacent}}$	$\frac{1}{\tan(x)} = \cot(x)$
$\csc(x)$	$\frac{\sin(x)}{\cos(x)} = \frac{\text{hypotenuse}}{\text{opposite}}$	$\frac{1}{\csc(x)} = \sin(x)$
$\sec(x)$	$\frac{\text{hypotenuse}}{\text{adjacent}}$	$\frac{1}{\sec(x)} = \cos(x)$
$\cot(x)$	$\frac{\cos(x)}{\sin(x)} = \frac{\text{adjacent}}{\text{opposite}}$	$\frac{1}{\cot(x)} = \tan(x)$

Contents

0.1. Trigonometric Functions	1
1. Matrix	16
1.1. Matrix Vector Product	16
1.2. Matrix Multiplication	16
1.3. Transpose	17
2. Matrix Factorization	17
2.1. LU Decomposition	17
2.2. QR Decomposition	17
2.3. Cholesky Decomposition	17
2.4. Singular Value Decomposition (SVD)	17
3. Vectors	20
3.1. Real Coordinate Spaces	20
3.2. Vector Operations	20
3.2.1. Vector Addition	20
3.2.2. Vector Subtraction	21
3.2.3. Scalar Multiplication	22
3.3. Unit Vector	23
3.4. Parametric Representation of line	24
3.5. Vector Spaces	27
4. Matrices	28
4.1. Matrix-Vector Products	28
4.2. Null Space	29
4.2.1. Column Space	30
4.2.2. Dimension of a Subspace	31
4.2.3. Nullity	31
4.2.4. Rank	31
4.2.5. Matrix Representation of Systems of Equations	31
4.3. Matrix Multiplication	32
5. Linear Combinations	33
6. Span	33
7. Linear Independence	36
8. Subspace	38
9. Basis	41
9.1. Vector Dot Product	42
9.1.1. Magnitude (Length)	42
9.1.2. Properties	42
9.1.2.1. Commutative	42
9.1.2.2. Distributive	42
9.1.2.3. Associativity	43
9.1.3. Cauchy-Schwarz Inequality	43
9.1.4. Vector Triangle Inequality	45
9.2. Angles Between Vectors	45
9.2.1. Plane in \mathbb{R}^3	48
9.2.2. Point Distance to Plane	50
9.2.3. Distance Between Planes	50
9.2.4. Cross Product	50
9.2.5. Proof: Relationship Between Cross Product and Sin of Angle	51

9.2.6. Dot and Cross Products	51
9.3. Row Echelon Form (REF)	51
9.3.1. Solution Types in Linear Systems: Unique, Infinite, or None	53
9.3.2. Special Cases	54
10. Matrices	55
10.1. Matrix-Vector Products	55
10.2. Null Space	56
10.2.1. Column Space	57
10.2.2. Dimension of a Subspace	58
10.2.3. Nullity	58
10.2.4. Rank	58
10.2.5. Matrix Representation of Systems of Equations	58
10.3. Matrix Multiplication	59
10.4. Linear Transformation	60
10.4.1. Functions	60
10.5. Vector Transformation	61
10.6. Linear Transformation	62
10.7. Matrix Vector Products	63
10.8. Linear transformations as matrix vector products	66
10.9. Image of a subset under transformation	67
10.10. Image of a transformation	68
10.11. Preimage of a set	70
10.11.1. Kernel of a Transformation	71
10.11.2. Kernel and Null Space	72
10.12. Sum and Scalar Multiples of Linear Transformation	72
10.12.1. Sum	72
10.12.2. Scalar Multiplication	72
10.13. Projection	73
10.14. Composition of Linear Transformations	77
10.15. Matrix Product	79
10.16. Matrix Product Associativity	80
10.17. Eigenvectors	80
10.17.1. Transformation	80
10.18. Eigenvalues	81
10.18.1. LU Decomposition	81
10.19. Solving Systems of Linear Equations	82
10.19.1. Gaussian Elimination	85
10.19.2. Substitution	87
10.19.3. Addition or Subtraction Method	87
11. Cheatsheet	90
11.1. Limits	90
11.2. Derivatives	91
11.3. Integrals	92
11.4. Differential	92
11.5. Product Rule \rightarrow Integration by Parts	95
11.6. Chain Rule \rightarrow u -Substitution	95
11.7. Quotient Rule \rightarrow Algebraic Manipulation / Substitution	95
12. Limits & Continuity	96
13. Properties of Limits	96

13.1. Continuous	96
13.1.1. Addition, Subtraction, Multiplication, Division	96
13.1.2. Constant	96
13.2. Non-continuous	96
13.3. Composite Functions	97
13.4. Limits by Direct Substitution	99
13.4.1. Limits of Piecewise Functions	99
13.4.2. Absolute Value	99
13.5. Limits by Factoring	99
13.6. Limits by Rationalizing	99
13.7. Continuity & Differentiability at a Point	99
13.8. Power Rule	100
13.9. Constant Rule	101
13.10. Constant Multiple Rule	101
13.11. Sum Rule	101
13.12. Difference Rule	101
13.13. Square Root	102
13.14. Derivative of a Polynomial	102
13.15. Sin	103
13.16. Cos	104
13.17. e^x	104
13.18. $\ln(x)$	105
13.19. Product Rule	105
13.20. Quotient Rule	106
13.20.1. $\tan(x)$	106
13.20.2. $\cot(x)$	107
13.20.3. $\sec(x)$	108
13.20.4. $\csc(x)$	109
13.21. Chain Rule	110
13.22. Implicit Differentiation	111
13.23. Derivatives of Inverse Functions	112
13.23.1. Derivative Inverse Sin	115
13.23.2. Derivative Inverse Cos	115
13.23.3. Derivative Inverse Tan	115
13.24. Inverse Functions	115
13.25. L'Hôpital's Rule	117
13.26. Mean Value Theorem	119
13.27. Extreme Value Theorem	120
13.27.1. Critical points	120
13.27.2. Global vs. Local Extrema	121
13.27.3. First and Second Derivative Tests	121
13.27.3.1. First Derivative Test	121
13.27.3.2. Second Derivative Test	122
13.28. Differentiation Rules	123
14. Probability Theory	129
14.1. Probability Axioms	129
14.1.1. Non-Negativity	129
14.1.2. Normalization	129
14.1.3. Additivity	129

14.2. Rules	129
14.2.1. Complement Rule	129
14.2.2. Multiplication Rule	129
14.2.3. Addition Rule	130
14.2.4. Conditional Probability	130
14.2.5. Law of Total Probability	130
14.2.6. Law of Large Numbers	131
14.2.7. Central Limit Theorem	131
14.3. Bayes Theorem	131
15. Descriptive Statistics	131
15.1. Central Tendency	131
15.1.1. Mean	131
15.1.2. Median	132
15.1.3. Mode	132
15.2. Dispersion	132
15.2.1. Range	132
15.2.2. Variance	133
15.2.3. Standard deviation	133
15.2.4. Interquartile Range (IQR)	134
16. Probability Distributions	134
16.1. Gaussian (Normal) distribution	134
16.2. t-Distribution	135
16.3. Binomial distribution	135
16.4. Poisson distribution	136
16.5. Exponential distribution	136
17. Functions	137
17.1. PDF (Probability Density Function)	137
17.2. PMF (Probability Mass Function)	137
17.3. CDF (Cumulative Distribution Function)	137
17.4. PPF (Percent-Point Function)	138
17.5. SF (Survival Function)	138
18. Error Metrics	138
18.1. MAE (Mean Absolute Error)	138
18.2. MSE (Mean Squared Error)	138
18.3. RMSE (Root Mean Squared Error)	139
18.4. MAPE (Mean Absolute Percentage Error)	139
18.5. R-squared	139
18.6. Adj R-squared	140
18.7. MSLE (Mean Squared Logarithmic Error)	140
18.8. Cross-Entropy Loss (Log Loss)	140
19. Hypothesis Testing	141
19.1. Hypotheses	141
19.1.1. Null (H_0)	141
19.1.2. Alternative (H_1 or H_a)	141
19.2. Error Types	141
19.2.1. Type I (α)	141
19.2.2. Type II (β)	141
19.3. t-Tests	141
19.3.1. One-sample	141

19.3.2. Independent	143
19.3.3. Paired	143
19.4. Chi-square tests	143
19.4.1. Goodness of Fit Test	143
19.4.2. Test of independence	144
19.5. ANOVA (Analysis of Variance)	144
19.5.1. One-way	144
19.5.2. Two-way	145
20. Regression Analysis:	145
20.1. Simple linear regression	145
20.2. Multiple regression	147
20.3. Logistic Regression	147
20.4. Model diagnostics	147
20.4.1. p-Values	147
20.4.2. F-Statistic	148
20.4.3. Confidence Intervals (CI)	149
21. Correlation	149
21.1. Pearson	149
21.2. Spearman's Rank	149
22. Non-Parametric Statistics	150
22.1. Mann-Whitney U	150
22.2. Wilcoxon Signed-Rank	150
22.3. Kolmogorov-Smirnov	150
22.4. Kruskal-Wallis	150
23. Time Series	151
23.1. SMA (Simple Moving Averages)	151
23.2. WMA (Weighted Moving Average)	152
23.3. Exponential Smoothing	152
23.4. Seasonal Decomposition	153
23.5. ARMA (AutoRegressive Moving Average)	154
23.6. ARIMA (AutoRegressive Integrated Moving Average)	154
24. Causes of variation	156
24.1. Common	156
24.2. Special	156
25. Design of Experiments (DOE):	156
26. Control Charts:	156
26.1. P-charts (Proportion)	156
26.2. NP-charts (Number Proportion)	157
26.3. C-charts (Count)	157
26.4. U-charts (Unit)	158
26.5. \bar{X} -chart	159
26.6. R-chart	159
27. Process Capability Analysis	159
27.1. C_p (Process Capability Index)	159
27.2. C_{pk} (Process Capability Index with Centering)	162
27.3. C_{pm} (Taguchi Capability Index)	164
27.4. P_p (Process Performance Index)	164
27.5. P_{pk} (Process Performance Index with Centering)	166
28. Inventory Management	167

28.1. Newsvendor	167
28.2. ABC Analysis	169
28.3. Fill Rate	170
28.4. (OCT) Order Cycle Time	170
28.5. ROP (Reorder Point)	170
28.6. XYZ Analysis	171
28.7. EOQ (Economic Order Quantity)	172
28.7.1. Perfect Order Rate	172
28.8. Safety Stock	172
29. Queuing Theory	173
29.1. M/M/1	173
30. Network Optimization	178
30.1. Shortest Path	178
30.2. Dijkstra's	178
30.3. Maximum Flow	179
30.4. Network Flow Optimization	179
30.5. Ford-Fulkerson	183
31. Optimization	186
32. Linear Programming	187
33. Integer Programming	187
34. Non-Linear Programming	189
34.1. Mathematical Programs	190
34.1.1. Formulation	190
34.1.2. Transformation	191
34.1.3. Constraints	191
34.1.4. Feasible Solutions	191
34.1.5. Feasible Region and Optimal Solution	192
34.1.6. Binding Constraint	192
34.1.7. Strict and Weak Inequalities	192
34.1.8. Graphical Approach	192
34.1.9. Types of LP	193
34.1.10. LP Formulation	193
34.2. IP (Integer Programming)	202
34.3. Selection and Logical Relations on Binary Variables $x_i \in \{0, 1\}$	202
34.3.1. Cardinality Constraints	202
34.3.2. Logical OR	202
34.3.3. Conditional Selection: "If X, then Y"	203
34.3.4. Conditional OR: "If not X, then select Y and Z"	204
34.3.5. XOR / Exclusive Conditions	204
34.4. Constraint Activation and Relaxation via Binary Indicators $z_i \in \{1, 0\}$	205
34.4.1. Base Case: At Least One of Two Constraints Satisfied	205
34.4.2. General Formulation for m Constraints with At Least k Satisfied	205
34.5. Fixed Charge Constraints (Setup Costs)	206
34.6. Facility Location	206
34.6.1. Set Covering Problem	207
34.6.2. Maximum Covering Problem	207
34.7. Machine Scheduling	213
34.7.1. Single Machine Serial Production	214
34.7.2. Multiple Parallel Machines	216

34.7.3. Flow Shop	217
34.7.4. Job Shop	217
34.8. Vehicle Routing	217
34.8.1. Traveling Salesperson	217
34.9. MIP (Mixed Integer Programming)	219
34.10. Non-Linear Programming	219
34.10.1. EOQ	219
34.10.2. Portfolio Optimization	221
34.11. Linearizing Maximum/Minimum Functions	221
34.12. Linearize Objective Function	222
34.13. Linearizing Products of Decision Variables	225
34.14. Simplex Method	225
34.14.1. Linear Algebra Review	225
34.14.2. Gaussian Elimination	226
34.14.3. Inverse	228
34.14.4. Linear Dependence and Independence	229
34.14.5. Extreme Points	230
34.14.6. Slack and Suplus	231
34.14.7. Standard Form	257
34.14.8. Simplex Method	258
34.15. Branch-and-Bound	261
34.16. Gradient Descent	266
34.16.1. Gradient	266
34.16.2. Algorithm	267
34.17. Newton's Method	275
34.18. Monte Carlo	277

Linear Algebra

Vector	$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$	
Scalar Vector Multiplication	$c \in \mathbb{R} \quad \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$ $c\vec{x} = \begin{bmatrix} cx_1 \\ cx_2 \\ \vdots \\ cx_n \end{bmatrix}$	
Dot Product	$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$ $\vec{x} \cdot \vec{y} = \sum_{i=1}^n x_i y_i$ $= x_1 y_1 + x_2 y_2 + \dots + x_n y_n$ $= \vec{x}^T \vec{y}$	
Cross Product (\mathbb{R}^3)	$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$ $\vec{c} = \vec{a} \times \vec{b}$ $\vec{c} = \begin{bmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{bmatrix}$	Returns a vector orthogonal to the two vectors
Vector Space	<p>Closure under addition</p> $\vec{u}, \vec{v} \in V \Rightarrow \vec{u} + \vec{v} \in V$ <p>Closure under scalar multiplication</p> $\vec{v} \in V \wedge c \in \mathbb{R} \Rightarrow c\vec{v} \in V$ <p>Commutativity of addition</p> $\vec{u} + \vec{v} = \vec{v} + \vec{u}$ <p>Associativity of addition</p> $(\vec{u} + \vec{v}) + \vec{w} = \vec{u} + (\vec{v} + \vec{w})$ <p>Additive identity</p> $\exists \mathbf{0} \in V \mid \vec{v} + \mathbf{0} = \vec{v}$ <p>Additive inverse</p> $\forall \vec{v} \in V \exists -\vec{v} \in V \mid \vec{v} + (-\vec{v}) = \mathbf{0}$ <p>Scalar multiplication (compatibility)</p> $a(b\vec{v}) = (ab)\vec{v}$ <p>Distributivity over vector addition</p> $a(\vec{u} + \vec{v}) = a\vec{u} + a\vec{v}$	

	Distributivity over scalar addition $(a + b)\vec{v} = a\vec{v} + b\vec{v}$ Multiplicative identity $1\vec{v} = \vec{v}$	
Subspace	Non-emptiness $\mathbf{0} \in V$ Closure under addition If $\vec{u}, \vec{v} \in V$, then $\vec{u} + \vec{v} \in V$ Closure under scalar multiplication If $\vec{v} \in V, c \in \mathbb{R}$, then $c\vec{v} \in V$	A subspace is a subset of a vector space that is itself a vector space, satisfying the same axioms as the original. If V is a vector space in \mathbb{R}^n , then the subspace U is always contained in \mathbb{R}^n , meaning $U \subseteq \mathbb{R}^n$
Vector Addition	$\vec{u} = (u_1, u_2, \dots, u_n)$ $\vec{v} = (v_1, v_2, \dots, v_n)$ $\vec{u} + \vec{v} = (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n)$	
Dot Product	$\vec{u} \cdot \vec{v} = \sum_{i=1}^n u_i v_i$	
Orthogonality	$\vec{u} \cdot \vec{v} = \mathbf{0}$	Angle between the two vectors is 90°
Angle between vectors	$\Theta = \arccos\left(\frac{\vec{u} \cdot \vec{v}}{\ \vec{u}\ _2 \cdot \ \vec{v}\ _2}\right)$	
L_1 Norm (Manhattan)	$\ \vec{u}\ _1 = \sum_{i=1}^n u_i $	
L_2 Norm (Euclidean)	$\ \vec{u}\ _2 = \sqrt{\sum_{i=1}^n u_i^2}$	
L_1 Distance (Manhattan)	$d(\vec{u}, \vec{v}) = \sum_{i=1}^n u_i - v_i $	
L_2 Distance (Euclidean)	$d(\vec{u}, \vec{v}) = \sqrt{\sum_{i=1}^n (u_i - v_i)^2}$	
Projection	$\text{proj}_{\vec{w}}(\vec{v}) = \frac{\vec{v} \cdot \vec{w}}{\vec{w} \cdot \vec{w}} \vec{w}$	
Linear Independence		<p>A set of vectors is linearly independent if no vector in the set can be written as a linear combination of the others</p> <p>A set of vectors $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ is linearly independent if the only solution to the equation</p> $c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_n \vec{v}_n = \mathbf{0}$ <p>is $c_1 = c_2 = \dots = c_n = 0$</p>
Transformation	$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ $T(\vec{v}) = A\vec{v}$ <ul style="list-style-type: none"> Additivity 	<ul style="list-style-type: none"> Surjective (onto) <p>Every element in B is the image of at least one element in A. The transformation covers the entire codomain.</p>

	$T(\vec{u} + \vec{v}) = T(\vec{u}) + T(\vec{v})$ <ul style="list-style-type: none"> Homogeneity $T(c\vec{u}) = cT(\vec{u})$	$\text{Range}(T) = B$ <ul style="list-style-type: none"> Injective (one-to-one) <p>Different inputs in A map to different outputs in B. The transformation is information-preserving – doesn't collapse distinct vectors together</p> $T(\vec{x}_1) = T(\vec{x}_2) \Rightarrow x_1 = x_2$ <p>Or equivalently:</p> $\ker(T) = \{\mathbf{0}\}$
Domain	$T : V \rightarrow W$ $\text{Domain}(T) = V$	Set of all input vectors V that the transformation acts on
Codomain	$T : V \rightarrow W$ $\text{Codomain}(T) = W$	Set of all possible output vectors W to which elements of the domain V are mapped under the transformation
Transpose	$\det(A) = \det(A^T)$ $(AB)^T = B^T A^T$ $(A^T)^{-1} = (A^{-1})^T$	
Image	$T : V \rightarrow W$ $\text{Im}(T) = \{\vec{w} \in W \mid \vec{w} = T(\vec{v}) \text{ for some } \vec{v} \in V\}$	<p>The image of a transformation $T : V \rightarrow W$ is the set of all possible outputs $T(v)$ for $v \in V$:</p> <ul style="list-style-type: none"> It is a subspace of the codomain W If T is represented by a matrix A, the image of T is the column space of A
Preimage	$T : V \rightarrow W$ $T^{-1}(\vec{w}) = \{\vec{v} \in V \mid T(\vec{v}) = \vec{w}\}$	The preimage of a transformation refers to the set of all elements in the domain that map to a particular element or subset in the codomain
Span	$\text{Span}(\{v_1, v_2, \dots, v_k\}) = \left\{ \sum_{i=1}^n c_i \vec{v}_i \mid c_i \in \mathbb{R} \right\}$	The span of a set of vectors is the collection of all possible linear combinations of those vectors
Composition	$T_1 : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad T_2 : \mathbb{R}^m \rightarrow \mathbb{R}^p$ $T_1(\vec{v}) = A\vec{v} \quad T_2(\vec{v}) = B\vec{v}$ $T \circ S(\vec{v}) = T_2(T_1(\vec{v}))$ $T_2 \circ T_1(\vec{v}) = B(A\vec{v}) = (BA)\vec{v}$ <ul style="list-style-type: none"> Additivity $(T_3 \circ T_2) \circ T_1 = T_3 \circ (T_2 \circ T_1)$ <ul style="list-style-type: none"> Homogeneity $T_2 \circ T_1(c\vec{u}) = c(T_2 \circ T_1)(\vec{u})$ <ul style="list-style-type: none"> Identity Transformation 	

	$I \circ T = T \qquad T \circ I = T$					
Column Space (Range)	$\text{Col}(A) = \{Ax \mid \vec{x} \in \mathbb{R}^n\}$ Or equivalently $A = [\vec{c}_1 \ \vec{c}_2 \ \dots \ \vec{c}_3]$ $\text{Col}(A) = \text{span}(\vec{c}_1, \vec{c}_2, \dots, \vec{c}_3)$	The column space (or range) of a matrix A is the set of all linear combinations of its columns				
Determinant	$\det(A)$	<p>The determinant of a square matrix A measure of the "scale factor" by which the matrix A transforms a space</p> <ul style="list-style-type: none">$\det(A) \neq 0$<ul style="list-style-type: none">A does not collapse the spaceA has full rankA's columns are linearly independentA is invertable$\det(A) = 0$<ul style="list-style-type: none">A collapses the space into lower dimensionA does not have full rankA's columns are linearly dependentA is non-invertable (singular)				
Invertibility	$\det(A) \neq 0 \implies \text{Invertible}$ $\det(A) = 0 \implies \text{Non-Invertible}$ $AA^{-1} = A^{-1}A = I_n$ $(AB)^{-1} = B^{-1}A^{-1}$ $(A^T)^{-1} = (A^{-1})^T$					
Basis	<p>Linear Independence</p> $c_1\vec{v}_1 + c_2\vec{v}_2 + \dots + c_k\vec{v}_k = \mathbf{0}$ $\implies c_1 = c_2 = \dots = c_k = 0$ <p>Spanning</p> $\forall \vec{v} \in V, \exists c_1, \dots, c_k \in \mathbb{R} \quad s.t.$ $\vec{v} = c_1\vec{v}_1 + \dots + c_k\vec{v}_k$	<ul style="list-style-type: none">A basis of a vector space V is a set of linearly independent vectors that span the spaceEvery vector in V can be uniquely written as a linear combination of the basis vectors <p>E.g.:</p>				
Dimension	$\dim(V)$	<p>Number of linearly independent vectors (basis) in a vector space V</p> <p>$V \subseteq \mathbb{R}^n$</p> <table><tr><td>$\dim(V) = 0$</td><td>$V = \{\mathbf{0}\}$</td></tr><tr><td>$\dim(V) = 1$</td><td>V is a line through the origin in \mathbb{R}^n</td></tr></table>	$\dim(V) = 0$	$V = \{\mathbf{0}\}$	$\dim(V) = 1$	V is a line through the origin in \mathbb{R}^n
$\dim(V) = 0$	$V = \{\mathbf{0}\}$					
$\dim(V) = 1$	V is a line through the origin in \mathbb{R}^n					

		<table><tr><td>$\dim(V) = 2$</td><td>V is a plane through the origin in \mathbb{R}^n</td></tr><tr><td>$\dim(V) = k$</td><td>V is a k-dimensional flat subspace of \mathbb{R}^n</td></tr><tr><td>$\dim(V) = n$</td><td>$V = \mathbb{R}^n$</td></tr></table> <ul style="list-style-type: none">\mathbb{R}^2 has dimension 2:<ul style="list-style-type: none">A basis: $\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$\mathbb{R}^3$ has dimension 3:<ul style="list-style-type: none">A basis: $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$	$\dim(V) = 2$	V is a plane through the origin in \mathbb{R}^n	$\dim(V) = k$	V is a k -dimensional flat subspace of \mathbb{R}^n	$\dim(V) = n$	$V = \mathbb{R}^n$
$\dim(V) = 2$	V is a plane through the origin in \mathbb{R}^n							
$\dim(V) = k$	V is a k -dimensional flat subspace of \mathbb{R}^n							
$\dim(V) = n$	$V = \mathbb{R}^n$							
Rank	$\text{Rank}(A) = \dim(\text{Col}(A)) = \dim(\text{Row}(A))$	<ul style="list-style-type: none">The rank of a matrix A is the dimension of its column space (or row space)Number of linearly independent columns (or rows)						
Eigen	$Ax = \lambda x, \quad x \neq \mathbf{0}$	<p>Set of all nonzero vectors \vec{x} such that when the transformation represented by matrix A is applied to \vec{x}, the result is a scaled version of \vec{x} itself</p> <p>These vectors lie along directions that are preserved by the transformation:</p> <ul style="list-style-type: none">$\lambda > 1$: stretched$0 < \lambda < 1$: shrunk$\lambda < 0$: flipped$\lambda = 1$: stay the same						
Null Space (kernel)	$\text{Null}(A) = \{ \vec{x} \in \mathbb{R}^n \mid A\vec{x} = \mathbf{0} \}$	The null space of a matrix A is the set of all input vectors that get mapped to the zero vector when you multiply them by A						
Identity Matrix	$I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}$							
Matrix Inverse	$A \cdot A^{-1} = I$							
RREF	<p>1. Row Swapping (Interchange)</p> $R_1 \leftrightarrow R_2$ <p>2. Row Scaling (Multiplication)</p> $R_1 \rightarrow \frac{1}{3}R_1$ <p>3. Row Addition (Replacement)</p> $R_1 \rightarrow R_1 - 2R_2$							

1. Matrix

$$m \begin{matrix} & & n \\ \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \end{matrix})$$

1.1. Matrix Vector Product

$$m \begin{matrix} & & n \\ \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \end{matrix} \begin{matrix} n \\ \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \end{matrix}$$

$$\begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix}$$

1.2. Matrix Multiplication

$$A = \begin{matrix} & & n \\ m & \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \end{matrix} \qquad B = \begin{matrix} & & p \\ n & \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1p} \\ b_{21} & b_{22} & \cdots & b_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{np} \end{bmatrix} \end{matrix}$$

$$A = \begin{matrix} & & n \\ m & \begin{bmatrix} \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \end{bmatrix} \\ \begin{bmatrix} a_{21} & a_{22} & \cdots & a_{2n} \end{bmatrix} \\ \vdots \\ \begin{bmatrix} a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} \end{bmatrix} \end{matrix} \qquad B = \begin{matrix} & & p \\ n & \left[\begin{bmatrix} b_{11} \\ b_{21} \\ \vdots \\ b_{n1} \end{bmatrix} \begin{bmatrix} b_{12} \\ b_{22} \\ \vdots \\ b_{n2} \end{bmatrix} \cdots \begin{bmatrix} b_{1p} \\ b_{2p} \\ \vdots \\ b_{np} \end{bmatrix} \right] \end{matrix}$$

A: Row Representation

$$A = \begin{bmatrix} r_1 \\ r_2 \\ \vdots \\ r_m \end{bmatrix}$$

$$r_i = [a_{i1} \ a_{i2} \ \cdots \ a_{in}], \quad \text{for } i = 1, 2, \dots, m$$

B: Column Representation

$$B = [c_1 \ c_2 \ \cdots \ c_p]$$
$$c_j = \begin{bmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{nj} \end{bmatrix}, \quad \text{for } j = 1, 2, \dots, p$$

$$C = \begin{matrix} & & p \\ m & \begin{bmatrix} r_1 \cdot c_1 & r_1 \cdot c_2 & \cdots & r_1 \cdot c_p \\ r_2 \cdot c_1 & r_2 \cdot c_2 & \cdots & r_2 \cdot c_p \\ \vdots & \vdots & \ddots & \vdots \\ r_m \cdot c_1 & r_m \cdot c_2 & \cdots & r_m \cdot c_p \end{bmatrix} \end{matrix}$$

1.3. Transpose

A^T

n

m

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

n

m

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{mn} \end{bmatrix}$$

n

m

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1j} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2j} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mj} & \cdots & a_{mn} \end{bmatrix}$$

n

m

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \cdots & a_{i1} & \cdots & a_{m1} \\ a_{12} & a_{22} & \cdots & a_{i2} & \cdots & a_{m2} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{1j} & a_{2j} & \cdots & a_{ij} & \cdots & a_{mj} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{1n} & a_{2n} & \cdots & a_{in} & \cdots & a_{mn} \end{bmatrix}$$

2. Matrix Factorization

2.1. LU Decomposition

$A = LU$

where:

- L : lower triangular matrix (entries above the diagonal are zero)
- U : upper triangular matrix (entries below the diagonal are zero)

1. $m \times n$ Matrix (with $m \geq n$)

n

m

$$L = \begin{bmatrix} l_{11} & l_{12} & \cdots & l_{1n} \\ l_{21} & l_{22} & \cdots & l_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ l_{m1} & l_{m2} & \cdots & l_{mn} \end{bmatrix}$$

n

m

$$U = \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ u_{21} & u_{22} & \cdots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ u_{n1} & u_{n2} & \cdots & u_{nn} \end{bmatrix}$$

2. $m \times n$ Matrix (with $m < n$)

see QR Decomposition

2.2. QR Decomposition

2.3. Cholesky Decomposition

2.4. Singular Value Decomposition (SVD)

$A = U \Sigma V^T$

m

m

$$U = \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1m} \\ u_{21} & u_{22} & \cdots & u_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ u_{m1} & u_{m2} & \cdots & u_{mm} \end{bmatrix}$$

n

m

$$\Sigma = \begin{bmatrix} \varepsilon_{11} & \varepsilon_{12} & \cdots & \varepsilon_{1m} \\ \varepsilon_{21} & \varepsilon_{22} & \cdots & \varepsilon_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ \varepsilon_{n1} & \varepsilon_{n2} & \cdots & \varepsilon_{nm} \end{bmatrix}$$

n

$$V^T = \begin{bmatrix} v_{11} & v_{12} & \cdots & v_{1n} \\ v_{21} & v_{22} & \cdots & v_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ v_{n1} & v_{n2} & \cdots & v_{nn} \end{bmatrix}$$

	$\left[\begin{array}{ccc c} 1 & 2 & -1 & 2 \\ 2 & 3 & 1 & 5 \\ 3 & 4 & -2 & 4 \end{array} \right]$
$R_2 \rightarrow R_2 - 2R_1$ $R_3 \rightarrow R_3 - 3R_1$	$\left[\begin{array}{ccc c} 1 & 2 & -1 & 2 \\ 0 & -1 & 3 & 1 \\ 0 & -2 & 1 & -2 \end{array} \right]$
$R_2 \rightarrow -R_2$	$\left[\begin{array}{ccc c} 1 & 2 & -1 & 2 \\ 0 & 1 & -3 & -1 \\ 0 & -2 & 1 & -2 \end{array} \right]$
$R_1 \rightarrow R_1 - 2R_2$ $R_3 \rightarrow R_3 + 2R_2$	$\left[\begin{array}{ccc c} 1 & 0 & 5 & 4 \\ 0 & 1 & -3 & -1 \\ 0 & 0 & -5 & -4 \end{array} \right]$
$R_3 \rightarrow \frac{1}{-5}R_3$	$\left[\begin{array}{ccc c} 1 & 0 & 5 & 4 \\ 0 & 1 & -3 & -1 \\ 0 & 0 & 1 & \frac{4}{5} \end{array} \right]$
$R_1 \rightarrow R_1 - 5R_3$ $R_2 \rightarrow R_2 + 3R_3$	$\left[\begin{array}{ccc c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & \frac{7}{5} \\ 0 & 0 & 1 & \frac{4}{5} \end{array} \right]$

$$A = \begin{bmatrix} 2 & -1 & -3 \\ -4 & 2 & 6 \end{bmatrix}$$

1. Null Space

The **null-space** of A , denoted $N(A)$, consists of all vectors $\vec{x} \in \mathbb{R}^3$ such that $A\vec{x} = \mathbf{0}$. The set of all such vectors is the **pre-image** of the zero vector under the transformation defined by A . In other words, $N(A) = \{x \in \mathbb{R}^3 \mid A\vec{x} = \mathbf{0}\}$, which represents the set of vectors that A maps to zero.

$$N(A) = \{\vec{x} \in \mathbb{R}^3 \mid A\vec{x} = \mathbf{0}\}$$

To find the null space $N(A)$ of the matrix A , we can use the **row-reduced echelon form (RREF)**. By augmenting the matrix A with a zero column and performing row operations, we reduce it to the form:

$$\begin{bmatrix} 2 & -1 & -3 \\ -4 & 2 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

	$\left[\begin{array}{ccc c} 2 & -1 & -3 & 0 \\ -4 & 2 & 6 & 0 \end{array} \right]$
--	---

$R_1 \rightarrow \frac{R_1}{2}$ $R_2 \rightarrow \frac{R_2}{4}$	$\left[\begin{array}{ccc c} 1 & -\frac{1}{2} & -\frac{3}{2} & 0 \\ -1 & \frac{1}{2} & \frac{3}{2} & 0 \end{array} \right]$
$R_2 \rightarrow R_2 - R_1$	$\left[\begin{array}{ccc c} 1 & -\frac{1}{2} & -\frac{3}{2} & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$

$$\begin{bmatrix} 1 & -\frac{1}{2} & -\frac{3}{2} \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$x_1 - \frac{1}{2}x_2 - \frac{3}{2}x_2 = 0$$

$$x_1 = \frac{1}{2}x_2 + \frac{3}{2}x_2$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = x_2 \begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} \frac{3}{2} \\ 0 \\ 1 \end{bmatrix}$$

$$N(a) = \text{span} \left(\left\{ \begin{bmatrix} \frac{1}{2} \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} \frac{3}{2} \\ 0 \\ 1 \end{bmatrix} \right\} \right)$$

The dimension of the **null-space** is the number of vectors in this basis, which is 2. This is important because the dimension of the null space gives us insight into how many degrees of freedom exist in the system of equations $Ax = 0$

2. Column Space

$$\begin{aligned} C(A) &= \text{span} \left(\left\{ \begin{bmatrix} 2 \\ -4 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix}, \begin{bmatrix} -3 \\ 6 \end{bmatrix} \right\} \right) \\ &= \text{span} \left(\left\{ \begin{bmatrix} 2 \\ -4 \end{bmatrix} \right\} \right) \end{aligned}$$

3. Basis

$$\begin{bmatrix} 2 \\ -4 \end{bmatrix}$$

4. Rank

Number of vector in the basis of our column space


$$\text{Rank}(A) = 1$$

3. Vectors

$$\text{Vector} = \text{Magnitude} + \text{Direction}$$

A car is moving:

$$\underbrace{\underbrace{3 \text{ MPH}}_{\substack{\text{Magnitude} \\ (\text{Speed} \rightarrow \text{Scalar})}} \quad \underbrace{\text{East}}_{\text{Direction}}}_{\text{Velocity (Vector)}}$$



$$\vec{v} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

3.1. Real Coordinate Spaces

N-dimensional Real Coordinate Space

$$\mathbb{R}^n$$

$$\vec{x} \in \mathbb{R}^n$$

All possible real-valued n-tuples

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$



3.2. Vector Operations

3.2.1. Vector Addition

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{bmatrix}$$

$$\vec{a} = \begin{bmatrix} 6 \\ -2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} -4 \\ 4 \end{bmatrix}$$

$$\vec{a} + \vec{b} = \begin{bmatrix} 6 + (-4) \\ -2 + 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$




3.2.2. Vector Subtraction

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 - y_1 \\ x_2 - y_2 \\ x_3 - y_3 \end{bmatrix}$$

$$\vec{u} = \begin{bmatrix} 1 \\ 5 \end{bmatrix} \quad \vec{v} = \begin{bmatrix} 5 \\ 1 \end{bmatrix}$$

$$\vec{u} - \vec{v} = \begin{bmatrix} 1 - 5 \\ 5 - 1 \end{bmatrix} = \begin{bmatrix} -4 \\ 4 \end{bmatrix}$$




3.2.3. Scalar Multiplication

$$c \times \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} c \times x_1 \\ c \times x_2 \\ c \times x_3 \end{bmatrix}$$


$$\vec{a} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$3\vec{a} = 3 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \cdot 2 \\ 3 \cdot 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$$



$$\vec{a} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$-1\vec{a} = -1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \cdot 2 \\ -1 \cdot 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -1 \end{bmatrix}$$



3.3. Unit Vector

A vector that has a magnitude (or length) of exactly 1

For a vector \vec{v} in n -dimensional space, a unit vector \hat{v} is defined as:

$$\hat{v} = \frac{\vec{v}}{\|\vec{v}\|}$$

Where:

- $\|\vec{v}\|$ is the **magnitude** (or **norm**) of the vector \vec{v} , computed as:

$$\|\vec{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

Key Properties

- **Magnitude:**

$$\|\hat{v}\| = 1$$

- **Direction:** A unit vector points in the same direction as the original vector \vec{v}

Finding unit vector (vector of magnitude 1) with given direction


$$\vec{a} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

Magnitude

$$\|\vec{a}\| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$

$$\hat{u} = \left(\frac{3}{\|\vec{a}\|}, \frac{4}{\|\vec{a}\|} \right) = \left(\frac{3}{5}, \frac{4}{5} \right)$$

$$\|\hat{u}\| = \sqrt{\left(\frac{3}{5}\right)^2 + \left(\frac{4}{5}\right)^2} = \sqrt{\frac{9}{25} + \frac{16}{25}} = \sqrt{\frac{25}{25}} = \sqrt{1} = 1$$



3.4. Parametric Representation of line

Set L of all points (i.e., line) equal to the set of all vectors \vec{x} plus some scalar t times the vector \vec{v} such that t can be any real number (\mathbb{R})

$$L = \{\vec{x} + t\vec{v} \mid t \in \mathbb{R}\}$$


$$\vec{a} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

$$t = 1$$


The line L can be defined as:

$$\begin{aligned} \vec{b} + t(\vec{b} - \vec{a}) &= \begin{bmatrix} 0 \\ 3 \end{bmatrix} + 1 \left(\begin{bmatrix} 0 \\ 3 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \\ &= \begin{bmatrix} 0 \\ 3 \end{bmatrix} + 1 \begin{bmatrix} -2 \\ 2 \end{bmatrix} \\ &= \begin{bmatrix} -2 \\ 5 \end{bmatrix} \end{aligned}$$



The line L can also be defined as:

$$\begin{aligned}\vec{a} + t(\vec{b} - \vec{a}) &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} + 1 \left(\begin{bmatrix} 0 \\ 3 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \\ &= \begin{bmatrix} 2 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} -2 \\ 2 \end{bmatrix} \\ &= \begin{bmatrix} 0 \\ 3 \end{bmatrix}\end{aligned}$$



Generalization

$$P_1 = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad P_2 = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$L = P_1 + t(P_1 - P_2) \mid t \in \mathbb{R}$$

$$\vec{P}_1 = \begin{bmatrix} -1 \\ 2 \end{bmatrix} \quad \vec{P}_2 = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

$$\vec{P}_1 - \vec{P}_2 = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

- L starts at P_1 and moves toward P_2 as t decreases

$$L = P_1 + t(P_1 - P_2) = \begin{bmatrix} -1 \\ 2 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

$$x = -1 + t(-1)$$

$$y = 2 + t(-1)$$

- L starts at P_2 and moves toward P_1 as t increases

$$L = P_2 + t(P_1 - P_2) = \begin{bmatrix} 0 \\ 3 \end{bmatrix} + t \begin{bmatrix} -1 \\ -1 \end{bmatrix}$$

$$x = 0 + t(-1)$$

$$y = 3 + t(-1)$$

- L starts at P_1 and moves toward P_2 as t increases

$$L = P_1 + t(P_2 - P_1) = \begin{bmatrix} -1 \\ 2 \end{bmatrix} + t \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$x = -1 + t(1)$$


$$y = 2 + t(1)$$

- L starts at P_2 and moves toward P_1 as t decreases

$$L = P_2 + t(P_1 - P_2) = \begin{bmatrix} 0 \\ 3 \end{bmatrix} + t \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$x = 0 + t(1)$$

$$y = 3 + t(-1)$$



3.5. Vector Spaces

Let's say your factory can produce up to 300 units of product 1, 500 units of product 2, and 400 units of product 3. The set of all possible production combinations forms a vector space:

$$x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$0 \leq x_1 \leq 300$$

$$0 \leq x_2 \leq 500$$

$$0 \leq x_3 \leq 400$$

4. Matrices

$m \times n$ matrix A

- m : rows
- n : columns

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

4.1. Matrix-Vector Products

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$A\vec{x} = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

For the dot product to be defined, the number of columns in the matrix A (which is n) must match the number of elements in the vector \vec{x} (also n).

The result of multiplying matrix A and vector \vec{x} will be a column vector with dimensions $m \times 1$, where m is the number of rows in the matrix A

$$(m \times n) \cdot (n \times 1) = m \times 1$$

1. As Row vectors

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$\vec{a}^T = [a_1, a_2, \dots, a_n]$$

$$\vec{b}^T = [b_1, b_2, \dots, b_n]$$

$$\mathbf{A} = \begin{bmatrix} [a_1, a_2, \dots, a_n] \\ [b_1, b_2, \dots, b_n] \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \vec{a} \\ \vec{b} \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$\begin{bmatrix} \vec{a}^T \\ \vec{b}^T \end{bmatrix} \cdot \vec{x} = \begin{bmatrix} \vec{a} \cdot \vec{x} \\ \vec{b} \cdot \vec{x} \end{bmatrix}$$

2. As Column Vectors

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} & \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} \end{bmatrix}$$

$$\mathbf{A} = [\vec{a} \ \vec{b}]$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\mathbf{A}\vec{x} = x_1\vec{a} + x_2\vec{b}$$

4.2. Null Space

The null space (or kernel) of a matrix \mathbf{A} is the set of all vectors \vec{x} that satisfy the equation:

$$A\vec{x} = \mathbf{0}$$

Where:

- A : $m \times n$ matrix
- \vec{x} : n -dimensional vector
- $\mathbf{0}$: zero vector in \mathbb{R}^m

$$N(A) = N(\text{rref}(A)) = \text{span}(\vec{v}_1, \vec{v}_2, \vec{v}_3)$$

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

We want to find the null space of A , which consists of all vectors $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$ that satisfy:

$$A\vec{x} = \mathbf{0}$$

This expands to the following system of linear equations:

$$\begin{cases} 1x_1 + 1x_2 + 1x_3 + 1x_4 = 0 \\ 1x_1 + 2x_2 + 3x_3 + 4x_4 = 0 \\ 4x_1 + 3x_2 + 2x_3 + 1x_4 = 0 \end{cases}$$

This can be represented as the augmented matrix:

$$\left[\begin{array}{cccc|c} 1 & 1 & 1 & 1 & 0 \\ 1 & 2 & 3 & 4 & 0 \\ 4 & 3 & 2 & 1 & 0 \end{array} \right]$$

4.2.1. Column Space

The **column space** (or range) of matrix A is span of its columns vectors

If the matrix A has columns $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$, then the column space of A is defined as:

$$\text{Col}(A) = \{\vec{y} \in \mathbb{R}^m \mid \vec{y} = A\vec{x} \text{ for some } \vec{x} \in \mathbb{R}^n\}$$

or equivalently,

$$\text{Col}(A) = \text{span}(\{\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n\})$$

Consider the simple example of a 2×2 matrix:

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$$

The matrix has two columns:

$$\vec{a}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \quad \text{and} \quad \vec{a}_2 = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$$

The column space, denoted $\text{Col}(A)$, is the span of these two vectors:

$$\text{Col}(A) = \text{span}\left(\left\{\begin{bmatrix} 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 6 \end{bmatrix}\right\}\right)$$

Finding the Column Space

We observe that the two columns \vec{a}_1 and \vec{a}_2 are **linearly dependent**:

$$\vec{a}_2 = k\vec{a}_1$$

This means that \vec{a}_2 is a scalar multiple of \vec{a}_1 , the the two columns are **linearly dependent**. As a result, the column space is spanned by just one vector, \vec{a}_1 , because any linear combination of \vec{a}_1 and \vec{a}_2 can be reduced to a multiple of \vec{a}_1 .

Therefore, the column space of A is:

$$\text{Col}(A) = \text{span}\left(\left\{\begin{bmatrix} 1 \\ 3 \end{bmatrix}\right\}\right)$$

which represents all vectors of the form:

$$c \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} c \\ 3c \end{bmatrix} \quad \text{for any scalar } c$$

In other words, the column space is a line in \mathbb{R}^2 through the origin in the direction of

Rank of A

The rank of A , which is the **dimension of its column space**, is 1 because there is only one linearly independent column

This means the column space is the span of the columns of A , or all vectors that can be formed by taking linear combinations of the columns of A .

4.2.2. Dimension of a Subspace

Number of elements in a basis for the subspace

4.2.3. Nullity

Dimension of the Null Space

$$\dim(N(A))$$

The nullity of A : number of non-pivot columns (i.e., free variables) in the rref of A

4.2.4. Rank

Dimension of the column space

$$\text{rank}(A) = \dim(C(A))$$

4.2.5. Matrix Representation of Systems of Equations

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1m}x_m &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2m}x_m &= b_2 \\ &\vdots \\ a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nm}x_m &= b_n \end{aligned}$$

Coefficient Matrix (A):

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{bmatrix}$$

Variable Vector (x):

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix}$$

Constant Vector (b):

$$\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$A\mathbf{x} = \mathbf{b}$$

The system of equations:

$$2x_1 + 3x_2 + 5x_3 = 100$$

$$4x_1 + 2x_2 + 1x_3 = 80$$

$$1x_1 + 5x_2 + 2x_3 = 60$$

Can be represented as a matrix equation:

$$\begin{bmatrix} 2 & 3 & 5 \\ 4 & 2 & 1 \\ 1 & 5 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 100 \\ 80 \\ 60 \end{bmatrix}$$

4.3. Matrix Multiplication

$m \times n$ matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{bmatrix}$$

$n \times p$ matrix:

$$B = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1p} \\ a_{21} & a_{22} & \dots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{np} \end{bmatrix}$$

Compute Each Element of Result Matrix C

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

Let A be an $n \times m$ matrix:

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

Let B be an $p \times n$ matrix:

$$B = \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$$

Calculate Each Element of C

$$c_{11} = (1 \cdot 7) + (2 \cdot 9) + (3 \cdot 11) = 58$$

$$c_{12} = (1 \cdot 8) + (2 \cdot 10) + (3 \cdot 12) = 64$$

$$c_{21} = (4 \cdot 7) + (5 \cdot 9) + (6 \cdot 11) = 138$$

$$c_{22} = (4 \cdot 8) + (5 \cdot 10) + (6 \cdot 12) = 154$$

C is a $m \times p$ matrix

$$C = \begin{bmatrix} 58 & 64 \\ 138 & 154 \end{bmatrix}$$

5. Linear Combinations

Set of vector

$$v_1, v_2, \dots, v_n \in \mathbb{R}^n$$

Where

- v_1, v_2, \dots, v_n : set of vectors
- \mathbb{R}^n : set of all ordered tuples of n real numbers

Linear combination of those vector

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n$$

$$c_1, c_2, \dots, c_n \in \mathbb{R}$$

Where:

- c_1, c_2, \dots, c_n : constants or weights

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

$$0\vec{a} + 0\vec{b}$$

$$3\vec{a} + 2\vec{b}$$

6. Span

Represents the subspace of the vector space that is “covered” by these vectors through their linear combinations

If you have a set of vectors v_1, v_2, \dots, v_n , the span of these vectors is the set of all vectors that can be written as:

$$\text{Span}(v_1, v_2, \dots, v_n) = \{c_1 v_1 + c_2 v_2 + \dots + c_n v_n \mid c_1, c_2, \dots, c_n \in \mathbb{R}\}$$


Any vector in \mathbb{R}^2 can be represented by a linear combination with some combination of these vectors

1. Spanning \mathbb{R}^2

$$\text{Span}(\vec{a}, \vec{b}) = \mathbb{R}^2$$

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

$$3\vec{a} + (-2)\vec{b} = \begin{bmatrix} 3 - 0 \\ 6 - 6 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$



Any point \vec{x} can be represented as a linear combination of \vec{a} and \vec{b}

1. Define the vectors

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 0 \\ 3 \end{bmatrix} \quad \vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

2. Express \vec{x} as a linear combinations

$$c_1 \vec{a} + c_2 \vec{b} = \vec{x}$$

Which expands to

$$c_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + c_2 \begin{bmatrix} 0 \\ 3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

3. Set up the system of equations

$$1c_1 + 0c_2 = x_1 \quad (1)$$

$$2c_1 + 3c_2 = x_2 \quad (2)$$

4. Express c_1 : From equation (1), we can directly express c_1

$$c_1 = x_1$$

5. Substitute c_1 into equation (2)

$$2x_1 + 3c_2 = x_2$$

Rearranging gives:

$$3c_2 = x_2 - 2x_1$$

6. Solve for c_2 : Dividing both sides by 3 yields

$$c_2 = \frac{x_2 - 2x_1}{3}$$

7. Example with specific values: Let's say we want to find c_1 and c_2 when $\vec{x} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$

Substitute $x_1 = 2$ and $x_2 = 2$

$$c_1 = x_1 = 2$$

$$c_2 = \frac{2 - 2 \cdot 2}{3} = -\frac{2}{3}$$

8. Final linear combination: Now, substituting c_1 and c_2 back into the linear combination

$$2\vec{a} - \frac{2}{3}\vec{b} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

Verifying

$$2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \frac{1}{3} \begin{bmatrix} 0 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$

9. This shows that


$$2\vec{a} - \frac{2}{3}\vec{b} = \vec{x}$$

2. Spanning Line in \mathbb{R}^2

Any linear combination of \vec{a} and \vec{b} will produce vectors that lie along the same line. This is the line through the origin in the direction of \vec{a} (or \vec{b}), with all points on the line being scalar multiples of \vec{a}

$$\vec{a} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} -2 \\ -2 \end{bmatrix}$$

$$3\vec{a} + (-2)\vec{b} = \begin{bmatrix} 6 - 4 \\ 6 - 4 \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$$



7. Linear Independence

1. Definition of Linear Independence

The set of vectors

$$S = \{v_1, v_2, \dots, v_n\}$$

is said to be **linearly independent** if the only solution to the equation

$$c_1 v_1 + c_2 v_2 + \dots + c_n v_n = \mathbf{0}$$

is $c_1 = c_2 = \dots = c_n = 0$. In other words, no vector in the set can be written as a linear combination of the others.

If at least one constant c_i is non-zero, the set is linearly dependent.

Example 1: Testing for Linear Independence

Problem: Is the following set of vectors **linearly dependent**?

$$S = \{\vec{v}_1, \vec{v}_2\}$$

Where:

$$\vec{v}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{v}_2 = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

For a set of vectors to be **linearly independent**, the only solution to the equation:

$$c_1 \vec{v}_1 + c_2 \vec{v}_2 = \mathbf{0}$$

must be $c_1 = 0$ and $c_2 = 0$

In this case:

$$c_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \mathbf{0}$$

If not only the zero solution exists (i.e., if c_1 or c_2 can be non-zero), the set is **linearly dependent**.

Step 1. Set up the system of equations:

1. $2c_1 + 3c_2 = 0$
2. $1c_1 + 2c_2 = 0$

2. Eliminate one variable

$$2 \times (1c_1 + 2c_2 = 0) \Rightarrow 2c_1 + 4c_2 = 0$$

Now the system is

$$2c_1 + 3c_2 = 0$$

$$2c_1 + 4c_2 = 0$$

3. Subtract the equations

$$(2c_1 + 3c_2) - (2c_1 + 4c_2) = 0$$

Simplifies to:

$$(2c_1 - 2c_1) + (4c_2 - 3c_2) = 0$$

So:

$$c_2 = 0$$

4. Substitute back to find c_1

Now that we know $c_2 = 0$, substitute this value into one of the original equations. Let's use the second equation:

$$1c_1 + 2c_2 = 0$$

Substitute $c_2 = 0$:

$$1c_1 + 2(0) = 0$$

$$c_1 = 0$$

Conclusion:

Since $c_1 = 0$ and $c_2 = 0$, the set of vectors S is **linearly independent**. These vectors span \mathbb{R}^2 .

Example 2: Testing for Linear Dependence

Problem: Is the following set of vectors **linearly dependent**?

$$S = \{\vec{v}_1, \vec{v}_2\}$$

Where:

$$\vec{v}_1 = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \quad \text{and} \quad \vec{v}_2 = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

The span of this set is the collection of all vectors that can be formed by linear combinations of \vec{v}_1 and \vec{v}_2 :

$$c_1 v_1 + c_2 v_2$$

Since $v_2 = 2v_1$, the linear combination becomes:

$$\begin{aligned} c_1 v_1 + c_2 (2v_1) &= (c_1 + 2c_2)v_1 \\ c_1 \begin{bmatrix} 2 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} 4 \\ 6 \end{bmatrix} \\ c_1 \begin{bmatrix} 2 \\ 3 \end{bmatrix} + c_2 2 \begin{bmatrix} 2 \\ 3 \end{bmatrix} \\ (c_1 + 2c_2) \begin{bmatrix} 2 \\ 3 \end{bmatrix} \\ c_3 \begin{bmatrix} 2 \\ 3 \end{bmatrix} \end{aligned}$$

Thus, any linear combination of these vectors is just a scalar multiple of v_1 . The span is a single line in \mathbb{R}^2 , and the vectors are **linearly dependent**.

For any two **colinear** vectors in \mathbb{R}^2 , their span reduces to a single line.

One vector in the set can be represented by some combination of other vectors in the set

2. General Rule

In \mathbb{R}^n , if you have more than n vectors, at least one vector must be linearly dependent on the others, meaning the set cannot be linearly independent.

8. Subspace

V is a linear subspace of \mathbb{R}^n :

- **Non-emptiness:** V contains the **0** vector

$$\mathbf{0} \in V$$

- **Closure under addition:** If u and v are any vectors in the subspace V , then their sum $u + v$ must also be in V .

$$\text{If } u, v \in V, \text{ then } u + v \in V$$

- **Closure under scalar multiplication:** If u is any vector in V and c is any scalar (real number), then the product cu must also be in V .

$$\text{If } u \in V \text{ and } c \in \mathbb{R}, \text{ then } cu \in V$$

Example 1: Subspace

Problem: Is V a subspace of \mathbb{R}^2

$$V = \{\mathbf{0}\} = \left\{ \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\}$$

- **Non-emptiness** ✓

$$\mathbf{0} \in V$$

- **Closure under addition** ✓

$$\begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- **Closure under scalar multiplication** ✓

$$c \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$


Conclusion

The subset V of \mathbb{R}^3 is a **subspace**

Example 2: Not Subspace

Problem: Is S a subspace of \mathbb{R}^2 ✓

$$S = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2 \mid x_1 \geq 0 \right\}$$



- **Non-emptiness** ✓

$$\mathbf{0} \in S$$

- **Closure under addition** ✓

$$\begin{bmatrix} a \\ b \end{bmatrix} + \begin{bmatrix} c \\ d \end{bmatrix} = \begin{bmatrix} a+c \\ b+d \end{bmatrix}$$

$$a \geq 0$$

$$b \geq 0$$

$$a+b \geq 0$$

- **Closure under scalar multiplication** ✗

$$-1 \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix}$$

Conclusion

Span and Subspace

The span of any set of vectors is a valid subspace

$$U = \text{Span}(v_1, v_2, \dots, v_n) = \text{Valid Subspace of } \mathbb{R}^n$$

- **Non-emptiness**

$$0v_1 + 0v_2 + \dots + 0v_n = \mathbf{0}$$

- **Closure under addition**

$$\vec{X} = a_1 v_1 + a_2 v_2 + \dots + a_n v_n$$

$$\vec{Y} = b_1 v_1 + b_2 v_2 + \dots + b_n v_n$$

$$\begin{aligned}\vec{X} + \vec{Y} &= (a_1 + b_1)v_1 + (a_2 + b_2)v_2 + \dots + (a_n + b_n)v_n \\ &= c_1 v_1 + c_2 v_2 + \dots + c_n v_n\end{aligned}$$

- **Closure under scalar multiplication**

$$\vec{X} = a_1 v_1 + a_2 v_2 + \dots + a_n v_n$$

$$\begin{aligned}b\vec{X} &= bc_1 v_1 + bc_2 v_2 + \dots + bc_n v_n \\ &= c_1 v_1 + c_2 v_2 + \dots + c_n v_n\end{aligned}$$

9. Basis

Non-redundant set of vectors that span \mathbb{R}^n

A basis of a vector space is a set of vectors that satisfies two conditions:

1. **Linear Independence:** No vector in the set can be written as a linear combination of the others. This means that **the only way to combine the vectors to get the zero vector is by using all zero coefficients.**

The vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly independent if the only solution to the equation

$$c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_n \vec{v}_n = \mathbf{0}$$

is $c_1 = c_2 = \dots = c_n = 0$, where c_i are scalar coefficients.

2. **Spanning:** The set of vectors can be linearly combined to form any vector in the vector space. In other words, **every vector in the vector space can be expressed as a linear combination of the basis vectors.**

The set $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ spans the vector space V if any vector $v \in V$ can be expressed as a linear combination of the basis vectors:

$$\vec{v} = c_1 \vec{v}_1 + c_2 \vec{v}_2 + \dots + c_n \vec{v}_n$$

for some scalars c_1, c_2, \dots, c_n .

Consider the vector space \mathbb{R}^2 (the 2-dimensional Euclidean space). A common basis for \mathbb{R}^2 is $\{e_1, e_2\}$, where:

$$e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

This set is a basis because:

- Linear Independence: The only solution to $c_1 e_1 + c_2 e_2 = \mathbf{0}$ is $c_1 = c_2 = 0$.
- Spanning: Any vector $\vec{v} = \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$ can be written as $\vec{v} = x e_1 + y e_2$


This means $\{e_1, e_2\}$ is a basis for \mathbb{R}^2 , and the dimension of \mathbb{R}^2 is 2.

9.1. Vector Dot Product

$$\vec{a} \cdot \vec{b} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} = a_1 b_1 + a_2 b_2 + \dots + a_n b_n$$

9.1.1. Magnitude (Length)

$$\|\vec{a}\| = \sqrt{a_1^2 + a_2^2 + \dots + a_n^2}$$



$$\vec{a} \cdot \vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = a_1^2 + a_2^2 + \dots + a_n^2$$

$$\|\vec{a}\| = \sqrt{\vec{a} \cdot \vec{a}}$$

$$\|\vec{a}\|^2 = \vec{a} \cdot \vec{a}$$

9.1.2. Properties

9.1.2.1. Commutative

$$\vec{v} \cdot \vec{w} = \vec{w} \cdot \vec{v}$$

9.1.2.2. Distributive

$$(\vec{v} + \vec{w}) \cdot \vec{x} = \vec{v} \cdot \vec{x} + \vec{w} \cdot \vec{x}$$

9.1.2.3. Associativity

$$(c\vec{v}) \cdot \vec{w} = c(\vec{v} \cdot \vec{w})$$

9.1.3. Cauchy-Schwarz Inequality

$$|\vec{u} \cdot \vec{v}| \leq \|\vec{u}\| \|\vec{v}\|$$

$$|\vec{u} \cdot \vec{v}| = \|\vec{u}\| \|\vec{v}\| \quad \text{when} \quad \vec{u} = c\vec{v}$$

Where:

- $\vec{u} \cdot \vec{v}$: dot product of vectors \vec{u} and \vec{v}
- $\|\vec{u}\|$ and $\|\vec{v}\|$: magnitudes (lengths) of vectors \vec{u} and \vec{v}

Step 1: Understand the dot product

The dot product of two vectors $\vec{u} = [u_1, u_2, \dots, u_n]$ and $\vec{v} = [v_1, v_2, \dots, v_n]$ is calculated as:

$$\vec{u} \cdot \vec{v} = u_1v_1 + u_2v_2 + \dots + u_nv_n$$

The magnitude (or norm) of a vector \vec{u} is:

$$\|\vec{u}\| = \sqrt{u_1^2 + u_2^2 + \dots + u_n^2}$$

Step 2: Define a new function

We introduce a parameter $t \in \mathbb{R}$ and define a new vector:

$$w(t) = \vec{u} - t\vec{v}$$

Now, consider the dot product of this new vector with itself, which is always non-negative because it represents the square of the magnitude of $w(t)$:

$$w(t) \cdot w(t) \geq 0$$

This inequality makes sense because the dot product of any vector with itself is the square of its magnitude, and **a square is always non-negative**.

Step 3: Expand the dot product

Expand $w(t) \cdot w(t)$:

$$w(t) \cdot w(t) = (\vec{u} - t\vec{v}) \cdot (\vec{u} - t\vec{v})$$

Now, we apply the distributive property of the dot product, which behaves similarly to the distributive property of multiplication. We expand each term:

$$(\vec{u} - t\vec{v}) \cdot (\vec{u} - t\vec{v}) = \vec{u} \cdot \vec{u} - t(\vec{u} \cdot \vec{v}) - t(\vec{v} \cdot \vec{u}) + t^2(\vec{v} \cdot \vec{v})$$

Since the dot product is commutative ($\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$), we can rewrite this as:

$$\vec{u} \cdot \vec{u} - 2t(\vec{u} \cdot \vec{v}) + t^2(\vec{v} \cdot \vec{v})$$

This simplifies to:

$$\|\vec{u}\|^2 - 2t(\vec{u} \cdot \vec{v}) + t^2 \|\vec{v}\|^2$$

We've now expressed the result of expanding the dot product as a quadratic expression in t , where

- $\|\vec{u}\|^2$ is a constant term,
- $-2t(\vec{u} \cdot \vec{v})$ is the linear term in t
- $t^2 \|\vec{v}\|^2$ is the quadratic term

Step 4: Treat as a quadratic equation

Now that we have the quadratic expression:

$$\|\vec{v}\|^2 t^2 - 2(\vec{u} \cdot \vec{v})t + \|\vec{u}\|^2 \geq 0$$

We recognize this as a standard quadratic inequality of the form $at^2 + bt + c \geq 0$, where:

- $a = \|\vec{v}\|^2$
- $b = -2(\vec{u} \cdot \vec{v})$
- $c = \|\vec{u}\|^2$

For any quadratic expression $at^2 + bt + c$ to always be non-negative, its discriminant must be less than or equal to zero. The discriminant of a quadratic equation $at^2 + bt + c = 0$ is given by:

$$\Delta = b^2 - 4ac$$

Substituting in the values of a , b , and c from our expression:

$$\Delta = (-2(\vec{u} \cdot \vec{v}))^2 - 4 \cdot \|\vec{v}\|^2 \cdot \|\vec{u}\|^2$$

Simplifying:

$$\Delta = 4(\vec{u} \cdot \vec{v})^2 - 4 \|\vec{v}\|^2 \|\vec{u}\|^2$$

Step 5: Apply the discriminant condition

For the quadratic inequality to hold, the discriminant must be less than or equal to zero:

$$\Delta = 4(\vec{u} \cdot \vec{v})^2 - 4 \|\vec{v}\|^2 \|\vec{u}\|^2 \leq 0$$

Divide by 4:

$$\Delta = (\vec{u} \cdot \vec{v})^2 \leq \|\vec{v}\|^2 \|\vec{u}\|^2$$

Take the square root of both sides:

$$|\vec{u} \cdot \vec{v}| \leq \|\vec{u}\| \|\vec{v}\|$$

$$\vec{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \vec{v} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

Step 1: Compute the dot product $\vec{u} \cdot \vec{v}$

$$\vec{u} \cdot \vec{v} = (1)(3) + (2)(4) = 3 + 8 = 11$$

Step 2: Compute the norms of \vec{u} and \vec{v}

- The norm $\|\vec{u}\|$ is:

$$\|\vec{u}\| = \sqrt{1^2 + 2^2} = \sqrt{1 + 4} = \sqrt{5}$$

- The norm $\|\vec{v}\|$ is:

$$\|\vec{v}\| = \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} = 5$$

Step 3: Verify the Cauchy-Schwarz inequality

The inequality states:

$$|\vec{u} \cdot \vec{v}| \leq \|\vec{u}\| \|\vec{v}\|$$

Substitute the values:


$$|11| \leq 5\sqrt{5} = 11.18$$

Since $11 \leq 11.18$, the inequality holds.


The Cauchy-Schwarz inequality is satisfied for the vectors $\vec{u} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\vec{v} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$

9.1.4. Vector Triangle Inequality

$$\|\vec{u} + \vec{v}\| \leq \|\vec{u}\| + \|\vec{v}\|$$



$$\|\vec{u} + \vec{v}\| = \|\vec{u}\| + \|\vec{v}\|$$





$$\vec{u} = c\vec{v} \quad c > 0$$

9.2. Angles Between Vectors

The scalar $\|\vec{u}\|$ is the length of the vector \vec{u}

Say $\vec{u}, \vec{v} \in \mathbb{R}^n$



Law of Cosines

$$c^2 = a^2 + b^2 - 2ab \cdot \cos(C)$$

Where:

- a, b and c : lengths of the sides of a triangle
- C : angle opposite the side c


$$\begin{aligned} \|\vec{u} - \vec{v}\|^2 &= \|\vec{u}\|^2 + \|\vec{v}\|^2 - 2 \|\vec{u}\| \|\vec{v}\| \cdot \cos(\Theta) \\ (\vec{u} - \vec{v}) \cdot (\vec{u} - \vec{v}) &= \\ \vec{u} \cdot \vec{u} - \vec{u} \cdot \vec{v} - \vec{v} \cdot \vec{u} + \vec{v} \cdot \vec{v} &= \\ \|\vec{u}\|^2 - 2(\vec{u} \cdot \vec{v}) + \|\vec{v}\|^2 &= \|\vec{u}\|^2 + \|\vec{v}\|^2 - 2 \|\vec{u}\| \|\vec{v}\| \cdot \cos(\Theta) \end{aligned}$$

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cdot \cos(\Theta)$$


$$\frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} = \cos(\Theta)$$

$$\Theta = \arccos\left(\frac{\vec{a} \cdot \vec{b}}{\|\vec{a}\| \|\vec{b}\|}\right)$$

So, if \vec{u} is a scalar multiple of \vec{v} ($\vec{u} = c\vec{v}$) where $c > 0$, then $\Theta = 0^\circ$



And, if \vec{u} is a scalar multiple of \vec{v} ($\vec{u} = c\vec{v}$) where $c < 0$, then $\Theta = 180^\circ$



\vec{u} and \vec{v} are perpendicular if the angle Θ between them is 90°


$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cdot \cos(90^\circ)$$

$$\vec{u} \cdot \vec{v} = 0$$

If \vec{u} and \vec{v} are perpendicular, then $\vec{u} \cdot \vec{v} = 0$

If \vec{u} and \vec{v} are non-zero and $\vec{u} \cdot \vec{v} = 0$, then they are perpendicular

If $\vec{u} \cdot \vec{v} = 0$ then \vec{u} and \vec{v} are **orthogonal**.



9.2.1. Plane in \mathbb{R}^3


Plane: Each point (x, y, z) on the satisfies the equation

$$ax + by + cz = d$$

Normal Vector: vector that is perpendicular (orthogonal) to a plane, line, or curve, at a specific point

1. Plane

If a plane is defined by the equation $ax + by + cz = d$, the vector $\vec{n} = \langle a, b, c \rangle$ is a normal vector to the plane because it is perpendicular to any vector that lies in the plane



$$\vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \quad \vec{x}_0 = \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix} \quad \vec{n} = \begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix}$$

$$\vec{x} - \vec{x}_0 = \begin{bmatrix} x - x_0 \\ y - y_0 \\ z - z_0 \end{bmatrix}$$

$\vec{x} - \vec{x}_0$ is a vector that lies on the plane, then \vec{n} is normal if:

$$\vec{n} \cdot \vec{x} - \vec{x}_0 = 0$$

$$\begin{bmatrix} n_1 \\ n_2 \\ n_3 \end{bmatrix} \cdot \begin{bmatrix} x - x_0 \\ y - y_0 \\ z - z_0 \end{bmatrix} = 0$$

$$n_1(x - x_0) + n_2(y - y_0) + n_3(z - z_0) = 0$$

Find the equation of the plane given the point on the plane \vec{x}_0 and a the normal vector \vec{n}

$$\vec{n} = \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \quad \vec{x}_0 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad \vec{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$


$$\vec{x} - \vec{x}_0 = \begin{bmatrix} x-1 \\ y-2 \\ z-3 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix} \cdot \begin{bmatrix} x-1 \\ y-2 \\ z-3 \end{bmatrix} = 0$$

$$(x-1) + 3(y-2) - 2(z-3) = 0$$

$$x-1 + 3y-6 - 2z+6 = 0$$

$$x + 3y - 2z = 1$$



The normal vector to a plane can be directly obtained from the coefficients of x , y , and z in the plane equation of the form:

$$Ax + By + Cz = D$$

$$\vec{n} = \begin{bmatrix} A \\ B \\ C \end{bmatrix}$$

Find the equation of the normal vector \vec{n} given the equation for the plane:

$$-3x + \sqrt{2}y + 7z = \pi$$

$$\vec{n} = -3\hat{i} + \sqrt{2}\hat{j} + 7\hat{k}$$

$$\vec{n} = \begin{bmatrix} -3 \\ \sqrt{2} \\ 7 \end{bmatrix}$$

2. Curve

For a curve described by a function $y = f(x)$, the normal vector at a point on the curve is perpendicular to the tangent line at that point. If the tangent vector has slope $f'(x)$, the normal vector will have a slope of $-\frac{1}{f'(x)}$

9.2.2. Point Distance to Plane

$$d = \frac{Ax_0 + By_0 + Cz_0 - D}{\sqrt{A^2 + B^2 + C^2}}$$

Given a the equation of the plane:

$$1x - 2y + 3z = 5$$

And a point **not** on the plane:

$$(2, 3, 1)$$

Find the shortest path (normal vector) from the plane to the point

$$\begin{aligned} d &= \frac{Ax_0 + By_0 + Cz_0 - D}{\sqrt{A^2 + B^2 + C^2}} \\ &= \frac{1 \cdot 2 - 2 \cdot 3 + 3 \cdot 1 - 5}{\sqrt{1^2 + 2^2 + 3^2}} \\ &= \frac{2 - 6 + 3 - 5}{\sqrt{1 + 4 + 9}} \\ &= -\frac{6}{\sqrt{14}} \end{aligned}$$

9.2.3. Distance Between Planes

9.2.4. Cross Product

Only defined in \mathbb{R}^3

Returns a vector orthogonal to the two vectors

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$\vec{c} = \vec{a} \times \vec{b}$$

$$\vec{c} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix}$$

$$\vec{a} = \begin{bmatrix} 1 \\ -7 \\ 1 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 5 \\ 2 \\ 4 \end{bmatrix}$$

$$\vec{c} = \vec{a} \times \vec{b}$$

$$\vec{c} = \begin{bmatrix} -7 \cdot 4 - 1 \cdot 2 \\ 1 \cdot 5 - 1 \cdot 4 \\ 1 \cdot 2 - (-7) \cdot 5 \end{bmatrix} = \begin{bmatrix} -30 \\ 1 \\ 37 \end{bmatrix}$$

\vec{c} is orthogonal to both \vec{a} and \vec{b}

Proof: \vec{c} is orthogonal to \vec{a} and \vec{b}

When the dot product of two vectors is equal to 0, it means that the two vectors are perpendicular (or orthogonal) to each other

1. Orthogonal to Vector \vec{a}

$$\vec{c} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix}$$

$$a_1a_2b_3 - a_1a_3b_2 + a_2a_3b_1 - a_2a_1b_3 + a_3a_1b_2 - a_3a_2b_1$$

$$\cancel{a_1a_2b_3} - \cancel{a_1a_3b_2} + \cancel{a_2a_3b_1} - \cancel{a_2a_1b_3} + \cancel{a_3a_1b_2} - \cancel{a_3a_2b_1} = 0$$

2. Orthogonal to Vector \vec{b}

$$\vec{c} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix} \cdot \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$b_1a_2b_3 - b_1a_3b_2 + b_2a_3b_1 - b_2a_1b_3 + b_3a_1b_2 - b_3a_2b_1$$

$$\cancel{b_1a_2b_3} - \cancel{b_1a_3b_2} + \cancel{b_2a_3b_1} - \cancel{b_2a_1b_3} + \cancel{b_3a_1b_2} - \cancel{b_3a_2b_1} = 0$$

9.2.5. Proof: Relationship Between Cross Product and Sin of Angle

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

$$\vec{c} = \vec{a} \times \vec{b}$$

$$\vec{c} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \times \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} a_2b_3 - a_3b_2 \\ a_3b_1 - a_1b_3 \\ a_1b_2 - a_2b_1 \end{bmatrix}$$

$$\vec{a} \cdot \vec{b} = \|\vec{a}\| \|\vec{b}\| \cos(\Theta)$$

$$\|\vec{a} \times \vec{b}\| = \|\vec{a}\| \|\vec{b}\| \sin(\Theta)$$

9.2.6. Dot and Cross Products

$$\vec{a} \cdot \vec{b} = \|\vec{a}\| \|\vec{b}\| \cos(\Theta)$$

$$\frac{\vec{a} \cdot \vec{b}}{\|\vec{a}\| \|\vec{b}\|} = \cos(\Theta)$$

$$\Theta = \arccos()$$

9.3. Row Echelon Form (REF)

Visual Structure:

1. Pivot (leading 1): The leading entry of each non-zero row is 1

2. Zeros below pivots: Every pivot has zeros below it in its column
3. Rightward movement of pivots: Each leading 1 in a lower row is further to the right than in the row above it
4. Rows of all zeros (if any) are at the bottom of the matrix

$$\left[\begin{array}{cccc|c} 1 & a_{12} & a_{13} & a_{14} & b_1 \\ 0 & 1 & a_{23} & a_{24} & b_2 \\ 0 & 0 & 1 & a_{34} & b_3 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

Elementary row operations:

1. **Row Swapping:** Swap two rows
2. **Row Multiplication:** Multiply a row by a non-zero scalar
3. **Row Addition / Subtraction:** Add or subtract a multiple of one row from another row

Consider the system of linear equations:

$$\begin{aligned} 2x + y + z &= 8 \\ -3x - y + 2z &= -11 \\ -2x + 1y + 2z &= -3 \end{aligned}$$

The augmented matrix for this system is:

$$\left[\begin{array}{ccc|c} 2 & 1 & 1 & 8 \\ -3 & -1 & 2 & -11 \\ -2 & 1 & 2 & -3 \end{array} \right]$$

Step 1: Make the leading entry of the first row a 1

We divide the first row by 2 (row multiplication):

$$R_1 \rightarrow \frac{1}{2}R_1 = \left[\begin{array}{ccc|c} 1 & 0.5 & 0.5 & 4 \\ -3 & -1 & 2 & -11 \\ -2 & 1 & 2 & -3 \end{array} \right]$$

Step 2: Eliminate the entries below the first pivot We now want the entries below the first pivot (1 in the first column) to become zeros. We use row addition:

1. $R_2 \rightarrow R_2 + 3R_1$
2. $R_3 \rightarrow R_3 + 2R_1$

This gives:

$$\left[\begin{array}{ccc|c} 1 & 0.5 & 0.5 & 4 \\ 0 & 0.5 & 3.5 & 1 \\ 0 & 2 & 3 & 5 \end{array} \right]$$

Step 3: Make the leading entry of the second row a 1

We divide the second row by 0.5 (row multiplication):

$$R_2 \rightarrow \frac{1}{0.5}R_2 = \left[\begin{array}{ccc|c} 1 & 0.5 & 0.5 & 4 \\ 0 & 1 & 7 & 2 \\ 0 & 2 & 3 & 5 \end{array} \right]$$

Step 4: Eliminate the entry below the second pivot

We now want the entry below the second pivot (1 in the second column) to become zero. We use row addition:

$$R_3 \rightarrow R_2 - 2R_2 = \left[\begin{array}{ccc|c} 1 & 0.5 & 0.5 & 4 \\ 0 & 1 & 7 & 2 \\ 0 & 0 & -11 & 1 \end{array} \right]$$

Step 5: Make the leading entry of the third row a 1

We divide the third row by -11 (row multiplication):

$$R_3 \rightarrow -\frac{1}{11}R_3 = \left[\begin{array}{ccc|c} 1 & 0.5 & 0.5 & 4 \\ 0 & 1 & 7 & 2 \\ 0 & 0 & 1 & -\frac{1}{11} \end{array} \right]$$

Step 6: Back-substitute to solve for the variables

From the third row:

$$z = -\frac{1}{11}$$

Substitute z into the second row:

$$\begin{aligned} y + 7z &= 2 \\ y + 7\left(-\frac{1}{11}\right) &= 2 \\ y &= 2 + \frac{7}{11} \\ y &= \frac{29}{11} \end{aligned}$$

Substitute y and z into the first row:

$$\begin{aligned} x + 0.5y + 0.5z &= 4 \\ x + 0.5\left(\frac{29}{11}\right) + 0.5\left(-\frac{1}{11}\right) &= 4 \\ x &= 4 - \frac{14}{11} + \frac{1}{11} \\ x &= \frac{31}{11} \end{aligned}$$

Final Solution:

$$x = \frac{31}{11} \quad y = \frac{29}{11} \quad z = -\frac{1}{11}$$

9.3.1. Solution Types in Linear Systems: Unique, Infinite, or None

1. Unique Solution

$$\left[\begin{array}{cccc|c} 1 & a_{12} & a_{13} & a_{14} & b_1 \\ 0 & 1 & a_{23} & a_{24} & b_2 \\ 0 & 0 & 1 & a_{34} & b_3 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

2. No Solution

$$0 = a$$

3. No Unique Solution (Infinite Number of Solutions)

Column 2 and 4 indicate free variables x_2 and x_4 because they have no pivot entries

$$\left[\begin{array}{cccc|c} 1 & a_{12} & a_{13} & a_{14} & b_1 \\ 0 & 0 & 1 & a_{24} & b_2 \\ 0 & 0 & 1 & a_{34} & b_3 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

9.3.2. Special Cases

Rows of all zeros appear in row echelon form (REF) in the following situations:

1. Dependent Equations

Some equations are multiples or linear combinations of others

$$2x + 4y = 8$$

$$x + 2y = 4$$

Augmented matrix:

$$\left[\begin{array}{cc|c} 2 & 4 & 8 \\ 1 & 2 & 4 \end{array} \right]$$

After row reduction:

$$\left[\begin{array}{cc|c} 1 & 2 & 4 \\ 0 & 0 & 0 \end{array} \right]$$

2. Underdetermined Systems

Number of variables is greater than the number of independent equations

$$x + y + z = 2$$

$$2x + 3y + z = 5$$

Augmented matrix:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 2 & 3 & 1 & 5 \end{array} \right]$$

After row reduction:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 2 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

3. Inconsistent Systems

No solution exists

Rows of zeros on the left side (coefficients of the variables) and a non-zero entry on the right side (augmented column)

$$\begin{aligned}x + y &= 3 \\2x + 2y &= 7\end{aligned}$$

Augmented matrix:

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 2 & 2 & 7 \end{array} \right]$$

After row reduction:

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 0 & 0 & 1 \end{array} \right]$$

10. Matrices

$m \times n$ matrix A

- m : rows
- n : columns

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

10.1. Matrix-Vector Products

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$A\vec{x} = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

For the dot product to be defined, the number of columns in the matrix A (which is n) must match the number of elements in the vector \vec{x} (also n).

The result of multiplying matrix A and vector \vec{x} will be a column vector with dimensions $m \times 1$, where m is the number of rows in the matrix A

$$(m \times n) \cdot (n \times 1) = m \times 1$$

1. As Row vectors

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$\vec{a}^T = [a_1, a_2, \dots, a_n]$$

$$\vec{b}^T = [b_1, b_2, \dots, b_n]$$

$$\mathbf{A} = \begin{bmatrix} [a_1, a_2, \dots, a_n] \\ [b_1, b_2, \dots, b_n] \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \vec{a} \\ \vec{b} \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$\begin{bmatrix} \vec{a}^T \\ \vec{b}^T \end{bmatrix} \cdot \vec{x} = \begin{bmatrix} \vec{a} \cdot \vec{x} \\ \vec{b} \cdot \vec{x} \end{bmatrix}$$

2. As Column Vectors

$$\vec{a} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

$$\vec{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} & \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} \end{bmatrix}$$

$$\mathbf{A} = [\vec{a} \ \vec{b}]$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\mathbf{A}\vec{x} = x_1\vec{a} + x_2\vec{b}$$

10.2. Null Space

The null space (or kernel) of a matrix \mathbf{A} is the set of all vectors \vec{x} that satisfy the equation:

$$A\vec{x} = \mathbf{0}$$

Where:

- A : $m \times n$ matrix
- \vec{x} : n -dimensional vector
- $\mathbf{0}$: zero vector in \mathbb{R}^m

$$N(A) = N(\text{rref}(A)) = \text{span}(\vec{v}_1, \vec{v}_2, \vec{v}_3)$$

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{bmatrix}$$

We want to find the null space of A , which consists of all vectors $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$ that satisfy:

$$A\vec{x} = \mathbf{0}$$

This expands to the following system of linear equations:

$$\begin{cases} 1x_1 + 1x_2 + 1x_3 + 1x_4 = 0 \\ 1x_1 + 2x_2 + 3x_3 + 4x_4 = 0 \\ 4x_1 + 3x_2 + 2x_3 + 1x_4 = 0 \end{cases}$$

This can be represented as the augmented matrix:

$$\left[\begin{array}{cccc|c} 1 & 1 & 1 & 1 & 0 \\ 1 & 2 & 3 & 4 & 0 \\ 4 & 3 & 2 & 1 & 0 \end{array} \right]$$

10.2.1. Column Space

The **column space** (or range) of matrix A is span of its columns vectors

If the matrix A has columns $\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n$, then the column space of A is defined as:

$$\text{Col}(A) = \{\vec{y} \in \mathbb{R}^m \mid \vec{y} = A\vec{x} \text{ for some } \vec{x} \in \mathbb{R}^n\}$$

or equivalently,

$$\text{Col}(A) = \text{span}(\{\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n\})$$

Consider the simple example of a 2×2 matrix:

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix}$$

The matrix has two columns:

$$\vec{a}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \quad \text{and} \quad \vec{a}_2 = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$$

The column space, denoted $\text{Col}(A)$, is the span of these two vectors:

$$\text{Col}(A) = \text{span}\left(\left\{\begin{bmatrix} 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 6 \end{bmatrix}\right\}\right)$$

Finding the Column Space

We observe that the two columns \vec{a}_1 and \vec{a}_2 are **linearly dependent**:

$$\vec{a}_2 = k\vec{a}_1$$

This means that \vec{a}_2 is a scalar multiple of \vec{a}_1 , the the two columns are **linearly dependent**. As a result, the column space is spanned by just one vector, \vec{a}_1 , because any linear combination of \vec{a}_1 and \vec{a}_2 can be reduced to a multiple of \vec{a}_1 .

Therefore, the column space of A is:

$$\text{Col}(A) = \text{span}\left(\left\{\begin{bmatrix} 1 \\ 3 \end{bmatrix}\right\}\right)$$

which represents all vectors of the form:

$$c \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} c \\ 3c \end{bmatrix} \quad \text{for any scalar } c$$

In other words, the column space is a line in \mathbb{R}^2 through the origin in the direction of

Rank of A

The rank of A , which is the **dimension of its column space**, is 1 because there is only one linearly independent column

This means the column space is the span of the columns of A , or all vectors that can be formed by taking linear combinations of the columns of A .

10.2.2. Dimension of a Subspace

Number of elements in a basis for the subspace

10.2.3. Nullity

Dimension of the Null Space

$$\dim(N(A))$$

The nullity of A : number of non-pivot columns (i.e., free variables) in the rref of A

10.2.4. Rank

Dimension of the column space

$$\text{rank}(A) = \dim(C(A))$$

10.2.5. Matrix Representation of Systems of Equations

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1m}x_m &= b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2m}x_m &= b_2 \\ &\vdots \\ a_{n1}x_1 + a_{n2}x_2 + \dots + a_{nm}x_m &= b_n \end{aligned}$$

Coefficient Matrix (A):

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{bmatrix}$$

Variable Vector (x):

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix}$$

Constant Vector (b):

$$\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$A\mathbf{x} = \mathbf{b}$$

The system of equations:

$$2x_1 + 3x_2 + 5x_3 = 100$$

$$4x_1 + 2x_2 + 1x_3 = 80$$

$$1x_1 + 5x_2 + 2x_3 = 60$$

Can be represented as a matrix equation:

$$\begin{bmatrix} 2 & 3 & 5 \\ 4 & 2 & 1 \\ 1 & 5 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 100 \\ 80 \\ 60 \end{bmatrix}$$

10.3. Matrix Multiplication

$m \times n$ matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{bmatrix}$$

$n \times p$ matrix:

$$B = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1p} \\ a_{21} & a_{22} & \dots & a_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{np} \end{bmatrix}$$

Compute Each Element of Result Matrix C

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

Let A be an $n \times m$ matrix:

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

Let B be a $p \times n$ matrix:

$$B = \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$$

Calculate Each Element of C

$$c_{11} = (1 \cdot 7) + (2 \cdot 9) + (3 \cdot 11) = 58$$

$$c_{12} = (1 \cdot 8) + (2 \cdot 10) + (3 \cdot 12) = 64$$

$$c_{21} = (4 \cdot 7) + (5 \cdot 9) + (6 \cdot 11) = 138$$

$$c_{22} = (4 \cdot 8) + (5 \cdot 10) + (6 \cdot 12) = 154$$

C is a $m \times p$ matrix

$$C = \begin{bmatrix} 58 & 64 \\ 139 & 154 \end{bmatrix}$$


10.4. Linear Transformation

10.4.1. Functions

A function f that maps elements from a set X (the domain) to a set Y (the codomain):

$$f : X \rightarrow Y$$

- Domain: The set X contains all possible inputs for the function f
- Codomain: The set Y is the space where all possible outputs of f reside, though not every element in Y must be an output of f



If

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

is defined by

$$f(x) = x^2$$

$$f : x \mapsto x^2$$

then:

- **Domain:** $X = \mathbb{R}$, any real number $((-\infty, \infty))$
- **Codomain:** $Y = \mathbb{R}$, any real number $((-\infty, \infty))$
- **Range:** the subset of the codomain $(\mathbb{R}), [0, \infty)$

10.5. Vector Transformation

A function f that maps an n -dimensional vector in \mathbb{R}^n to an m -dimensional vector in \mathbb{R}^m :

1. Function Definition

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

This means f takes as input a vector in \mathbb{R}^n (an n -dimensional space of real numbers) and maps it to a vector in \mathbb{R}^m (an m -dimensional space of real numbers).

2. Input Vector \vec{x}

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad \text{where } x_1, x_2, \dots, x_n \in \mathbb{R}$$

Here, \vec{x} is an n -dimensional vector, and each component x_i is a real number

3. Input Vector \vec{y}

$$\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_m \end{bmatrix} \quad \text{where } y_1, y_2, \dots, y_m \in \mathbb{R}$$

The output \vec{y} is an m -dimensional vector, and each component y_i is also a real number

Summary



The function f takes an n -dimensional vector of real numbers as input and produces an m -dimensional vector of real numbers as output

$$f(x_1, x_2, x_3) = (x_1 + 2x_2, 3x_3)$$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$$

$$f\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}\right) = \begin{bmatrix} x_1 + 2x_2 \\ 3x_3 \end{bmatrix}$$

$$f\left(\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 3 \end{bmatrix}$$



10.6. Linear Transformation

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$\vec{a}, \vec{b} \in \mathbb{R}^n$$

For a transformation **linear** it must satisfy two conditions:

1. Additivity (or linearity of addition)

$$T(\vec{a} + \vec{b}) = T(\vec{a}) + T(\vec{b})$$

Let's consider a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 2x \\ 3y \end{bmatrix}$$

Now let's take two vectors in \mathbb{R}^2 :

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \vec{b} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

Then the additivity property can be verified as follows:

1. First, find $T(\vec{a}) + T(\vec{b})$ separately:

$$T(\vec{a}) = T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 2 \cdot 1 \\ 3 \cdot 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$$

$$T(\vec{b}) = T\left(\begin{bmatrix} 3 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 2 \cdot 3 \\ 3 \cdot 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$$

$$T(\vec{a}) + T(\vec{b}) = \begin{bmatrix} 2 \\ 6 \end{bmatrix} + \begin{bmatrix} 6 \\ 3 \end{bmatrix} = \begin{bmatrix} 2+6 \\ 6+3 \end{bmatrix} = \begin{bmatrix} 8 \\ 9 \end{bmatrix}$$

2. Next, find $T(\vec{a} + \vec{b})$:

$$\vec{a} + \vec{b} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

$$T(\vec{a} + \vec{b}) = T\left(\begin{bmatrix} 4 \\ 3 \end{bmatrix}\right) = \begin{bmatrix} 2 \cdot 4 \\ 3 \cdot 3 \end{bmatrix} = \begin{bmatrix} 8 \\ 9 \end{bmatrix}$$

Since $T(\vec{a} + \vec{b}) = T(\vec{a}) + T(\vec{b})$, this confirms the additivity (linearity of addition) property of the transformation T

2. Homogeneity (or linearity of scalar multiplication):

$$T(c\vec{a}) = cT(\vec{a})$$

Let's consider a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 2x \\ 3y \end{bmatrix}$$

Now let's take one vectors in \mathbb{R}^2 :

$$\vec{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

and a scalar $c = 3$

Then the homogeneity property can be verified as follows:

1. First, find $cT(\vec{a})$

$$T(\vec{a}) = T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 2 \cdot 1 \\ 3 \cdot 2 \end{bmatrix} = \begin{bmatrix} 2 \\ 6 \end{bmatrix}$$

$$cT(\vec{a}) = 3 \cdot \begin{bmatrix} 2 \\ 6 \end{bmatrix} = \begin{bmatrix} 3 \cdot 2 \\ 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 6 \\ 18 \end{bmatrix}$$

2. Then, find $T(c\vec{a})$ by c to get $c\vec{a}$

$$c\vec{a} = 3 \cdot \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 3 \cdot 1 \\ 3 \cdot 2 \end{bmatrix} = \begin{bmatrix} 3 \\ 6 \end{bmatrix}$$

$$T(c\vec{a}) = T\left(\begin{bmatrix} 3 \\ 6 \end{bmatrix}\right) = \begin{bmatrix} 2 \cdot 3 \\ 3 \cdot 6 \end{bmatrix} = \begin{bmatrix} 6 \\ 18 \end{bmatrix}$$

Since $T(c\vec{a}) = cT(\vec{a})$, this confirms the homogeneity (linearity of scalar multiplication) property of the transformation T

10.7. Matrix Vector Products

Matrix product with vector is always a linear transformation

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$T(\vec{x}) = A\vec{x}$$

$$A = \begin{bmatrix} v_1 & v_2 & \dots & v_n \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$A\vec{x} = \begin{bmatrix} v_1 & v_2 & \dots & v_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1 v_1 + x_2 v_2 + \dots + x_n v_n$$

1. Additivity (or linearity of addition)

$$T(\vec{a} + \vec{b}) = T(\vec{a}) + T(\vec{b})$$

$$\begin{aligned} A \cdot (\vec{a} + \vec{b}) &= A \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ \vdots \\ a_n + b_n \end{bmatrix} = (a_1 + b_1)v_1 + (a_2 + b_2)v_2 + \dots + (a_n + b_n)v_n \\ &= a_1 v_1 + b_1 v_1 + a_2 v_2 + b_2 v_2 + \dots + a_n v_n + b_n v_n \\ &= (a_1 v_1 + a_2 v_2 + \dots + a_n v_n) + (b_1 v_1 + b_2 v_2 + \dots + b_n v_n) \\ &= A \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} + A \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix} \end{aligned}$$

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \quad u = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad v = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

1. Calculate $A(u + v)$

$$\begin{aligned} u + v &= \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} 4 \\ 6 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} A(u + v) &= \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 6 \end{bmatrix} \\ &= \begin{bmatrix} (2 \cdot 1) + (1 \cdot 6) \\ (0 \cdot 4) + (3 \cdot 6) \end{bmatrix} \\ &= \begin{bmatrix} 14 \\ 18 \end{bmatrix} \end{aligned}$$

2. Calculate $Au + Av$

$$\begin{aligned}
 A\mathbf{u} &= \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \\
 &= \begin{bmatrix} (2 \cdot 1) + (1 \cdot 2) \\ (0 \cdot 1) + (3 \cdot 2) \end{bmatrix} \\
 &= \begin{bmatrix} 4 \\ 6 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 A\mathbf{v} &= \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\
 &= \begin{bmatrix} (2 \cdot 3) + (1 \cdot 4) \\ (0 \cdot 3) + (3 \cdot 4) \end{bmatrix} \\
 &= \begin{bmatrix} 10 \\ 12 \end{bmatrix}
 \end{aligned}$$

$$\begin{aligned}
 A\mathbf{u} + A\mathbf{v} &= \begin{bmatrix} 4 \\ 6 \end{bmatrix} + \begin{bmatrix} 10 \\ 12 \end{bmatrix} \\
 &= \boxed{\begin{bmatrix} 14 \\ 18 \end{bmatrix}}
 \end{aligned}$$

2. Homogeneity (or linearity of scalar multiplication):

$$T(c\vec{a}) = cT(\vec{a})$$

$$\begin{aligned}
 A \cdot (c\vec{a}) &= \begin{bmatrix} v_1 & v_2 & \dots & v_n \end{bmatrix} \begin{bmatrix} ca_1 \\ ca_2 \\ \vdots \\ ca_n \end{bmatrix} \\
 &= ca_1v_1 + ca_2v_2 + \dots + ca_nv_n \\
 &= c \underbrace{(a_1v_1 + a_2v_2 + \dots + a_nv_n)}_{A\vec{a}}
 \end{aligned}$$

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \quad \mathbf{v} = \begin{bmatrix} 3 \\ 4 \end{bmatrix} \quad c = 5$$

1. Calculate $A(c\mathbf{v})$

$$c\mathbf{v} = 5 \cdot \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 15 \\ 20 \end{bmatrix}$$

$$\begin{aligned}
 A(c\mathbf{v}) &= \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 15 \\ 20 \end{bmatrix} \\
 &= \begin{bmatrix} (2 \cdot 15) + (1 \cdot 20) \\ (0 \cdot 15) + (3 \cdot 20) \end{bmatrix} \\
 &= \boxed{\begin{bmatrix} 50 \\ 60 \end{bmatrix}}
 \end{aligned}$$

2. Calculate $c(A\mathbf{v})$

$$\begin{aligned} A\mathbf{v} &= \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} (2 \cdot 3) + (1 \cdot 4) \\ (0 \cdot 3) + (3 \cdot 4) \end{bmatrix} \\ &= \begin{bmatrix} 10 \\ 12 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} c(A\mathbf{v}) &= 5 \cdot \begin{bmatrix} 10 \\ 12 \end{bmatrix} \\ &= \boxed{\begin{bmatrix} 50 \\ 60 \end{bmatrix}} \end{aligned}$$

10.8. Linear transformations as matrix vector products

The $n \times n$ matrix I_n :

$$I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}$$

$$\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

$$I_n \vec{x} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Standard Basis

$$I_n \vec{x} = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \\ \underbrace{}_{e_1} & \underbrace{}_{e_2} & \underbrace{}_{e_3} & & \underbrace{}_{e_n} \end{bmatrix}$$

$\{e_1, e_2, \dots, e_n\}$ is the standard basis for \mathbb{R}^n

$$\begin{aligned}
I_n \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} &= a_1 \vec{e}_1 + a_2 \vec{e}_2 + \dots + a_n \vec{e}_n \\
&= a_1 \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} + a_2 \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix} + \dots + a_n \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} \\
&= \begin{bmatrix} a_1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ a_2 \\ \vdots \\ 0 \end{bmatrix} + \dots + \begin{bmatrix} 0 \\ 0 \\ \vdots \\ a_n \end{bmatrix}
\end{aligned}$$

10.9. Image of a subset under transformation

$$\vec{x}_0 = \begin{bmatrix} -2 \\ -2 \end{bmatrix} \quad \vec{x}_1 = \begin{bmatrix} -2 \\ 2 \end{bmatrix} \quad \vec{x}_2 = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

$$L_0 = \{\vec{x}_0 + t(\vec{x}_1 - \vec{x}_0) \mid 0 \leq t \leq 1\}$$

$$L_1 = \{\vec{x}_1 + t(\vec{x}_2 - \vec{x}_1) \mid 0 \leq t \leq 1\}$$

$$L_2 = \{\vec{x}_2 + t(\vec{x}_0 - \vec{x}_2) \mid 0 \leq t \leq 1\}$$

The triangle T can be defines as the set of these points:

$$S = \{L_0, L_1, L_2\}$$

Let's define a transformation

$$T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


$$T(\vec{x}) = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$\begin{aligned}
T(L_0) &= \{T(\vec{x}_0 + t(\vec{x}_1 - \vec{x}_0)) \mid 0 \leq t \leq 1\} \\
&= \{T(\vec{x}_0) + T(t(\vec{x}_1 - \vec{x}_0)) \mid 0 \leq t \leq 1\} \\
&= \{T(\vec{x}_0) + tT(\vec{x}_1 - \vec{x}_0) \mid 0 \leq t \leq 1\} \\
&= \{T(\vec{x}_0) + tT(\vec{x}_1) - T(\vec{x}_0) \mid 0 \leq t \leq 1\}
\end{aligned}$$

$$T(\vec{x}_0) = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} -2 \\ -2 \end{bmatrix} = \begin{bmatrix} 0 \\ -4 \end{bmatrix}$$

$$T(\vec{x}_1) = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} -4 \\ -4 \end{bmatrix}$$

$$T(\vec{x}_2) = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 2 \\ -2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \end{bmatrix}$$



$T(L_0)$ is the image of L_0 under T

$T(S)$ is the image of S under T

10.10. Image of a transformation

The **image** of a transformation T is defined as:

$$\text{im}(T) = \{T(\vec{x}) \mid \vec{x} \in \mathbb{R}^n\}$$

or, equivalently,

$$T(\mathbb{R}^n)$$

$T(\mathbb{R}^n)$ is the image of \mathbb{R}^2 under T

This is the set of all possible outputs when T is applied to vectors in \mathbb{R}^n

Understanding $T(\mathbb{R}^n)$

1. Whole space transformation

The image of \mathbb{R}^n under T is the complete set of transformed vectors, often denoted as $\text{im}(T)$

2. Subset Transformation

For any subset $V \subseteq \mathbb{R}^n$, the image of V under T is the set of transformed vectors from V

Matrix representation of T

If T is represented by a $m \times n$ matrix A , then:

$$T(\vec{x}) = \{A\vec{x} \mid \vec{x} \in \mathbb{R}^n\}$$

where:

- A is the matrix associated with T
- $\vec{x} \in \mathbb{R}^n$ represents a vector in the input space

Transformation in terms of columns of A

If $A = \begin{bmatrix} \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \end{bmatrix}$, then for $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$:


$$A\vec{x} = x_1\vec{a}_1 + x_2\vec{a}_2 + \dots + x_n\vec{a}_n$$

Column Space of A

The image of T (or $\text{im}(T)$) is the column space of A :

$$C(A) = \text{span}(\vec{a}_1, \vec{a}_2, \dots, \vec{a}_n)$$

This is the set of all possible linear combinations of the columns of A , and thus represents all possible outputs of the transformation T



Suppose we have a matrix A :

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix}$$

Matrix A defines the transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that for any vector $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2$, the image under T is:

$$T(\vec{x}) = A\vec{x} = \begin{bmatrix} 2 & 1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Calculating Transformation

To see what T does to vectors in \mathbb{R}^2 , let's compute a few specific examples:

1. for $\vec{x} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$

$$T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = A\begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

2. for $\vec{x} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = A\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Image of T

The image of T , $\text{im}(T)$, is the set of all linear combinations of the vectors $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$:

$$\text{im}(T) = \text{span}\left(\begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \end{bmatrix}\right)$$

Thus, any vector in the image of T can be written as:

$$y = x_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 2x_1 + x_2 \\ x_1 + 3x_2 \end{bmatrix}$$

where $x_1, x_2 \in \mathbb{R}$.

Column space interpretation

The image of T is all the vectors in \mathbb{R}^2 that can be formed as linear combinations of $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

10.11. Preimage of a set

The preimage of a set S under a function T , denoted $T^{-1}(S)$, is the set of all elements in the domain of T that map to elements in S under the transformation T .

If $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a function from a set \mathbb{R}^n to a set \mathbb{R}^m , and $S \subseteq \mathbb{R}^m$ is a subset of a target space, then the preimage of S under T is:

$$T^{-1}(S) = \{\vec{x} \in \mathbb{R}^n \mid T(\vec{x}) \in S\}$$

This means that $T^{-1}(S)$ consists of all elements in \mathbb{R}^n that, when transformed by T , end up in S .

For any subset $S \subseteq \mathbb{R}^m$, the preimage $T^{-1}(S)$ collects all points in the domain that end up in S after applying T . If S is a single point, the preimage will be the set of all points in the domain that map to that specific point (this could be empty, a single point, or even a set of points, depending on the function).

Consider the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by the matrix:

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

This transformation T maps any vector $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ in \mathbb{R}^2 to:

$$T(\vec{x}) = A\vec{x} = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2x_1 \\ 3x_2 \end{bmatrix}$$

Now, let's find the preimage of a subset $S \subseteq \mathbb{R}^2$. Suppose we want the preimage of the set $S = \left\{ \begin{bmatrix} 4 \\ 6 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right\}$

The preimage of S under T , denoted $T^{-1}(S)$, consists of all vectors $\vec{x} \in \mathbb{R}^2$ such that $T(\vec{x}) \in \left\{ \begin{bmatrix} 4 \\ 6 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right\}$. To find this, we solve for \vec{x} in both cases: $T(\vec{x}) = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$ and $T(\vec{x}) = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$

Preimage of $\begin{bmatrix} 4 \\ 6 \end{bmatrix}$

$$A\vec{x} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

or

$$\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 4 \\ 6 \end{bmatrix}$$

Solving each component:

1. $2x_1 = 4 \Rightarrow x_1 = 2$
2. $3x_2 = 6 \Rightarrow x_2 = 2$

Thus, the preimage of S is the single point:

$$T^{-1}\left(\begin{bmatrix} 4 \\ 6 \end{bmatrix}\right) = \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix} \right\}$$

Preimage of $\begin{bmatrix} 2 \\ 3 \end{bmatrix}$

$$A\vec{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

or

$$\begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$$

Solving each component:

1. $2x_1 = 2 \Rightarrow x_1 = 1$
2. $3x_2 = 3 \Rightarrow x_2 = 1$

Thus, the preimage of S is the single point:

$$T^{-1}\left(\begin{bmatrix} 2 \\ 3 \end{bmatrix}\right) = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

Preimage of the set S

Since $S = \left\{ \begin{bmatrix} 4 \\ 6 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \end{bmatrix} \right\}$, the preimage of S is the union of the preimage of each vector in S :

$$T^{-1}(S) = \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$$

10.11.1. Kernel of a Transformation

The kernel of a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, denoted $\ker(T)$, is the set of all vectors in \mathbb{R}^n that T maps to the zero vector in \mathbb{R}^m . Formally, we define the kernel as:

$$\ker(T) = \{ \vec{x} \in \mathbb{R}^n \mid T(\vec{x}) = \vec{0} \}$$

The consists of all vectors that are “annihilated” by T , resulting in the zero vector after applying T .

The transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by the matrix

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$$

so that $T(\vec{x}) = A\vec{x} = \begin{bmatrix} 2x_1 \\ 3x_2 \end{bmatrix}$ for any $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^2$

To find the kernel of T , we need to find all the vectors $\vec{x} \in \mathbb{R}^2$ that satisfy:

$$T(\vec{x}) = \vec{0} \Rightarrow \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

This leads to the system of equations:

1. $2x_1 = 0 \Rightarrow x_1 = 0$

$$2. \quad 3x_2 = 0 \Rightarrow x_2 = 0$$

Thus, the only solution is $\vec{x} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$

10.11.2. Kernel and Null Space

The kernel of T

$$\ker(T) = \text{Null}(A)$$

10.12. Sum and Scalar Multiples of Linear Transformation

10.12.1. Sum

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m \quad S : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$(T + S) : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$A = \begin{bmatrix} \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \end{bmatrix} \quad B = \begin{bmatrix} \vec{b}_1 & \vec{b}_2 & \dots & \vec{b}_n \end{bmatrix}$$

$$\begin{aligned} (T + S)(\vec{x}) &= T(\vec{x}) + S(\vec{x}) \\ &= A\vec{x} + B\vec{x} \\ &= x_1\vec{a}_1 + x_2\vec{a}_2 + \dots + x_n\vec{a}_n + x_1\vec{b}_1 + x_2\vec{b}_2 + \dots + x_n\vec{b}_n \\ &= x_1(\vec{a}_1 + \vec{b}_1) + x_2(\vec{a}_2 + \vec{b}_2) + \dots + x_n(\vec{a}_n + \vec{b}_n) \\ &= \begin{bmatrix} \vec{a}_1 + \vec{b}_1 & \vec{a}_2 + \vec{b}_2 & \dots & \vec{a}_n + \vec{b}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= (A + B)\vec{x} \end{aligned}$$

10.12.2. Scalar Multiplication

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$cT : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

$$\begin{aligned} (cT)(\vec{x}) &= c(T(\vec{x})) \\ &= c(x_1\vec{a}_1 + x_2\vec{a}_2 + \dots + x_n\vec{a}_n) \\ &= x_1c\vec{a}_1 + x_2c\vec{a}_2 + \dots + x_nc\vec{a}_n \\ &= \begin{bmatrix} c\vec{a}_1 & c\vec{a}_2 & \dots & c\vec{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \\ &= cA\vec{x} \end{aligned}$$

10.13. Projection

The projection of a vector \vec{x} , onto a line L , denoted as $\text{Proj}_L(\vec{x})$, is a vector that lies on the line L , such that the difference between \vec{x} and its projection, $\text{Proj}_L(\vec{x}) - \vec{x}$, is orthogonal to L

$\text{Proj}_L(\vec{x})$ can be seen as the “shadow” cast by \vec{x} onto L when light shines perpendicularly to L .

$$\text{Proj}_L(\vec{x}) = c\vec{v} = \left(\frac{\vec{x} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \right) \vec{v}$$

Equivalently

$$\text{Proj}_L(\vec{x}) = c\vec{v} = \left(\frac{\vec{x} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v}$$

Where:

- \vec{v} is a direction vector for the line L
- $c = \frac{\vec{x} \cdot \vec{v}}{\vec{v} \cdot \vec{v}}$ is a scalar


$$\vec{x} = \begin{bmatrix} 2 \\ 3 \end{bmatrix}, \quad \vec{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

1. Define the line L as all vectors of the form $c\vec{v}$, where c is a scalar:

$$\begin{aligned} L &= \{c\vec{v} \mid c \in \mathbb{R}\} \\ &= \left\{ c \begin{bmatrix} 2 \\ 1 \end{bmatrix} \mid c \in \mathbb{R} \right\} \end{aligned}$$

2. Compute the projection using

$$\begin{aligned} \text{Proj}_L(\vec{x}) &= \frac{\vec{x} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \vec{v} \\ &= \frac{\begin{bmatrix} 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}}{\begin{bmatrix} 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \\ &= \frac{7}{5} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 2.8 \\ 1.4 \end{bmatrix} \end{aligned}$$



Projection as a Transformation

As a matrix vector product

$$L = \{c\vec{v} \mid c \in \mathbb{R}\}$$

$$\text{Proj}_L : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\vec{v} \cdot \vec{v} = \|\vec{v}\|^2$$

If \vec{v} is a unit vector:

$$\|\vec{v}\| = 1$$

Then


$$\begin{aligned} \text{Proj}_L(\vec{x}) &= \left(\frac{\vec{x} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \right) \vec{v} \\ &= \left(\frac{\vec{x} \cdot \vec{v}}{\|\vec{v}\|^2} \right) \vec{v} \end{aligned}$$

If we redefine our line L as all the scalar multiples of our unit vector \hat{u} :

$$L = \{c\hat{u} \mid c \in \mathbb{R}\}$$

Simplifies to:

$$(\vec{x} \cdot \hat{u})\hat{u}$$



Projection as a Linear Transformation

Let $\hat{u} \in \mathbb{R}^n$ be a unit vector:

$$\hat{u} = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix}, \quad \text{where } \|\hat{u}\| = 1$$

The projection matrix A is:

$$A = \hat{u}\hat{u}^T = \begin{bmatrix} u_1 \\ u_2 \\ \vdots \\ u_n \end{bmatrix} [u_1 \quad u_2 \quad \dots \quad u_n]$$

Expands to:

$$A = \begin{bmatrix} u_1 u_1 & u_1 u_2 & \dots & u_1 u_n \\ u_2 u_1 & u_2 u_2 & \dots & u_2 u_n \\ \vdots & \vdots & \ddots & \vdots \\ u_n u_1 & u_n u_2 & \dots & u_n u_n \end{bmatrix}$$

For any $\vec{x} \in \mathbb{R}^n$, the projection of \vec{x} onto the line spanned by \hat{u} is:

$$\text{Proj}_L(\vec{x}) = A\vec{x} = (\hat{u} \cdot \vec{x})\hat{u}$$

Consider a vector \vec{v} in \mathbb{R}^2 :

$$\vec{v} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

1. Construct the Unit Vector

$$\|\vec{v}\| = \sqrt{2^2 + 1^2} = \sqrt{5}$$

$$\hat{u} = \frac{\vec{v}}{\|\vec{v}\|} = \frac{1}{\sqrt{5}} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} \end{bmatrix}$$

This unit vector \hat{u} defines the line L , which consists of all the scalar multiples of \vec{v} :

$$L = \{c\vec{v} \mid c \in \mathbb{R}\}$$

2. Derive the Projection Matrix

The projection of any vector \vec{x} onto the line L is given by:

$$\text{Proj}_L(\vec{x}) = A\vec{x}$$


Where A is the projection matrix. To construct A we use the formula:

$$\begin{aligned} A &= \hat{u}\hat{u}^T \\ &= \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \begin{bmatrix} u_1 & u_2 \end{bmatrix} \\ &= \begin{bmatrix} u_1 \cdot u_1 & u_1 \cdot u_2 \\ u_2 \cdot u_1 & u_2 \cdot u_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{2}{\sqrt{5}} \\ \frac{1}{\sqrt{5}} \end{bmatrix} \begin{bmatrix} \frac{2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{2}{\sqrt{5}}\right)^2 & \frac{1}{\sqrt{5}} \frac{2}{\sqrt{5}} \\ \frac{2}{\sqrt{5}} \frac{1}{\sqrt{5}} & \left(\frac{1}{\sqrt{5}}\right)^2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix} \end{aligned}$$

3. Applying the Projection

To project any vector \vec{x} onto L , we multiply \vec{x} by the matrix A :

$$\begin{aligned} \text{Proj}_L(\vec{x}) &= A\vec{x} \\ &= \begin{bmatrix} \frac{4}{5} & \frac{2}{5} \\ \frac{2}{5} & \frac{1}{5} \end{bmatrix} \vec{x} \end{aligned}$$



1. Additivity of Projections (Linearity with respect to addition)

$$\begin{aligned}
 \text{Proj}_L(\vec{a} + \vec{b}) &= ((\vec{a} + \vec{b}) \cdot \hat{u})\hat{u} \\
 &= (\vec{a} \cdot \hat{u} + \vec{b} \cdot \hat{u})\hat{u} \\
 &= (\vec{a} \cdot \hat{u})\hat{u} + (\vec{b} \cdot \hat{u})\hat{u} \\
 &= \text{Proj}_L(\vec{a}) + \text{Proj}_L(\vec{b})
 \end{aligned}$$

2. Homogeneity of Projections (Linearity with respect to scalar multiplication)

$$\begin{aligned}
 \text{Proj}_L(c\vec{a}) &= (c\vec{a} \cdot \hat{u})\hat{u} \\
 &= c(\vec{a} \cdot \hat{u})\hat{u} \\
 &= c\text{Proj}_L(\vec{a})
 \end{aligned}$$

General Properties of A

1. Idempotence

$$A^2 = A$$

2. Symmetry

$$A^T = A$$

3. Rank

$$\text{rank}(A) = 1$$


Because $\hat{u}\hat{u}^T$ projects onto a one-dimensional subspace spanned by \hat{u}

10.14. Composition of Linear Transformations

$$S : X \rightarrow Y \quad T : Y \rightarrow Z$$

$$T \circ S : X \rightarrow Z$$

$$S(\vec{x}) = \underbrace{A}_{m \times n} \vec{x} \quad T(\vec{x}) = \underbrace{B}_{l \times m} \vec{x}$$



Consider two linear transformations T and S , where:

- T maps $\mathbb{R}^m \rightarrow \mathbb{R}^l$
- S maps $\mathbb{R}^n \rightarrow \mathbb{R}^m$

The composition $T \circ S$ is a linear transformation mapping $\mathbb{R}^n \rightarrow \mathbb{R}^l$ defined by:

$$T \circ S(\vec{x}) = T(S(\vec{x}))$$

Key Properties of $T \circ S$

1. Additivity

$$\begin{aligned}
 T \circ S(\vec{x} + \vec{y}) &= T(S(\vec{x} + \vec{y})) \\
 &= T(S(\vec{x}) + S(\vec{y})) \\
 &= T(S(\vec{x})) + T(S(\vec{y})) \\
 &= T \circ S(\vec{x}) + T \circ S(\vec{y})
 \end{aligned}$$

2. Homogeneity

$$\begin{aligned}
 T \circ S(c\vec{x}) &= T(S(c\vec{x})) \\
 &= T(cS(\vec{x})) \\
 &= cT(S(\vec{x})) \\
 &= c(T \circ S)(\vec{x})
 \end{aligned}$$

Matrix Representation of $T \circ S$

Let S be represented by the matrix A ($m \times n$), and let T be represented by the matrix B ($l \times m$)

For a vector $\vec{x} \in \mathbb{R}^n$

$$\begin{aligned}
 T \circ S(\vec{x}) &= T(S(\vec{x})) = T(A\vec{x}) \\
 &= \underbrace{B}_{l \times m} \left(\underbrace{A}_{m \times n} \vec{x} \right) \\
 &= \underbrace{C}_{l \times n} \vec{x}
 \end{aligned}$$

The composition $T \circ S$ is therefore represented by the matrix $C = A \cdot B$, where C is of size $l \times n$

Column-Wise Interpretation

The matrix A can be decomposed column-wise:

$$A = \begin{bmatrix} \vec{a}_1 & \vec{a}_2 & \dots & \vec{a}_n \end{bmatrix}$$

where \vec{a}_i is the i -th column of A and I_n is the identity matrix in \mathbb{R}^n , and its columns are the standard basis vectors $\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n$.

$$I_n = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \\ \underbrace{}_{\vec{e}_1} & \underbrace{}_{\vec{e}_2} & & \underbrace{}_{\vec{e}_n} \end{bmatrix}$$

To compute C :

1. For each \vec{e}_i in the basis of \mathbb{R}^n , $A\vec{e}_i = \vec{a}_i$, the i -th column of A

$$\begin{aligned} C &= \begin{bmatrix} B(A\vec{e}_1) & B(A\vec{e}_2) & \dots & B(A\vec{e}_n) \end{bmatrix} \\ &= \begin{bmatrix} B\left(A \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}\right) & B\left(A \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}\right) & \dots & B\left(A \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}\right) \end{bmatrix} \\ &= \begin{bmatrix} B\vec{a}_1 & B\vec{a}_2 & \dots & B\vec{a}_n \end{bmatrix} \end{aligned}$$

The composition $T \circ S$ is the linear map represented by $C = B \cdot A$

Each column of C reflects how T transforms the action of S on a standard basis vector

10.15. Matrix Product

$$\underbrace{A}_{m \times n} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \quad \underbrace{B}_{n \times p} = \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1p} \\ b_{21} & b_{22} & \dots & b_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \dots & b_{np} \end{bmatrix}$$

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

$$\underbrace{A}_{2 \times 3} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \quad \underbrace{B}_{3 \times 2} = \begin{bmatrix} 7 & 8 \\ 9 & 10 \\ 11 & 12 \end{bmatrix}$$

$$AB = \left[A \begin{bmatrix} 7 \\ 9 \\ 11 \end{bmatrix} \quad A \begin{bmatrix} 8 \\ 10 \\ 12 \end{bmatrix} \right]$$


$$= \left[\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 7 \\ 9 \\ 11 \end{bmatrix} \quad \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \begin{bmatrix} 8 \\ 10 \\ 12 \end{bmatrix} \right]$$

$$= \left[\begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \begin{bmatrix} 7 \\ 9 \\ 11 \end{bmatrix} \quad \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} \begin{bmatrix} 8 \\ 10 \\ 12 \end{bmatrix} \right]$$

10.16. Matrix Product Associativity

10.17. Eigenvectors

10.17.1. Transformation



$$A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}$$

$$T(\hat{i}) = A\hat{i}$$

$$= \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

$$T(\hat{j}) = A\hat{j}$$

$$= \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$\begin{aligned}
T(\vec{x}) &= A\vec{x} \\
&= \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \\
&= \begin{bmatrix} 4 \\ 2 \end{bmatrix}
\end{aligned}$$

10.18. Eigenvalues

10.18.1. LU Decomposition

Given a matrix A , LU decomposition aims to express A as:

$$A = LU$$

Where:

- L : Lower triangular matrix (all elements above the diagonal are zero)
- U : Upper triangular matrix (all elements below the diagonal are zero)

1. Solve $Ly = b$ Using Forward Substitution

$$Ly = b$$

Where:

- L : Lower triangular matrix (all elements above the diagonal are zero)
- y : Intermediate vector we are solving for
- b : Right-hand side vector

$$\begin{pmatrix} L_{11} & 0 & 0 \\ L_{21} & L_{22} & 0 \\ L_{31} & L_{32} & L_{33} \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$$

- First row: $L_{11}y_1 = b_1$, so $y_1 = \frac{b_1}{L_{11}}$
- Second row: $L_{21}y_1 + L_{22}y_2 = b_2$, substitute y_1 into this equation and solve for y_2 :

$$y_2 = \frac{b_2 - L_{21}y_1}{L_{22}}$$

- Third row: $L_{31}y_1 + L_{32}y_2 + L_{33}y_3 = b_3$, substitute y_1 and y_2 into this equation, solve for y_3 :

$$y_3 = \frac{b_3 - L_{31}y_1 - L_{32}y_2}{L_{33}}$$

2. Solve $Ux = y$ Using Backward Substitution

$$Ux = y$$

Where:

- U : Upper triangular matrix (all elements below the diagonal are zero)
- y : Vector of unknowns (solution)
- b : Vector computed from the forward substitution step

$$\begin{pmatrix} U_{11} & U_{12} & U_{13} \\ 0 & U_{22} & U_{23} \\ 0 & 0 & U_{33} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$$

- Third row: $U_{33}x_3 = y_3$, so $x_3 = \frac{y_3}{U_{33}}$
- Second row: $U_{22}x_2 + U_{23}x_3 = y_2$, substitute x_3 from the previous step and solve for x_2 :

$$x_2 = \frac{y_2 - U_{23}x_3}{U_{22}}$$

- First row: $U_{11}x_1 + U_{12}x_2 + U_{13}x_3 = y_1$, substitute x_2 and x_3 from the previous step and solve for x_1 :

$$x_1 = \frac{y_1 - U_{12}x_2 - U_{13}x_3}{U_{11}}$$

$$A = \begin{pmatrix} 2 & 3 & 1 \\ 4 & 7 & 3 \\ 6 & 18 & 5 \end{pmatrix} \quad b = \begin{pmatrix} 5 \\ 12 \\ 31 \end{pmatrix}$$

1. Factor A into L and U :

$$L = \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 6 & 1 \end{pmatrix} \quad U = \begin{pmatrix} 2 & 3 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{pmatrix}$$

2. Solve $Ax = b$

- $y_1 = \frac{b_1}{L_{11}} = \frac{5}{1} = 5$
- $y_2 = \frac{b_2 - L_{21}y_1}{L_{22}} = \frac{12 - 2 \times 5}{1} = 2$
- $y_3 = \frac{b_3 - L_{31}y_1 - L_{32}y_2}{L_{33}} = \frac{31 - 4 \times 5 - 3 \times 2}{1} = 5$

So,

$$y = \begin{pmatrix} 5 \\ 2 \\ 5 \end{pmatrix}$$

3. Solve $Ux = y$

- $x_3 = \frac{y_3}{U_{33}} = \frac{5}{-2} = -2.5$
- $x_2 = \frac{y_2 - U_{23}x_3}{U_{22}} = \frac{2 - 1 \times -2.5}{1} = 4.5$
- $x_1 = \frac{y_1 - U_{12}x_2 - U_{13}x_3}{U_{11}} = \frac{5 - 3 \times 4.5 - 1 \times -2.5}{2} = -4$

So,

$$x = \begin{pmatrix} -4 \\ 4.5 \\ -2.5 \end{pmatrix}$$

10.19. Solving Systems of Linear Equations

Linear Equation

$$y = a_1x_1 + a_2x_2 + \dots + a_nx_n$$

1. Consistency

Whether a system of linear equations has at least one solution

Consistent System

$$x + y = 3$$

$$x - y = 1$$

This system has a unique solution

$$(x, y) = (2, 1)$$

Inconsistent System

$$x + y = 3$$

$$x + y = 5$$

This system is inconsistent (equations contradict each other, no solution can satisfy both)

2. Independence

Whether the equations in the system provide unique and non-redundant information about the variables

Independent Equations

$$x + y = 3$$

$$x - y = 1$$

Neither equation can be derived from the other (they provide unique information and intersect at a single point)

Dependent Equations

$$x + y = 3$$

$$2x + 2y = 6$$


Second equation is just a multiple of the first equation (they describe the same line)

3. Recognizing Systems with No Solution or Infinite Solutions


$$3x + 2y = 6 \quad (\text{Equation 1})$$

$$6x + 4y = 12 \quad (\text{Equation 2})$$


Unique Solution (Consistent and Independent):



No Solution (Inconsistent):



Infinitely Many Solutions (Consistent and Dependent):



2. Matrix Representation

System of Equations

$$\begin{aligned}
 a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\
 a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\
 &\vdots \\
 a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m
 \end{aligned}$$

Matrix Representation

Coefficient vector (A)

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

Variable vector (x)

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Constant vector (b)

$$b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

Matrix equation

$$Ax = b$$

```

from scipy.linalg import solve

X = np.array([
    [1, 1, 1],
    [2, -1, 3],
    [3, 4, -1]
])

Y = np.array([6, 14, 1])

intersection_point = solve(X, Y)

```

10.19.1. Gaussian Elimination

Convert a matrix into its row echelon form (REF) or reduced row echelon form (RREF)

$$\begin{aligned}
 a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &= b_1 \\
 a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &= b_2 \\
 &\vdots \\
 a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m
 \end{aligned}$$

1. Create an augmented matrix

$$A = \left(\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & \ddots & \vdots & b_3 \\ a_{2m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right)$$

2. Forward Elimination

Eliminate the element in the i -th of the k -th column ($k > i$)

$$R_k \leftarrow R_k - \frac{a_{ki}}{a_{ii}} R_i$$

Where

- a_{ii} : Pivot element
- R_k : k -th row
- R_i : i -th row

3. Back Substitution

4. Reduced Row Echelon Form (RREF)

$$2x_1 + 3x_2 = 5$$

$$4x_1 + 5x_2 = 5$$

1. Create an augmented matrix

$$\left(\begin{array}{cc|c} 2 & 3 & 5 \\ 4 & 5 & 5 \end{array} \right)$$

2. Forward Elimination

$$R_k \leftarrow R_k - \frac{a_{ki}}{a_{ii}} R_i$$

$$R_k \leftarrow R_k - \frac{a_{21}}{a_{11}} R_1$$

$$R_2 \leftarrow R_2 - \frac{4}{2} R_1$$

$$R_2 \leftarrow R_2 - 2 \times R_1$$

$$\left(\begin{array}{cc|c} 2 & 3 & 5 \\ 4 - 2 \times 2 & 5 - 2 \times 3 & 5 - 2 \times 5 \end{array} \right)$$

Simplifies to:

$$\left(\begin{array}{cc|c} 2 & 3 & 5 \\ 0 & -1 & -4 \end{array} \right)$$

System is now:

$$2x_1 + 3x_2 = 5$$

$$-1x_2 = -4$$

3. Back Substitution

$$-1x_2 = -4$$

$$x_2 = 4$$

Substitute:

$$2x_1 + 3(4) = 5$$

$$2x_1 + 12 = 5$$

$$x_1 = -3.5$$

Solution:

$$x_1 = -3.5$$

$$x_2 = 4$$

10.19.2. Substitution

$$x + y = 10 \quad (\text{Equation 1})$$

$$2x - y = 5 \quad (\text{Equation 2})$$

1. Solve Equation 1 for y

$$y = 10 - x$$

2. Substitute into Equation 2

$$2x - (10 - x) = 5$$

3. Solve for x :

$$2x - 10 + x = 5$$

$$3x - 10 = 5$$

$$3x = 15$$

$$x = 5$$

4. Find y using value of x

$$y = 10 - x$$

$$y = 10 - 5$$

$$y = 5$$

10.19.3. Addition or Subtraction Method

$$3x + 2y = 12 \quad (\text{Equation 1})$$

$$2x - 2y = 4 \quad (\text{Equation 2})$$

1. Add the equations

$$(3x + 2y) + (2x - 2y) = 12 + 4$$

$$5x = 16$$

$$x = \frac{16}{5}$$

$$x = 3.2$$

2. Substitute

$$3(3.2) + 2y = 12$$

$$9.6 + 2y = 12$$

$$2y = 12 - 9.6y = \frac{2.4}{2}$$

$$y = 1.2$$

Calculus I

11. Cheatsheet

11.1. Limits

Epsilon-Delta

For every distance ε around L , there's a δ -range around a that keeps $f(x)$ within ε of L .

$$\lim_{x \rightarrow x_0} f(x) = L \iff \forall \varepsilon > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$$

The limit of $f(x)$ as x approaches x_0 equals L if and only if, for every $\varepsilon > 0$, there exists a $\delta > 0$ such that, whenever $0 < |x - x_0| < \delta$, implies that $|f(x) - L| < \varepsilon$

Limit Type	Name	Quantifiers
$x \rightarrow x_0, f(x) \rightarrow L$	Epsilon-Delta	$\forall \varepsilon, \exists \delta$
$x \rightarrow x_0, f(x) \rightarrow \infty$	M-Delta	$\forall M, \exists \delta$
$x \rightarrow \infty, f(x) \rightarrow L$	epsilon-N	$\forall \varepsilon, \exists N$
$x \rightarrow \infty, f(x) \rightarrow \infty$	M-N	$\forall M, \exists N$

Finite \rightarrow Finite (Epsilon-Delta)

$$\lim_{x \rightarrow x_0} f(x) = L \iff \forall \varepsilon > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$$

Finite \rightarrow Infinity (M-Delta)

$+\infty$

$$\lim_{x \rightarrow x_0} f(x) = +\infty \iff \forall M > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - x_0| < \delta \Rightarrow f(x) > M$$

$-\infty$

$$\lim_{x \rightarrow x_0} f(x) = -\infty \iff \forall M > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - x_0| < \delta \Rightarrow f(x) < -M$$

Infinity \rightarrow Finite (Epsilon-N)

$+\infty$

$$\lim_{x \rightarrow +\infty} f(x) = L \iff \forall \varepsilon > 0, \exists N > 0 \text{ s.t. } x > N \Rightarrow |f(x) - L| < \varepsilon$$

$-\infty$

$$\lim_{x \rightarrow +\infty} f(x) = L \iff \forall \varepsilon > 0, \exists N > 0 \text{ s.t. } x < -N \Rightarrow |f(x) - L| < \varepsilon$$

Infinity \rightarrow Infinity (M-N)

$$\lim_{x \rightarrow \pm\infty} f(x) = \pm\infty \iff \forall M > 0, \exists N > 0 \text{ s.t. } x < N \Rightarrow f(x) > M$$

11.2. Derivatives

$$\frac{df}{dx}(x) = f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Let $f(x) = x^2$

a. Find $f'(x)$

$$f'(x) = \boxed{2x}$$

b. Prove a

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{(x+h) - x} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{h(2x + h)}{h} \\ &= \lim_{h \rightarrow 0} (2x + h) \\ &= \lim_{h \rightarrow 0} (2x + 0) \\ &= \boxed{2x} \end{aligned}$$

c. Prove b

$$\lim_{x \rightarrow x_0} f(x) = L \iff \forall \varepsilon > 0, \exists \delta > 0 \text{ s.t. } 0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon$$

p.f.:

Let $\varepsilon > 0$

Choose $\delta = \varepsilon$

Suppose $0 < |h - 0| < \delta$

Check

$$\left| \frac{(x+h)^2 - x^2}{h} - 2x \right|$$

$$\begin{aligned}
&= \left| \frac{x^2 + 2xh + h^2 - x^2}{h} - 2x \right| \\
&= \left| \frac{h(2x + h)}{h} - 2x \right| \\
&= |2x + h - 2x| \\
&= |h| < \delta = \varepsilon
\end{aligned}$$

11.3. Integrals

11.4. Differential

If you have a function:

$$y = f(x)$$

Then the differential du is defined as:

$$dy = f'(x) \cdot dx$$

This means:

- $f'(x) = \frac{dy}{dx}$ is the derivative
- dx is a small change in x
- dy is the corresponding small change in y

So:

$$dy = \frac{dy}{dx} \cdot dx$$

Let's say“

$$y = x^2$$

Then:

$$\frac{dy}{dx} = 2x$$

$$\frac{d}{dx} \left(\int f(x) dx \right) = f(x)$$

$$\int \left(\frac{d}{dx} f(x) \right) dx = f(x)$$

Rule	$\frac{d}{dx}$ Rule	$\frac{d}{dx}$ Example	\int Rule	\int Example
Constant	$\frac{d}{dx}[c] = 0$	$\frac{d}{dx}[7] = 0$	$\int c dx = cx + C$	$\int 7 dx = 7x + C$
Power	$\frac{d}{dx}[x^n] = nx^{n-1}$	$\frac{d}{dx}[x^4] = 4x^3$	$\int x^n dx = \frac{x^{n+1}}{n+1} + C$ ($n \neq -1$)	$\int x^3 dx = \frac{x^4}{4} + C$
Constant Multiple	$\frac{d}{dx}[cf] = cf'$	$\frac{d}{dx}[3x^2] = 3 \cdot 2x$ $= 6x$	$\int cf dx = c \int f dx$	$\int 3x^2 dx = 3 \cdot \frac{x^3}{3}$ $= x^3 + C$
Sum	$\frac{d}{dx}[f + g] = f' + g'$	$\frac{d}{dx}(x^2 + x)$ $= \frac{d}{dx}(x^2) + \frac{d}{dx}(x)$ $= 2x + 1$	$\int (f + g) dx$ $= \int f dx + \int g dx$	$\int (x^2 + x) dx$ $= \int x^2 dx + \int x dx$ $= \frac{x^3}{3} + \frac{x^2}{2} + C$
Difference	$\frac{d}{dx}[f - g] = f' - g'$	$\frac{d}{dx}(x^2 - x)$ $= \frac{d}{dx}(x^2) - \frac{d}{dx}(x)$ $= 2x - 1$	$\int (f - g) dx$ $= \int f dx - \int g dx$	$\int (x^2 - x) dx$ $= \int x^2 dx - \int x dx$ $= \frac{x^3}{3} - \frac{x^2}{2} + C$
Product	$\frac{d}{dx}[fg] = f'g + fg'$	$\frac{d}{dx}[x \sin x]$ $= 1 \cdot \sin x + x \cdot \cos x$	Integration by Parts $\int u dv = uv - \int v du$	
Quotient	$\frac{d}{dx} \left[\frac{f}{g} \right] = \frac{f'g - fg'}{g^2}$	$\frac{d}{dx} \left[\frac{x^2}{x+1} \right]$ $= \frac{2x(x+1) - x^2(1)}{(x+1)^2}$ $= \frac{x^2 + 2x}{(x+1)^2}$	Algebraic Manipulation / Substitution	
Chain	$\frac{d}{dx}[f(g(x))]$ $= f'(g(x)) \cdot g'(x)$	$\frac{d}{dx}[\sin(x^2)]$ $= \cos(x^2) \cdot 2x$	Integration by Substitution $\int f(g(x))g'(x) dx$ $= \int f(u) du$	

Rule	$\frac{d}{dx}$ Rule	$\frac{d}{dx}$ Example	\int Rule	\int Example
Exponential	$\frac{d}{dx}[e^x] = e^x$	$\frac{d}{dx}[e^x] = e^x$	$\int e^x dx = e^x + C$	$\int e^x dx = e^x + C$
	$\frac{d}{dx}[a^x] = a^x \ln(a)$	$\frac{d}{dx}[2^x] = 2^x \ln 2$	$\int a^x dx = \frac{a^x}{\ln a} + C$	$\int 2^x dx = \frac{2^x}{\ln 2} + C$
Logarithmic	$\frac{d}{dx}[\ln(x)] = \frac{1}{x}$	$\frac{d}{dx}[\ln(x)] = \frac{1}{x}$	$\int \ln(x) dx$ $= x \ln x - x + C$	$\int \ln(x) dx$ $= x \ln x - x + C$
	$\frac{d}{dx}[\log_a(x)] = \frac{1}{x \ln(a)}$	$\frac{d}{dx}[\log_2(x)] = \frac{1}{x \ln(2)}$	$\int \log_a(x) dx =$ $= \frac{x \ln x}{\ln a} - \frac{x}{\ln a} + C$	$\int \log_{10}(x) dx$ $= \frac{x \ln x}{\ln 10} - \frac{x}{\ln 10} + C$
Sin	$\frac{d}{dx}[\sin(x)] = \cos(x)$	$\frac{d}{dx}[\sin(x)] = \cos(x)$	$\int \sin(x) dx$ $= -\cos x + C$	$\int \sin(x) dx$ $= -\cos x + C$
Cos	$\frac{d}{dx}[\cos(x)] = -\sin(x)$	$\frac{d}{dx}[\cos(x)] = -\sin(x)$	$\int \cos(x) dx$ $= \sin x + C$	$\int \cos(x) dx$ $= \sin x + C$
Tan	$\frac{d}{dx}[\tan(x)]$ $= \sec^2(x)$	$\frac{d}{dx}[\tan(x)]$ $= \sec^2(x)$	$\int \tan(x) dx$ $= -\ln \cos x + C$	$\int \tan(x) dx$ $= -\ln \cos x + C$

11.5. Product Rule \longrightarrow Integration by Parts

$$\int u \, dv = uv - \int v \, du$$

Given a function:

$$f(x) = x \cdot e^x$$

Integrate (by parts):

$$\int x \cdot e^x \, dx$$

Step 1: Choose u and dv

We choose:

- $u = x$ (easy to **differentiate**)
- $dv = e^x$ (easy to **integrate**)

Step 2: Compute du and v

- $u = x \longrightarrow du = dx$
- $dv = e^x \, dx \longrightarrow v = \int e^x \, dx = e^x$

Step 3: Plug into formula

$$\begin{aligned}\int x \cdot e^x \, dx &= u \cdot v - \int v \cdot du \\ &= x \cdot e^x - \int e^x \cdot dx\end{aligned}$$

Step 4: Compute the remaining integral:

$$\int e^x \, dx = e^x$$

Step 5: Finish the expression:

$$\begin{aligned}\int x \cdot e^x &= x \cdot e^x - e^x + C \\ &= e^x(x - 1) + C\end{aligned}$$

11.6. Chain Rule \longrightarrow u -Substitution

$$\int f(g(x))g'(x) \, dx = \int f(u) \, du$$

11.7. Quotient Rule \longrightarrow Algebraic Manipulation / Substitution

12. Limits & Continuity

13. Properties of Limits

13.1. Continuous

13.1.1. Addition, Subtraction, Multiplication, Division

$$\lim_{x \rightarrow c} (f(x) * g(x)) = \lim_{x \rightarrow c} f(x) * \lim_{x \rightarrow c} g(x)$$
$$* \in \{+, -, \times, \div\}$$



13.1.2. Constant

$$\lim_{x \rightarrow c} k f(x) = k \lim_{x \rightarrow c} f(x)$$

13.2. Non-continuous

Even though the limit for either function may not exist, their * can exist as long as

$$\lim_{x \rightarrow c^-} (f(x) * g(x)) = \lim_{x \rightarrow c^+} (f(x) * g(x))$$
$$* \in \{+, -, \times, \div\}$$



Problem 1: Limit Exists

$$\lim_{x \rightarrow -2} (f(x) + g(x))$$

1. Left-hand limit ($x \rightarrow -2^-$)

$$\lim_{x \rightarrow -2^-} f(x) = 1 \qquad \lim_{x \rightarrow -2^-} g(x) = 3$$

Adding these

$$\begin{aligned} \lim_{x \rightarrow -2^-} (f(x) + g(x)) &= \lim_{x \rightarrow -2^-} f(x) + \lim_{x \rightarrow -2^-} g(x) \\ &= 1 + 3 \\ &= 4 \end{aligned}$$

2. Right-hand limit ($x \rightarrow -2^+$)

$$\lim_{x \rightarrow -2^+} f(x) = 3 \qquad \lim_{x \rightarrow -2^+} g(x) = 1$$

Adding these

$$\begin{aligned} \lim_{x \rightarrow -2^+} (f(x) + g(x)) &= \lim_{x \rightarrow -2^+} f(x) + \lim_{x \rightarrow -2^+} g(x) \\ &= 3 + 1 \\ &= 4 \end{aligned}$$

3. Since both the left-hand and right-hand limits agree

$$\lim_{x \rightarrow -2} (f(x) + g(x)) = 4$$

Problem 2: Limit Does Not Exist

$$\lim_{x \rightarrow 1} (f(x) + g(x))$$

1. Left-hand limit ($x \rightarrow 1^-$)

$$\lim_{x \rightarrow 1^-} f(x) = 2 \qquad \lim_{x \rightarrow 1^-} g(x) = 0$$

Adding these

$$\begin{aligned} \lim_{x \rightarrow 1^-} (f(x) + g(x)) &= \lim_{x \rightarrow 1^-} f(x) + \lim_{x \rightarrow 1^-} g(x) \\ &= 2 + 0 \\ &= 2 \end{aligned}$$

2. Right-hand limit ($x \rightarrow 1^+$)

$$\lim_{x \rightarrow 1^+} f(x) = -1 \qquad \lim_{x \rightarrow 1^+} g(x) = 0$$

Adding these

$$\begin{aligned} \lim_{x \rightarrow 1^+} (f(x) + g(x)) &= \lim_{x \rightarrow 1^+} f(x) + \lim_{x \rightarrow 1^+} g(x) \\ &= -1 + 0 \\ &= -1 \end{aligned}$$



3. Since both the left-hand and right-hand limits do not agree, the limit $\lim_{x \rightarrow c} (f(x) + g(x))$ does not exist

13.3. Composite Functions

$$\lim_{x \rightarrow c} f(g(x)) = f\left(\lim_{x \rightarrow c} g(x)\right)$$

For this to hold true, two important conditions must be satisfied:

- **Inner limit exists:** The limit $\lim_{x \rightarrow c}$ must exist and equal some value L . That is, as x gets arbitrarily close to c , $g(x)$ approaches a well-defined number L
- **Continuity of the outer function:** The function f must be continuous at the point L . Continuity ensures that f behaves predictably near L , without any jumps, gaps, or undefined points



Problem 1: Inner Limit & Continuity Exist

$$\lim_{x \rightarrow -3} f(g(x))$$

1. Inner limit $\lim_{x \rightarrow -3} g(x)$

Observing $g(x)$, as $x \rightarrow -3$, $g(x) \rightarrow 3$. The inner limit $L = 3$ exists

$$\lim_{x \rightarrow -3} g(x) = 3$$

2. Continuity of $f(x)$ at $x = 3$

Observing $f(x)$, $f(3) = -1$. Since $f(x)$ is continuous at $x = 3$, the composite limit holds

$$\begin{aligned} f\left(\lim_{x \rightarrow -3} g(x)\right) &= f(3) \\ &= -1 \end{aligned}$$

Problem 2: Inner Limit Does Not Exist

$$\lim_{x \rightarrow 2} f(g(x))$$

1. Inner limit $\lim_{x \rightarrow 2} g(x)$

Observing $g(x)$, as $x \rightarrow 2$, $g(x) \rightarrow 2$. The inner limit does not exist

Problem 3: Continuity Does Not Exist

$$\lim_{x \rightarrow 0.5} f(g(x))$$

1. Inner limit $\lim_{x \rightarrow 0.5} g(x)$

Observing $g(x)$, as $x \rightarrow 0.5$, $g(x) \rightarrow 1$. The inner limit $L = 1$ exists

$$\lim_{x \rightarrow 0.5} g(x) = 1$$

2. Continuity of $f(x)$ at $x = 1$

Observing $f(x)$, $f(1)$ is not continuous. Since $f(x)$ is not continuous at $x = 1$, the composite limit does not hold

13.4. Limits by Direct Substitution

Limit exists

$$\begin{aligned}\lim_{x \rightarrow -1} (6x^2 + 5x - 1) &= 6(-1)^2 + 5(-1) - 1 \\ &= 6 - 5 - 1 \\ &= 0\end{aligned}$$

Limit does not exist (Undefined)

$$\begin{aligned}\lim_{x \rightarrow 1} \frac{x}{\ln(x)} &= \frac{1}{\ln(1)} \\ &= \frac{1}{0}\end{aligned}$$

13.4.1. Limits of Piecewise Functions

13.4.2. Absolute Value

13.5. Limits by Factoring

13.6. Limits by Rationalizing

13.7. Continuity & Differentiability at a Point

Piecewise function:

$$f(x) = \begin{cases} x^2 & \text{if } x < 3 \\ 6x - 9 & \text{if } x \geq 3 \end{cases}$$

1. Check for **Continuity**

- Value of $f(3)$

$$f(3) = 6(3) - 9 = 9$$

- Left-Hand Limit (LHL)

$$\lim_{x \rightarrow 3^-} f(x) = 3^2 = 9$$

- Right-Hand Limit (RHL)

$$\lim_{x \rightarrow 3^+} f(x) = 6(3) - 9 = 9$$

Since $f(3) = \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x)$, $f(x)$ is **continuous** at $x = 3$

2. Check for **Differentiability**

- Left-Hand Derivative (LHD)

$$\begin{aligned}
 \lim_{x \rightarrow 3^-} \frac{f(x) - f(3)}{x - 3} &= \frac{x^2 - 3^2}{x - 3} \\
 &= \frac{x^2 - 9}{x - 3} \\
 &= \frac{(x + 3)(x - 3)}{x - 3} \\
 &= x + 3 \\
 &= \boxed{6}
 \end{aligned}$$

- Right-Hand Derivative

$$\begin{aligned}
 \lim_{x \rightarrow 3^+} \frac{f(x) - f(3)}{x - 3} &= \frac{(6x - 9) - 3^2}{x - 3} \\
 &= \frac{6x - 9 - 9}{x - 3} \\
 &= \frac{6x - 18}{x - 3} \\
 &= \frac{6(x - 3)}{x - 3} \\
 &= \boxed{6}
 \end{aligned}$$

Since the left-hand and right-hand derivatives are equal, $f(x)$ is **differentiable** at $x = 3$

Conclusion: $f(x)$ is both continuous & differentiable at $x = 3$

13.8. Power Rule

$$\begin{aligned}
 f(x) &= x^n, \quad n \neq 0 \\
 f'(x) &= nx^{n-1}
 \end{aligned}$$

$$\begin{aligned}
 f(x) &= x^3 \\
 f'(x) &= 3x^2
 \end{aligned}$$

$$\begin{aligned}
 \frac{d}{dx}(\sqrt[3]{x^2}) &= \frac{d}{dx}\left((x^2)^{\frac{1}{3}}\right) \\
 &= \frac{d}{dx}\left(x^{2 \times \frac{1}{3}}\right) \\
 &= \frac{d}{dx}\left(x^{\frac{2}{3}}\right) \\
 &= \frac{d}{dx}\left(\frac{2}{3}x^{-\frac{1}{3}}\right)
 \end{aligned}$$

13.9. Constant Rule

$$\frac{d}{dx}[k] = 0$$

$$\frac{d}{dx}[-3] = 0$$

13.10. Constant Multiple Rule

$$\begin{aligned}\frac{d}{dx}[kf(x)] &= k \frac{d}{dx}[f(x)] \\ &= kf'(x)\end{aligned}$$

$$\begin{aligned}\frac{d}{dx}[2x^5] &= 2 \frac{d}{dx}[x^5] \\ &= 2 \cdot 5x^4 \\ &= 10x^4\end{aligned}$$

13.11. Sum Rule

$$\begin{aligned}\frac{d}{dx}[f(x) + g(x)] &= \frac{d}{dx}[f(x)] + \frac{d}{dx}[g(x)] \\ &= f'(x) + g'(x)\end{aligned}$$

$$\begin{aligned}\frac{d}{dx}[x^3 + x^{-4}] &= \frac{d}{dx}[x^3] + \frac{d}{dx}[x^{-4}] \\ &= 3x^2 + (-4x^{-5}) \\ &= 3x^2 - 4x^{-5}\end{aligned}$$

13.12. Difference Rule

$$\begin{aligned}\frac{d}{dx}[f(x) - g(x)] &= \frac{d}{dx}[f(x)] - \frac{d}{dx}[g(x)] \\ &= f'(x) - g'(x)\end{aligned}$$

$$\begin{aligned}\frac{d}{dx}[x^4 - x^3] &= \frac{d}{dx}[x^4] - \frac{d}{dx}[x^3] \\ &= 4x^3 - 3x^2\end{aligned}$$

13.13. Square Root

$$\begin{aligned}\frac{d}{dx} \sqrt[4]{x} &= \frac{d}{dx} x^{\frac{1}{4}} \\ &= \frac{1}{4} x^{\frac{1}{4}-1} \\ &= \frac{1}{4} \cdot x^{-\frac{3}{4}} \\ &= \frac{1}{4} \cdot \frac{1}{x^{3/4}} \\ &= \frac{1}{4x^{3/4}}\end{aligned}$$

13.14. Derivative of a Polynomial

$$\begin{aligned}f(x) &= 2x^3 - 7x^2 + 3x - 100 \\ f'(x) &= 2 \cdot 3x^2 - 7 \cdot 2x + 3 + 0\end{aligned}$$

$$h(x) = 3f(x) + 2g(x)$$

Evaluate $\frac{d}{dx} h(x)$ at $x = 9$

$$\begin{aligned}\frac{d}{dx}(h(x)) &= \frac{d}{dx}(3f(x) + 2g(x)) \\ &= \frac{d}{dx} 3f(x) + \frac{d}{dx} 2g(x) \\ &= 3 \frac{d}{dx} f(x) + 2 \frac{d}{dx} g(x)\end{aligned}$$

Evaluate $h'(9)$

$$h'(9) = 3f'(9) + 2g'(9)$$

$$\begin{aligned}g(x) &= \frac{2}{x^3} - \frac{1}{x^2} \\ \frac{d}{dx}(g(x)) &= \frac{d}{dx}(2x^{-3} - 1x^{-2}) \\ g'(x) &= 2 \cdot (-3)x^{-4} - (-2)x^{-3} \\ &= -6x^{-4} + 2x^{-3}\end{aligned}$$


$$\begin{aligned}
 g'(2) &= -6 \cdot 2^{-4} + 2 \cdot 2^{-3} \\
 &= -\frac{6}{2^4} + \frac{2}{2^3} \\
 &= -\frac{3}{8} + \frac{2}{8} \\
 &= -\frac{1}{8}
 \end{aligned}$$

$$-9 \quad -8$$

$$\begin{aligned}
 f(x) &= x^3 - 6x^2 + x - 5 \\
 y &= mx + b \\
 f'(1) &= -8 \quad y = -8x + b \\
 -9 &= -8 \cdot 1 + b \\
 -9 &= -8 + b \\
 -9 + 8 &= -8 + 8 + b \\
 b &= -1
 \end{aligned}$$


Tangent line to $f(x)$ at $x = 1$

$$y = -8x - 1$$




13.15. Sin

$$\frac{d}{dx} \sin(x) = \cos(x)$$




13.16. Cos

$$\frac{d}{dx} \cos(x) = -\sin(x)$$




13.17. e^x

$$\frac{d}{dx} e^x = e^x$$



13.18. $\ln(x)$

$$\ln(x) = \frac{1}{x}$$



13.19. Product Rule

$$\frac{d}{dx}[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$$

$$\frac{d}{dx}[x^2 \sin(x)]$$

$$\begin{aligned} f(x) &= x^2 & g(x) &= \sin(x) \\ f'(x) &= 2x & g'(x) &= \cos(x) \end{aligned}$$

$$\frac{d}{dx}[x^2 \sin(x)] = 2x \sin(x) + x^2 \cos(x)$$

13.20. Quotient Rule

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$$


$$\frac{d}{dx} \left[\frac{x^2}{\cos(x)} \right]$$

$$\begin{aligned} f(x) &= x^2 & g(x) &= \cos(x) \\ f'(x) &= 2x & g'(x) &= -\sin(x) \end{aligned}$$

$$\frac{d}{dx} \left[\frac{x^2}{\cos(x)} \right] = \frac{2x \cos(x) - x^2(-\sin(x))}{[\cos(x)]^2}$$


13.20.1. $\tan(x)$

$$\begin{aligned} \frac{d}{dx}[\tan(x)] &= \frac{d}{dx} \left[\frac{\sin(x)}{\cos(x)} \right] \\ &= \frac{\cos(x) \cdot \cos(x) - \sin(x) \cdot -\sin(x)}{\cos^2(x)} \\ &= \frac{\cos^2(x) + \sin^2(x)}{\cos^2(x)} \\ &= \boxed{\frac{1}{\cos^2(x)}} \\ &= \boxed{\sec^2(x)} \end{aligned}$$




13.20.2. $\cot(x)$

$$\begin{aligned}
 \frac{d}{dx}[\cot(x)] &= \frac{d}{dx} \left[\frac{\cos(x)}{\sin(x)} \right] \\
 &= \frac{\cos(x) \cdot \cos(x) - \sin(x) \cdot -\sin(x)}{\cos^2(x)} \\
 &= \frac{-\sin^2(x) - \cos^2(x)}{\cos^2(x)} \\
 &= -\frac{1}{\sin^2(x)} \\
 &= -\csc^2(x)
 \end{aligned}$$




13.20.3. $\sec(x)$

$$\begin{aligned}
 \frac{d}{dx}[\sec(x)] &= \frac{d}{dx} \left[\frac{1}{\cos(x)} \right] \\
 &= \frac{0 \cdot \cos(x) - 1 \cdot -\sin(x)}{\cos^2(x)} \\
 &= \frac{0 + 1 \cdot \sin(x)}{\cos^2(x)} \\
 &= \frac{\sin(x)}{\cos^2(x)} \\
 &= \frac{\sin(x)}{\cos(x)} \cdot \frac{1}{\cos(x)} \\
 &= \tan(x) \cdot \sec(x)
 \end{aligned}$$



13.20.4. $\csc(x)$

$$\begin{aligned}
 \frac{d}{dx}[\csc(x)] &= \frac{d}{dx}\left[\frac{1}{\sin(x)}\right] \\
 &= \frac{0 \cdot \sin(x) - 1 \cdot \cos(x)}{\sin^2(x)} \\
 &= \frac{0 - 1 \cdot \cos(x)}{\sin^2(x)} \\
 &= \frac{-\cos(x)}{\sin^2(x)} \\
 &= -\frac{\cos(x)}{\sin(x)} \cdot \frac{1}{\sin(x)} \\
 &= \cot(x) \cdot \csc(x)
 \end{aligned}$$



13.21. Chain Rule

Compute the derivative of a composite function

$$\begin{aligned} h'(x) &= \frac{d}{dx}[f(g(x))] \\ &= f'(g(x)) \cdot g'(x) \end{aligned}$$

More generally:

$$y = f_1(f_2(f_3(\dots f_n(x)\dots)))$$

$$\frac{dy}{dx} = f'_1(f_2(f_3(\dots f_n(x)\dots))) \cdot f'_2(f_3(\dots f_n(x)\dots)) \cdot f'_3(\dots f_n(x)\dots) \cdot \dots \cdot f'_n(x)$$

$$\frac{d}{dx}[f(g(x))] = f'(g(x)) \cdot g'(x)$$

$$\frac{d}{dx}[e^{\sin(x)}]$$

$$\frac{d}{dx}[e^x] = \frac{1}{x} \quad \frac{d}{dx}[\sin(x)] = \cos(x)$$

$$f'(g(x)) = \frac{1}{\sin(x)} \quad g'(x) = \cos(x)$$

$$\frac{d}{dx} \left[\underbrace{\ln \left(\overbrace{\sin(x)}^{g(x)} \right)}_{f(g(x))} \right] = \frac{1}{\sin(x)} \cdot \cos(x)$$

$$f(x) = \cos^3(x) = (\cos(x))^3$$

$$f(x) = v(u(x))$$

$$f'(x) = v'(u(x)) \cdot u'(x)$$

$$\begin{aligned} f'(x) &= \frac{dv}{du} \cdot \frac{du}{dx} \\ &= \frac{d(\cos(x))^3}{d\cos(x)} \cdot \frac{d\cos(x)}{dx} \\ &= 3(\cos(x))^2 \cdot -\sin(x) \\ &= -3(\cos(x))^2 \sin(x) \end{aligned}$$

13.22. Implicit Differentiation

When a function is not explicitly solved for one variable in terms of another. E.g.:

$$x^2 + y^2 = 1$$

Instead of solving for y explicitly in terms of x , implicit differentiation allows you to differentiate both sides of an equation directly, treating y as an implicit function of x .

Steps for Implicit Differentiation

1. Differentiate both sides of the equation with respect to x , treating y as a function of x
2. Apply the chain rule whenever differentiating y , since $y = y(x)$
2. Solve for $\frac{dx}{dy}$

$$x^2 + y^2 = 1$$

1. Differentiate both sides


$$\begin{aligned} \frac{d}{dx}[x^2 + y^2] &= \frac{d}{dx}[1] \\ \frac{d}{dx}[x^2] + \frac{d}{dx}[y^2] &= \frac{d}{dx}[1] \\ \frac{d}{dx}[x^2] + \frac{d}{dx}[y^2] &= 0 \end{aligned}$$

2. Apply the chain rule to y^2

$$2x + 2y \frac{dy}{dx} = 0$$

3. Solve for $\frac{dy}{dx}$

$$\begin{aligned} 2y \cdot \frac{dy}{dx} &= -2x \\ \frac{dy}{dx} &= \frac{-2x}{2y} \\ \frac{dy}{dx} &= -\frac{x}{y} \end{aligned}$$



$$x^2 + y^2 = 5^2$$

$$2x dx + 2y dy = 0$$

$$\frac{dy}{dx} = -\frac{x}{y}$$

13.23. Derivatives of Inverse Functions

Finding the derivative of the inverse function $f^{-1}(x)$ at a given point directly from the function $f(x)$. Instead of explicitly computing the inverse function $f^{-1}(x)$, we use the inverse function derivative formula:


$$\frac{d}{dx}[f^{-1}(x)] = \frac{1}{f'(f^{-1}(x))}$$

This approach allows us to determine the derivative of the inverse function without needing to express $f^{-1}(x)$ explicitly. Instead, we find the value of x that satisfies $f(x) = a$ (where a is the given point), evaluate $f'(x)$, and apply the formula.

1. Definition of Inverse Function

$$f(f^{-1}(x)) = x$$

$$f^{-1}(f(x)) = x$$



2. Differentiate Both Sides

Differentiate both sides with respect to x

$$\frac{d}{dx}[f(f^{-1}(x))] = \frac{d}{dx}[x]$$

The right-hand side simplifies to:

$$\frac{d}{dx}[f(f^{-1}(x))] = 1$$

Using chain-rule:

$$\frac{d}{dx}[f(g(x))] = f'(g(x)) \cdot g'(x)$$

The left side expands as:

$$\frac{d}{dx}[f(f^{-1}(x))] = f'(f^{-1}(x)) \cdot \frac{d}{dx}[f^{-1}(x)]$$

This we get:

$$f'(f^{-1}(x)) \cdot \frac{d}{dx}[f^{-1}(x)] = 1$$

3. Solve for $\frac{d}{dx}[f^{-1}(x)]$

Rearrange to isolate $\frac{d}{dx}[f^{-1}(x)]$:

$$\frac{d}{dx}[f^{-1}(x)] = \frac{1}{f'(f^{-1}(x))}$$

Given the function:

$$f(x) = x^3$$

We want to find $\frac{d}{dx}f^{-1}(x)$ at $x = 0.5$ using the inverse function derivative formula:

$$\frac{d}{dx}f^{-1}(x) = \frac{1}{f'(f^{-1}(x))}$$

1. Compute $f'(x)$

Differentiate $f(x)$:

$$f'(x) = 3x^2$$

2. Solve for x such that $f(x) = 0.5$

We need to find x such that:

$$x^3 = 0.5$$

Solving for x :

$$x = \sqrt[3]{0.5}$$

3. Compute $f'(\sqrt[3]{0.5})$


Evaluate the derivative at $x = \sqrt[3]{0.5}$:

$$f'(\sqrt[3]{0.5}) = 3(\sqrt[3]{0.5})^2$$

4. Use the formula

$$\frac{d}{dx} f^{-1}(0.5) = \frac{1}{3(\sqrt[3]{0.5})^2}$$

5. Interpretation



The expression:

$$\frac{d}{dx} f^{-1}(0.5) = \frac{1}{3(\sqrt[3]{0.5})^2}$$

represents the derivative of the inverse function $f^{-1}(x)$ evaluated at $x = 0.5$. This means it gives the slope of the tangent line to the inverse function at $x = 0.5$.

```
from sympy import symbols, solve

# y = x**3 + x
x, y = symbols('x y')

f = x**3 + x - y

inverse = solve(f, x)

print(inverse)
```

13.23.1. Derivative Inverse Sin

$$\frac{d}{dx}[\sin^{-1}(x)] = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}[\arcsin(x)] = \frac{1}{\sqrt{1-x^2}}$$

13.23.2. Derivative Inverse Cos

$$\frac{d}{dx}[\cos^{-1}(x)] = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}[\arccos(x)] = -\frac{1}{\sqrt{1-x^2}}$$

13.23.3. Derivative Inverse Tan

$$\frac{d}{dx}[\tan^{-1}(x)] = \frac{1}{1+x^2}$$


$$\frac{d}{dx}[\arctan(x)] = \frac{1}{1+x^2}$$

13.24. Inverse Functions

1. Definition

$$f(f^{-1}(x)) = x$$

$$f^{-1}(f(x)) = x$$



A function $f : A \rightarrow B$ has an inverse function f^{-1} if and only if f is **bijective** (i.e., both one-to-one and onto):

- **Injective (One-to-One):**

$$f(x_1) = f(x_2) \text{ implies } x_1 = x_2$$

No two inputs map to the same output

- **Surjective (Onto):**

Every element in B is mapped from some element in A




2. Finding Inverse Function




To determine f^{-1} :

- Express y in terms of x : $y = f(x)$
- Solve for x in terms of y
- Swap x and y , remaining y as $f^{-1}(x)$

3. Graphical Representation

The graph of f^{-1} is a reflection of the graph of f across the line $x = y$





13.25. L'Hôpital's Rule

Evaluating limits that result in an indeterminate form like $\frac{0}{0}$ or $\frac{\infty}{\infty}$

If $\lim_{x \rightarrow a} f(x) = 0$ and $\lim_{x \rightarrow a} g(x) = 0$ (or both go to $\pm\infty$), and $f(x)$ and $g(x)$ are differentiable near a , then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Consider:

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x^2}$$

Step 1: Direct Substitution

Substituting $x = 0$:

$$\frac{1 - \cos(0)}{0^2} = \frac{1 - 1}{0} = \frac{0}{0}$$

Since this is an indeterminate form, we apply L'Hôpital's Rule.

Step 2: First Application of L'Hôpital's Rule

Differentiate the numerator and denominator:

- Numerator: $f(x) = 1 - \cos(x) \Rightarrow f'(x) = \sin(x)$
- Denominator: $g(x) = x^2 \Rightarrow g'(x) = 2x$

Thus, applying L'Hôpital's Rule:

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x^2} = \lim_{x \rightarrow 0} \frac{\sin(x)}{2x}$$

Step 3: Second Check for Indeterminate Form

Substituting $x = 0$:

$$\frac{\sin(0)}{2(0)} = \frac{0}{0}$$

Since this is still an indeterminate form, we apply L'Hôpital's Rule again.

Step 4: Second Application of L'Hôpital's Rule

Differentiate again:

- Numerator: $f'(x) = \sin(x) \Rightarrow f''(x) = \cos(x)$
- Denominator: $g'(x) = 2x \Rightarrow g''(x) = 2$

Applying L'Hôpital's Rule again:

$$\lim_{x \rightarrow 0} \frac{\sin(x)}{2x} = \lim_{x \rightarrow 0} \frac{\cos(x)}{2}$$

Step 5: Evaluate the Limit

Now, substituting $x = 0$:

$$\frac{\cos(0)}{2} = \frac{1}{2}$$

Final Answer:

$$\lim_{x \rightarrow 0} \frac{1 - \cos(x)}{x^2} = \frac{1}{2}$$

13.26. Mean Value Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function that satisfies the following conditions:

1. f is **continuous** on the closed interval $[a, b]$
2. f is **differentiable** on the open interval (a, b)

Then there exists at least one point $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

This means that the instantaneous rate of change (derivative) at some point c is equal to the average rate of change over the entire interval

Consider $f(x) = x^2$ on $[1, 3]$


- The average rate of change is:

$$\frac{f(3) - f(1)}{3 - 1} = \frac{9 - 1}{2} = 4$$

- The derivative for $f(x)$ is $f'(x) = 2x$
- Setting $f'(c) = 4$, we solve:

$$2c = 4 \Rightarrow c = 2$$

Thus, at $c = 2$, the instantaneous rate of change matches the average rate of change




13.27. Extreme Value Theorem

The Extreme Value Theorem states that if a function $f(x)$ is continuous on a closed interval $[a, b]$, then $f(x)$ must attain both a maximum and a minimum value within that interval. This means there exist points $c, d \in [a, b]$ such that:


$$f(c) \geq f(x) \quad \text{and} \quad f(d) \leq f(x) \quad \text{for all } x \in [a, b]$$

1. **Continuity:** The function must be continuous on $[a, b]$. Discontinuities (jumps, asymptotes, holes) can prevent the function from attaining an extreme value



2. **Closed Interval:** If the function is defined on an open interval (a, b) , an extremum may not exist

$f(x) = \frac{1}{x}$ on $(0, 1]$ has no maximum because it keeps increasing as $x \rightarrow 0$.





13.27.1. Critical points

A critical point of a function $f(x)$ is a point in the domain where either:

- $f'(x) = 0$
- $f'(x)$ is undefined

$f(x) = \frac{1}{x}$ on $(0, 1]$ has no maximum because it keeps increasing as $x \rightarrow 0$.



13.27.2. Global vs. Local Extrema

- $f(c)$ is a **relative maximum** if $f(c) \geq f(x)$ for all $x \in (c - h, c + h)$ for $h > 0$
- $f(d)$ is a **relative minimum** if $f(d) \leq f(x)$ for all $x \in (d - h, d + h)$ for $h > 0$

13.27.3. First and Second Derivative Tests

First Derivative Test and the Second Derivative Test are used to classify critical points, and both aim to determine whether the point is a local maximum, local minimum, or neither

13.27.3.1. First Derivative Test

If $f'(x)$ changes sign around a critical point c , we can determine if $f(c)$ is a local maximum or minimum:

- If $f'(x)$ changes from **positive to negative** at c , then $f(c)$ is a **local maximum**
- If $f'(x)$ changes from **negative to positive** at c , then $f(c)$ is a **local minimum**
- If $f'(x)$ does **not** change sign, $f(c)$ is **not** a local extremum

Function:

$$f(x) = x^3 - 3x$$

Derivative:

$$\begin{aligned} f'(x) &= 3x^2 - 3 \\ &= 3(x - 1)(x + 1) \end{aligned}$$

Critical points ($f'(x) = 0$):

- $x = -1$
- $x = 1$

Analysis of $f'(x)$

- Pick point left of -1 , say $x = -2$:

$$f'(-2) = 3(4 - 1) = 9 > 0 \rightarrow \text{increasing}$$

- Pick a point between -1 and 1 , say $x = 0$:


$$f'(0) = -3 < 0 \rightarrow \text{decreasing}$$

- Pick a point right of 1 , say $x = 2$:

$$f'(2) = 3(4 - 1) = 9 > 0 \rightarrow \text{increasing}$$

Conclusion

- At $x = -1$, $f'(x)$ changes from **positive to negative** \rightarrow **local maximum**
- At $x = 1$, $f'(x)$ changes from **negative to positive** \rightarrow **local minimum**



13.27.3.2. Second Derivative Test

If $f''(x)$ is continuous near a critical point c , and $f'(c) = 0$, then:

- If $f''(c) > 0$, then $f(c)$ is a **local minimum** (concave up)
- If $f''(c) < 0$, then $f(c)$ is a **local maximum** (concave down)
- If $f''(c) = 0$, the test is **inconclusive** — use the first derivative test or other methods

Function:

$$f(x) = x^4 - 4x^2$$

First Derivative:

$$f'(x) = 4x^3 - 8x$$

Second Derivative:

$$f''(x) = 12x^2 - 8$$

Critical points ($f'(x) = 0$):

- $x = 0$
- $x = \sqrt{2}$

- $x = -\sqrt{2}$


Evaluate $f''(x)$ at each point

- At $x = 0$

$$f''(0) = 12(0)^2 - 8 = -8 \rightarrow \text{concave down} \rightarrow \text{local maximum}$$

- At $x = \sqrt{2}$ or $x = -\sqrt{2}$

$$f''(\sqrt{2}) = f''(-\sqrt{2}) = 12(2)^2 - 8 = 16 \rightarrow \text{concave up} \rightarrow \text{local minimum}$$




13.28. Differentiation Rules

Power	$\frac{d}{dx}[x^n] = n \cdot x^{n-1}$
Sum	$\frac{d}{dx}[f(x) + g(x)] = f'(x) + g'(x)$
Product	$\frac{d}{dx}[f(x) \cdot g(x)] = f'(x)g(x) + f(x)g'(x)$
Quotient	$\frac{d}{dx}\left[\frac{f(x)}{g(x)}\right] = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$
Chain	$\frac{d}{dx}[f(g(x))] = f'(g(x)) \cdot g'(x)$

Calculus II

$$z = f(u, v)$$

$$u = f(x, y) \quad v = f(x, y)$$



The derivative of z with respect to x is the sum of all possible paths from z to x :

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial x}$$

Let:

- $u = x + y$
- $v = xy$
- $z = u^2 + \sin(v)$

We want to compute:

$$\frac{\partial z}{\partial x}$$

Step 1: Apply the multivariable chain rule

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial z}{\partial v} \frac{\partial v}{\partial x}$$

Step 2: Compute partial derivatives

- $\frac{\partial u}{\partial x} = \frac{\partial}{\partial x}(x + y) = 1 + 0 = 1$
- $\frac{\partial v}{\partial x} = \frac{\partial}{\partial x}(xy) = 1y = y$
- $\frac{\partial z}{\partial u} = \frac{\partial}{\partial u}(u^2 + \sin(v)) = 2u + 0 = 2u$
- $\frac{\partial z}{\partial v} = \frac{\partial}{\partial v}(u^2 + \sin(v)) = 0 + \cos(v) = \cos(v)$

Step 3: Plug in:

$$\begin{aligned} \frac{\partial z}{\partial x} &= 2u \cdot 1 + \cos(v) \cdot y \\ &= 2(x + y) + y \cos(xy) \end{aligned}$$

Calculus III

Statistics

14. Probability Theory

14.1. Probability Axioms

14.1.1. Non-Negativity

Probability of any event cannot be negative

$$P(A) \geq 0$$

14.1.2. Normalization

The probability of the sample space is always 1

$$P(S) = 1$$

Where:

- S is the sample space (the set of all possible outcomes)

14.1.3. Additivity

If two events are mutually exclusive (cannot happen at the same time), the probability of either occurring is the sum of their individual probabilities

$$P(A \cup B) = P(A) + P(B)$$

14.2. Rules

14.2.1. Complement Rule

The probability of the complement of an event A is $P(A^c) = 1 - P(A)$

How likely it is that the event does not occur.

Consider a fair die:

- Let A be the event “rolling a 4”. Then $P(A) = \frac{1}{6}$
- The complement of A , denoted $P(A^c)$, is “not rolling 4”
- $P(A^c) = 1 - P(A) = 1 - \frac{1}{6} = \frac{5}{6}$

14.2.2. Multiplication Rule

- Independent Events

If A and B are independent, then $P(A \cap B) = P(A) \times P(B)$. The probability of both events occurring is the product of their individual probabilities.

Consider flipping two fair coins. Let A be the event “the first coin is heads” and B be the event “the second coin is heads”

- Since the flips are independent, $P(A \cap B) = P(A) \times P(B)$
- $P(A) = \frac{1}{2}$ and $P(B) = \frac{1}{2}$
- Thus, $P(A \cap B) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$

- Dependent Events

If A and B are dependent, $P(A \cap B) = P(A) \times P(B | A)$, where $P(B | A)$ is the conditional probability of B given A .

Draw two cards from a standard deck without replacement. Let A be the event “drawing an Ace on the first draw” and B be the event “drawing an Ace on the second draw”

- $P(A) = \frac{4}{52} = \frac{1}{13}$
- If A occurs (i.e., an Ace is drawn first), there are 3 Aces left out of 51 cards. So, $P(B | A) = \frac{3}{51} = \frac{1}{17}$
- Thus, $P(A \cap B) = P(A) \times P(B | A) = \frac{1}{13} \times \frac{1}{17} = \frac{1}{221}$

14.2.3. Addition Rule

For Any Two Events: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$. This accounts for the overlap between the two events to avoid double counting.

Suppose you roll a die, and you want to find the probability of rolling a 2 or a 4.

- Let A be the event “rolling a 2,” and B be the event “rolling a 4”
- $P(A) = \frac{1}{6}$ and $P(B) = \frac{1}{6}$
- Since A and B are mutually exclusive, $P(A \cap B) = 0$
- $P(A \cup B) = P(A) + P(B) - P(A \cap B) = \frac{1}{6} + \frac{1}{6} - 0 = \frac{2}{6} = \frac{1}{3}$

14.2.4. Conditional Probability

The probability of an event A given that B has occurred is $P(A | B) = \frac{P(A \cap B)}{P(B)}$, provided $P(B) > 0$.

In a deck of 52 cards, if you know a card is a spade, what is the probability that it is an Ace?

- Let A be the event “drawing an Ace,” and B be the event “drawing a spade”
- $P(B) = \frac{13}{52} = \frac{1}{4}$
- There is 1 Ace of Spades out of 13 spades, so $P(A \cap B) = \frac{1}{52}$
- Thus, $P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{1}{52}}{\frac{13}{52}} = \frac{1}{13}$

14.2.5. Law of Total Probability

If $\{B_i\}$ is a partition of the sample space, then for any event A :

$$P(A) = \sum_i P(A \cap B_i) = \sum_i P(A | B_i) \times P(B_i)$$

Suppose you want to calculate the probability of raining on a given day. You know that it’s either sunny or cloudy, and the probability of rain is different in each condition.

- Let B_1 be the event “sunny” and B_2 be the event “cloudy”
- $P(B_1) = \frac{3}{4}$ and $P(B_2) = \frac{1}{4}$

- The probability of rain given sunny is $P(A | B_1) = \frac{1}{10}$, and given cloudy is $P(A | B_2) = \frac{2}{5}$
- The total probability of rain is:

$$\begin{aligned}
 P(A) &= P(A | B_1) \times P(B_1) + P(A | B_2) \times P(B_2) \\
 &= \frac{1}{10} \times \frac{3}{4} + \frac{2}{5} \times \frac{1}{4} \\
 &= \frac{3}{40} + \frac{2}{20} \\
 &= \frac{3}{40} + \frac{4}{40} \\
 &= \frac{7}{40}
 \end{aligned}$$

14.2.6. Law of Large Numbers

As the number of trials of a random experiment increases, the sample mean will converge to the expected value (mean) of the random variable

You flip a fair coin 1000 times. As you increase the number of flips, the proportion of heads should get closer to the probability of heads (which is 0.5).

14.2.7. Central Limit Theorem

For a sufficiently large number of trials, the distribution of the sample mean approaches a normal distribution, regardless of the distribution of the population

Suppose you are rolling a fair die repeatedly and calculating the average of the results. If you take 30 samples, each being the average of 10 die rolls, the distribution of those sample means will be approximately normal with a mean of 3.5 (the mean of a fair die) and a standard deviation that can be computed based on the original die's distribution.

14.3. Bayes Theorem

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

- $P(A)$: prior probability (initial belief about event A)
- $P(B | A)$: likelihood (probability of observing B given that A is true)
- $P(B)$: marginal likelihood (overall probability of observing B under all possible conditions)
- $P(A | B)$: posterior probability (updated belief after considering the evidence B)

15. Descriptive Statistics

15.1. Central Tendency

15.1.1. Mean

Sum of all the values divided by the number of values

$$\mu = \frac{\sum_{i=1}^n x_i}{n}$$

$$[1, 2, 3]$$

$$\bar{x} = \frac{1 + 2 + 3}{3} = 150$$

15.1.2. Median

Middle value in a set of values when they are arranged in ascending or descending order

1. **Odd** Number of Values

$$[1, 2, 3]$$

- **Step 1:** Arrange the Data in Ascending Order

$$[1, 2, 3]$$

- **Step 2:** Identify the Median

$$\text{Median} = 2$$

2. **Even** Number of Values

$$[1, 2, 3, 4]$$

- **Step 1:** Arrange the Data in Ascending Order

$$[1, 2, 3, 4]$$

- **Step 2:** Identify the Median

$$\text{Median} = \frac{2 + 3}{2} = \frac{5}{2} = 2.5$$

15.1.3. Mode

Value that appears most frequently

$$[1, 1, 2, 3]$$

- **Step 1:** Identify the Most Frequent Number

$$1 : 2$$

$$2 : 1$$

$$3 : 1$$

- **Step 2:** Determine the Mode

$$\text{Mode} = 1$$

15.2. Dispersion

15.2.1. Range

$$\text{Range} = \text{max} - \text{min}$$

[1, 2, 3, 4, 5]

- **Step 1:** Identify the Maximum and Minimum Values

$$\text{max} = 5$$

$$\text{min} = 1$$

- **Step 2:** Calculate the Range

$$\text{Range} = 5 - 1 = 4$$

15.2.2. Variance

Quantifies the spread or dispersion of a set of data points in a dataset

- Population

$$\sigma^2 = \frac{1}{N} \sum_{i=1}^n (x_i - \mu)^2$$

- Sample

$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

[70, 75, 80, 85, 90]

Step 1: Find mean

$$\bar{x} = \frac{70 + 75 + 80 + 85 + 90}{5} = \frac{400}{5} = 80$$

Step 2: Subtract the Mean and Square the result

$$(70 - 80)^2 = (-10)^2 = 100$$

$$(75 - 80)^2 = (-5)^2 = 25$$

$$(80 - 80)^2 = 0^2 = 0$$

$$(85 - 80)^2 = 5^2 = 25$$

$$(90 - 80)^2 = 10^2 = 100$$

Step 3: Calculate variance

$$s^2 = \frac{100 + 25 + 0 + 25 + 100}{5 - 1} = \frac{250}{4} = 62.5$$

15.2.3. Standard deviation

Quantifies the amount of variation or dispersion in a set of data points

- Population

$$\sigma = \sqrt{\sigma^2}$$

$$\sigma = \sqrt{\frac{1}{N} \sum_{i=1}^N (x_i - \mu)^2}$$

- Sample

$$s = \sqrt{s^2}$$

$$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^N (x_i - \bar{x})^2}$$

15.2.4. Interquartile Range (IQR)

$$\text{IQR} = Q3 - Q1$$

[1, 2, 3, 4, 5, 6, 7]

Step 1: Arrange the Data in Ascending Order

[1, 2, 3, 4, 5, 6, 7]

Step 2: Find the Quartiles

1. Calculate the Median (Q2)

$$\text{Median (Q2)} = 4$$

2. Find First Quartile (Q1)

Q1 is the median of the first half of the dataset

$$Q1 = 2$$

3. Find Third Quartile (Q3)

Q3 is the median of the second half of the dataset

$$Q3 = 5$$


Step 3: Calculate the Interquartile Range (IQR)

$$\text{IQR} = 5 - 2 = 3$$

16. Probability Distributions

16.1. Gaussian (Normal) distribution

$$f(x | \mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$




16.2. t-Distribution

$$f(t \mid \nu) = \frac{\Gamma(\frac{\nu+1}{2})}{\sqrt{\nu\pi}\Gamma(\frac{\nu}{2})} \left(1 + \frac{t^2}{\nu}\right)^{-\frac{\nu+1}{2}}$$

Where:

- t : t-statistic
- ν (or df): degrees of freedom
- Γ : Gamma function (generalizes the factorial function)



Continuous probability distribution for estimating the mean of a normally distributed population in situations where:


- Sample size is small
- Population standard deviation is unknown

Similar in shape to the normal distribution but has heavier tails, which means it gives more probability to values further from the mean

16.3. Binomial distribution

Discrete probability distribution that describes the number of successes in a fixed number of independent Bernoulli trials, each with the same probability of success


$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$



16.4. Poisson distribution


Used to model the number of events that occur within a fixed interval of time or space, given a constant mean rate and assuming that these events occur independently of each other

$$P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$$



16.5. Exponential distribution


$$f(x | \lambda) = \lambda e^{-\lambda x}$$



17. Functions

17.1. PDF (Probability Density Function)

Function that describes the likelihood of a continuous random variable taking on a particular value



Properties:

- The area under the curve of a PDF over the entire range of possible values equals 1
- The PDF itself is non-negative everywhere
- The probability that the variable falls within a certain range is given by the integral of the PDF over that range


17.2. PMF (Probability Mass Function)

Frequency distribution that provides the probability that a categorical variable takes on each of its possible values

17.3. CDF (Cumulative Distribution Function)

Gives the probability that X will take a value less than or equal to x

$$F(x) = P(X \leq x)$$



1. Categorical

$$F(x) = \int_{-\infty}^x f(t)dt$$

2. Continuous

$$F(x) = \sum_{t \leq x} P(X = t)$$

```
from scipy.stats import norm
```

```
x = 1  
mu = 0  
sigma = 1
```

```
norm.cdf(x, loc=mu, scale=sigma)
```

17.4. PPF (Percent-Point Function)

Gives the value x such that the probability of a random variable being less than or equal to x is equal to a given probability p

```
from scipy.stats import norm


p = 0.95
mu = 0
sigma = 1

x_value = norm.ppf(p, loc=mu, scale=sigma)
```

17.5. SF (Survival Function)

Probability that a certain event has not occurred by a certain time

$$S(t) = P(T > t)$$



Relationship to PDF:

$$S(t) = 1 - F(t)$$

```
from scipy.stats import norm

z = 3.4
mu = 0
sigma = 1

norm.sf(z, loc=mu, scale=sigma)
```

18. Error Metrics

18.1. MAE (Mean Absolute Error)

Average of squared differences between predicted (\hat{y}_i) and actual values (y_i)

$$\text{MAE} = \frac{1}{n} \sum_{i=1}^n |y_i - \hat{y}_i|$$

18.2. MSE (Mean Squared Error)

Average of squared differences between predicted (\hat{y}_i) and actual (y_i) values

$$\text{MSE} = \frac{1}{2} \sum_{i=1}^n (y_i - \hat{y}_i)$$

18.3. RMSE (Root Mean Squared Error)

square root of the average squared differences between predicted (\hat{y}_i) and actual (y_i) values

$$\text{RMSE} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)^2}$$

18.4. MAPE (Mean Absolute Percentage Error)

Average percentage difference between predicted (\hat{y}_i) and actual (y_i) values

$$\text{MAPE} = \frac{100}{n} \sum_{i=1}^n \left| \frac{y_i - \hat{y}_i}{y_i} \right|$$

18.5. R-squared

Proportion of variance explained by the model


$$R^2 = 1 - \frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y})^2}$$

- 1: model explains all the variance in the dependent variable
- 0: model explains none of the variance in the dependent variable

Step 1: Fit the Regression Model


Step 2: Compute **Total Sum of Squares** (SST)

$$\text{SST} = \sum_{i=1}^n (y_i - \bar{y})^2$$




Step 3: Compute **Regression Sum of Squares** (SSR)

$$\text{SSR} = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$$



Step 4: Compute Residual Sum of Squares (SSE)

$$SSE = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$



Step 5: Calculate R^2

$$R^2 = \frac{SSR}{SST} = 1 - \frac{SSE}{SST}$$

18.6. Adj R-squared

Adjusts the R^2 value based on the number of predictors (penalty for adding non-informative variables)

$$\text{Adj } R^2 = 1 - (1 - R^2) \frac{n - 1}{n - p - 1}$$

18.7. MSLE (Mean Squared Logarithmic Error)

When predictions and actual values span several orders of magnitude

$$\text{MSLE} = \frac{1}{n} \sum_{i=1}^n (\log(1 + y_i) - \log(1 + \hat{y}_i))^2$$

18.8. Cross-Entropy Loss (Log Loss)

Binary and multi-class classification

$$\text{Log Loss} = -\frac{1}{n} \sum_{i=1}^n [y_i \log(\hat{y}_i) + (1 - y_i) \log(1 - \hat{y}_i)]$$

19. Hypothesis Testing

19.1. Hypotheses

19.1.1. Null (H_0)

No effect or difference (observed effect is due to sampling variability)

$$H_0 : \mu = \mu_0$$

19.1.2. Alternative (H_1 or H_a)

Presence of an effect or a difference

- Two-Tailed

$$H_1 : \mu \neq \mu_0$$

- Right-Tailed

$$H_1 : \mu > \mu_0$$

- Left-Tailed

$$H_1 : \mu < \mu_0$$

19.2. Error Types

19.2.1. Type I (α)

$$\alpha = P(\text{Reject } H_0 \mid H_0 \text{ is true})$$

19.2.2. Type II (β)

$$\beta = P(\text{Fail to Reject } H_0 \mid H_1 \text{ is true})$$

19.3. t-Tests

19.3.1. One-sample

Tests if the mean of a single sample differs from a known or hypothesized population mean.

$$t = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}}$$

- \bar{x} : sample mean
- μ_0 : hypothesized population mean
- s : sample standard deviation
- n : sample size

[78, 82, 89]

Step 1: State Hypotheses

$$H_0 : \mu = 85 \text{ cm}$$

$$H_1 : \mu \neq 85 \text{ cm}$$

Step 2: Summarize Data

- Sample values:

$$x_1 = 78, x_2 = 82, x_3 = 89$$

- Sample size:

$$n = 3$$

Step 3: Calculate Sample Mean (\bar{x})

$$\bar{x} = \frac{x_1 + x_2 + x_3}{n} = \frac{78 + 82 + 89}{3} = 83$$

Step 4: Calculate Sample Standard Deviation:

$$s = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}}$$

- Find the deviations from the mean and square them

$$(x_1 - \bar{x})^2 = (78 - 83)^2 = (-5)^2 = 25$$

$$(x_2 - \bar{x})^2 = (82 - 83)^2 = (-1)^2 = 1$$

$$(x_3 - \bar{x})^2 = (89 - 83)^2 = (6)^2 = 36$$

- Sum of squared deviations

$$\text{SSD} = 25 + 1 + 36$$

- Calculate variance of sample

$$s^2 = \frac{\text{SSD}}{n - 1} = \frac{62}{3 - 1} = \frac{62}{2} = 31$$

- Calculate Sample Standard Deviation

$$s = \sqrt{s^2} = \sqrt{31} = 5.57$$

Step 5: Calculate the Test Statistic

$$t = \frac{\bar{x} - \mu_0}{\frac{s}{\sqrt{n}}} = \frac{83 - 85}{\frac{5.57}{\sqrt{3}}} = -\frac{2}{3.22} = -0.62$$

Step 6: Determine the Degrees of Freedom

$$df = n - 1 = 3 - 1 = 2$$

Step 7: Find Critical t-value

- For a two-tailed test at a significance level (α) of 0.05 and 2 degrees of freedom (df)

$$4.303$$

Step 8: Compare the t-Value to the Critical t-Value

- If the absolute value of the test statistic is greater than the critical t-value, reject the null hypothesis.
- If the absolute value of the test statistic is less than the critical t-value, fail to reject the null hypothesis.

Step 8: Find the p-Value

$$t = 4.303 \text{ and } df = 3$$

$$\text{p-value} = 0.58$$

```
from scipy import stats
rvs = stats.uniform.rvs(size=50)
stats.ttest_1samp(rvs, popmean=0.5)
```

19.3.2. Independent

Compares the means of two independent samples.

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\left(\frac{s_1^2}{n_1}\right) + \left(\frac{s_2^2}{n_2}\right)}}$$

- \bar{x}_1 and \bar{x}_2 : sample means
- s_1^2 and s_2^2 : sample variances
- n_1 and n_2 : sample sizes

```
from scipy import stats
rvs1 = stats.norm.rvs(loc=5, scale=10, size=500)
rvs2 = stats.norm.rvs(loc=5, scale=10, size=500)
stats.ttest_ind(rvs1, rvs2)
```

19.3.3. Paired

$$t = \frac{\bar{D}}{\frac{s_D}{\sqrt{n}}}$$

- \bar{D} : mean of the differences between paired observations
- s_D : standard deviation of the differences
- n : number of pairs

19.4. Chi-square tests

19.4.1. Goodness of Fit Test

Compares an **observed** categorical distribution to a **theoretical** categorical distribution.

$$\chi^2 = \sum_{i=1}^k \frac{(o_i - e_i)^2}{e_i}$$

- o_i : observed frequency in category i
- e_i : expected frequency in category i

```

from scipy import stats
f_obs = np.array([43, 52, 54, 40])
f_exp = np.array([47, 47, 47, 47])
stats.chisquare(f_obs=f_obs, f_exp=f_exp)

```

19.4.2. Test of independence

Compares two **observed** categorical distributions.

$$X^2 = \sum_{i=1}^k \frac{(o_{ij} - e_{ij})^2}{e_{ij}}$$

- o_{ij} : observed frequency in cell (i, j)
- e_{ij} : expected frequency in category (i, j) calculated as:

$$E_{ij} = \frac{R_i \times C_j}{N}$$

- R_i : Row total for row i
- C_j : Column total for column j
- N : Total number of observations

19.5. ANOVA (Analysis of Variance)

19.5.1. One-way

Compares the means of three or more groups based on one independent variable

- $H_0: \mu_1 = \mu_2 = \dots = \mu_k$
- H_1 : At least one μ_i differs from the others

Step 1: Calculate Between-Group Variation (SS_{between})

$$SS_{\text{between}} = \sum_{i=1}^k n_i (\bar{X}_i - \bar{X}_{\text{overall}})^2$$

- n_i : Number of observations in group i
- \bar{X}_i : Mean of group i
- \bar{X}_{overall} : Overall mean of all groups

Step 2: Calculate Within-Group Variation (SS_{within})

$$SS_{\text{within}} = \sum_{i=1}^k \sum_{j=1}^{n_i} (X_{ij} - \bar{X}_i)^2$$

- X_{ij} : Observation j in group i

Step 3: Calculate Total Variation (SS_{total})

$$SS_{\text{total}} = SS_{\text{between}} + SS_{\text{within}}$$

Step 4: Calculate Mean Squares

- Mean Square Between (MS_{between})

$$MS_{\text{between}} = \frac{SS_{\text{between}}}{k - 1}$$

- Mean Square Within (MS_{within})

$$MS_{\text{within}} = \frac{SS_{\text{within}}}{N - k}$$

- N : total number of observations
- k : number of groups

Step 5: Calculate the F-statistic

$$F = \frac{MS_{\text{between}}}{MS_{\text{within}}}$$

Step 6: Decision Rule

- Compare the F-statistic to the critical value from the F-distribution table (based on chosen significance level α).
- Alternatively, compare the p-value to the significance level α .
- Reject H_0 if the F-statistic is greater than the critical value or if the p-value is less than α , indicating that at least one group mean is significantly different.

19.5.2. Two-way

20. Regression Analysis:

20.1. Simple linear regression

$$y = \beta_0 + \beta_1 x_1 + \varepsilon$$

- y : dependent variable
- x : independent variables
- β_0 : intercept (value of y when $x = 0$)
- β_1 : slope (change in y for a one-unit change in x)
- ε : error term (difference between the actual data points and the predicted values)

Data

x (Hours Studied)	y (Test Score)
1	2
2	3
3	5
4	4
5	6

Step 1: Calculate Means

$$\bar{x} = 3$$

$$\bar{y} = 4$$

Step 2: Calculating Slope β_1

$$\beta_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

- The numerator

$$\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = 9$$

- The denominator

$$\sum_{i=1}^n (x_i - \bar{x})^2 = 10$$

- The slope β_1 is

$$\beta_1 = \frac{9}{10} = 0.9$$

Step 3: Calculate Intercept β_0

$$\beta_0 = \bar{y} - \beta_1 \bar{x} = 1.3$$

Step 4: Calculate p-Value

- Calculate Standard Error of the Slope (SE_{β_1})

$$SE_{\beta_1} = \sqrt{\frac{\sum_{i=1}^n (y_i - \hat{y}_i)^2}{(n-2) \cdot \sum_{i=1}^n (x_i - \bar{x})^2}}$$

- Calculate the Residual Sum of Squares (RSS)

$$RSS = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

- Calculate the t-statistic for the Slope

$$t = \frac{\beta_1}{SE_{\beta_1}}$$

- Determine Degrees of Freedom

$$df = n - 2$$

- Look up the p-value corresponding to t with 3 degrees of freedom

Step 5: Calculate R^2 (R_{adj}^2)

$$R^2 = \frac{SS_{reg}}{SS_{total}}$$

SS_{reg} (Regression Sum of Squares): sum of the squared differences between the predicted \hat{y} values and the mean of the observed y values.

$$SS_{\text{reg}} = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$$

SS_{total} (Total Sum of Squares): sum of the squared differences between the observed y values and the mean of the observed y values.

$$SS_{\text{total}} = \sum_{i=1}^n (y_i - \bar{y})^2$$

Adjusting for the number of independent variables

$$R_{\text{adj}}^2 = 1 - \left(\frac{(1 - R^2)(n - 1)}{n - k - 1} \right)$$

- n : number of observations
- k : number of independent variables

20.2. Multiple regression

$$Y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_n x_n + \varepsilon$$

Matrix form:

$$Y = X\beta + \varepsilon$$

Estimating Coefficients (OLS):

$$\hat{\beta} = (X^T X)^{-1} X^T Y$$

Where:

- $\hat{\beta}$: vector of estimated coefficients

20.3. Logistic Regression

Binary classification (1 or 0, true or false, yes or no) based on one or more predictor variables

Sigmoid function:

$$\sigma(z) = \frac{1}{1 + e^{-z}}$$

Where:

$$z = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_n x_n$$


20.4. Model diagnostics

20.4.1. p-Values

Probability of obtaining results at least as extreme as the observed results


1. One-Tailed

$$p = P(Z > z_{\text{observed}})$$



2. Two-Tailed

$$p = 2 \cdot P(Z > |z_{\text{observed}}|)$$



```
z = 2.1
df = 3
scipy.stats.norm.sf(abs(z), df=df)
```

20.4.2. F-Statistic

$$F = \frac{\text{MSR}}{\text{MSE}}$$

Where:

- Mean Square Regression (MSR):

$$\text{MSR} = \frac{\text{SSR}}{\text{df}_{\text{regression}}}$$

- $\text{df}_{\text{regression}} = p$

Where:

- p : Number of independent
- Mean Square Error (MSE):

$$\text{MSE} = \frac{\text{SSE}}{\text{df}_{\text{error}}}$$

- $\text{df}_{\text{error}} = n - p - 1$

Where:

- p : Number of independent
- n : Number of observations
- 1: Constant (for intercept)

20.4.3. Confidence Intervals (CI)

Range within which we can be confident that the true value (population parameter) lies, based on the sample data

1. Known population standard deviation (σ):

$$CI = \bar{x} \pm z \frac{\sigma}{\sqrt{n}}$$

Where:

- \bar{x} : sample mean
- z : z-score corresponding to the desired confidence level
- σ : population standard deviation
- n : sample size

2. Unknown population standard deviation (σ):

$$CI = \bar{x} \pm t \frac{s}{\sqrt{n}}$$

Where:

- \bar{x} : sample mean
- t : critical value from t-distribution
- s : sample standard deviation
- n : sample size

21. Correlation

21.1. Pearson

Linear relationships

$$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \times (y_i - \bar{y})^2}}$$

Interpretation:

- 1: perfect positive linear relationship
- -1: perfect negative linear relationship
- 0: no linear relationship

$$X : [1, 2, 3, 4, 5]$$

$$Y : [2, 4, 6, 8, 10]$$

Step 1: Calculate the Means of X and Y

$$\bar{X} = 3$$

$$\bar{Y} = 6$$

Step 2: Calculate the Differences from the Mean

21.2. Spearman's Rank

Non-linear relationships

$$\rho = 1 - \frac{6 \sum_{i=1}^n d_i^2}{n(n^2 - 1)}$$

Where

- ρ : Spearman rank correlation coefficient
- d_i : difference between the ranks of corresponding values
- n : number of observations

Rank: position of a value within a data set when the values are ordered in ascending order

Observation i	X	Rank X	Y	Rank Y	$d_i = \text{Rank } X - \text{Rank } Y$	d_i^2
1	3	3	8	5	-2	4
2	1	1	6	3	-2	4
3	4	4	7	4	0	0
4	2	2	4	1	1	1
5	5	5	5	2	3	9

Interpretation

- $\rho = 1$: Perfect positive correlation
- $\rho = -1$: Perfect negative correlation
- $\rho = 0$: No correlation

22. Non-Parametric Statistics

22.1. Mann-Whitney U

Determine whether there is a significant difference between the distributions of two independent samples (used as an alternative to the independent samples t-test when the assumptions of normality are not met)

22.2. Wilcoxon Signed-Rank

Compare two paired samples or to assess whether the median of a single sample is different from a specified value (used as a non-parametric alternative to the paired t-test when the data does not meet the assumptions of normality)

22.3. Kolmogorov-Smirnov

Determine if a sample is drawn from a population with a specific distribution.

It compares the empirical distribution function (EDF) of the sample with the cumulative distribution function (CDF) of the reference distribution.

22.4. Kruskal-Wallis

Determine if there are significant differences between the medians of three or more independent groups (extension of the Mann-Whitney U test, which is used for comparing two groups)


23. Time Series

23.1. SMA (Simple Moving Averages)

For a time series y_t , the SMA of window size k at time t is:

$$s_t = \frac{1}{k} \sum_{i=0}^{k-1} y_{t-i}$$

This is the average of the last k values.



The $k = 3$ day SMA

Data

t	x (\$)
1	10
2	12
3	14
4	16
5	18

$$s_3 = \frac{x_{t_1} + x_{t_2} + x_{t_3}}{k} = \frac{10 + 12 + 14}{3} = 12$$

$$s_4 = \frac{x_{t_2} + x_{t_3} + x_{t_4}}{k} = \frac{12 + 14 + 16}{3} = 14$$

$$s_5 = \frac{x_{t_3} + x_{t_4} + x_{t_5}}{k} = \frac{14 + 16 + 18}{3} = 16$$

```
pl.col('X').rolling_mean(window_size)
```

23.2. WMA (Weighted Moving Average)

Each data point in the window is assigned a specific weight (usually decrease linearly)

$$\text{WMA} = \frac{\sum_{i=1}^n (x_i w_i)}{\sum_{i=1}^n w_i}$$

Data

- Day 1: \$10
- Day 2: \$12
- Day 3: \$14
- Day 4: \$13
- Day 5: \$15

Weights

- 1st most recent: 3
- 2nd most recent: 2
- 3rd most recent: 1

$$\text{WMA for Day 3} = \frac{(14 \times 3) + (12 \times 2) + (10 \times 1)}{3 + 2 + 1} = 12.67$$

$$\text{WMA for Day 4} = \frac{(13 \times 3) + (14 \times 2) + (12 \times 1)}{3 + 2 + 1} = 13.17$$

$$\text{WMA for Day 5} = \frac{(15 \times 3) + (13 \times 2) + (14 \times 1)}{3 + 2 + 1} = 14.7$$

23.3. Exponential Smoothing

Gives more weight to recent data points (more responsive to new information)


$$s_t = \alpha x_t + (1 - \alpha)s_{t-1}$$

Where:

- s_t : the smoothed value at time t
- x_t : actual value at time t
- α : smoothing factor between 0 and 1

Common way to choose α based on the smoothing window length N :

$$\alpha = \frac{2}{N + 1}$$



Data

t	x (\$)
1	10
2	12
3	14
4	13
5	15

Calculate Smoothing Factor

$$\alpha = \frac{2}{3 + 1} = 0.5$$

Calculate SMA for First Value

$$s_3 = \frac{10 + 12 + 14}{3}$$

Calculate EMA

$$s_4 = (0.5 \times 13) + (12 \times 0.5) = 12.5$$

$$s_5 = (0.5 \times 15) + (12.5 \times 0.5) = 13.75$$

```
pl.col("X").ewm_mean(span=window_size, adjust=False)
```

23.4. Seasonal Decomposition

- Level: baseline value around which the time series fluctuates
- Trend: ong-term progression or direction of the time series
- Seasonality: regular, repeating patterns or cycles in the time series

- Noise: random fluctuations or irregular variations (cannot be explained by the trend, seasonality, or other components)

Additive Decomposition

When the magnitude of seasonal fluctuations and trend does not change over time

$$Y_t = L_t + T_t + S_t + N_t$$

Where

- L_t : Level at time t
- T_t : Trend at time t
- S_t : Seasonal component at time t
- N_t : Noise (residuals) at time t

Multiplicative Decomposition

When the magnitude of seasonal fluctuations and trend change proportionally over time

$$Y_t = L_t \times T_t \times S_t \times N_t$$

Where

- L_t : Level at time t
- T_t : Trend at time t
- S_t : Seasonal component at time t
- N_t : Noise (residuals) at time t

23.5. ARMA (AutoRegressive Moving Average)

23.6. ARIMA (AutoRegressive Integrated Moving Average)

Industrial Engineering & Operations Research

24. Causes of variation

24.1. Common

Common Cause Variation: Normal, expected variation inherent to the process; predictable and stable.

- Temperature
- Humidity
- Material properties

24.2. Special

Special Cause Variation: Abnormal, unexpected variation due to specific causes; unpredictable and requires immediate action.

External disruptions

- Machine malfunction
- Faulty raw materials
- Staff shortages
- Weather events

25. Design of Experiments (DOE):

26. Control Charts:

26.1. P-charts (Proportion)

The p-chart monitors the proportion of defectives in each sample, accounting for variable sample sizes.

$$UCL_{p_i} = \bar{p} + 3\sqrt{\frac{\bar{p}(1 - \bar{p})}{n_i}}$$
$$LCL_{p_i} = \bar{p} - 3\sqrt{\frac{\bar{p}(1 - \bar{p})}{n_i}}$$

Where:


- d_i : The number of defective items observed in sample i
- p_i : sample proportion defective

$$p_i = \frac{d_i}{n_i}$$

- \bar{p} : overall proportion defective

$$\bar{p} = \frac{\sum d_i}{\sum n_i}$$

- n_i : sample size for sample i



26.2. NP-charts (Number Proportion)

Number of defective items (constant sample size)

$$UCL_{np} = n\bar{p} + 3\sqrt{n\bar{p}(1-\bar{p})}$$


$$LCL_{np} = n\bar{p} - 3\sqrt{n\bar{p}(1-\bar{p})}$$

Where:

- n : count of defective items in each sample
- p : proportion of defectives
- \bar{p} :

$$\begin{aligned}\bar{p} &= \frac{1}{k} \sum_{i=1}^k \frac{d_i}{n} \\ &= \frac{\sum_{i=1}^k d_i}{n \cdot k}\end{aligned}$$

- $n\bar{p} = \bar{np}$



26.3. C-charts (Count)

Count of defects (fixed unit size)

Number of defects observed in each sample or inspection unit


$$\bar{c} = \frac{1}{k} \sum_{i=1}^k c_i$$

$$UCL_c = \bar{c} + 3\sqrt{\bar{c}}$$

$$LCL_c = \bar{c} - 3\sqrt{\bar{c}}$$

Where:

- c : number of defects
- k : number of samples



This C-chart displays the number of defects identified in each of 25 inspected units, where each unit is of fixed size. The center line represents the average number of defects across all samples, calculated as:

$$\bar{c} = \frac{1}{k} \sum_{i=1}^k c_i$$

Control limits are drawn at ± 3 standard deviations from the mean, using the square root of \bar{c} to account for Poisson-distributed defect counts:

$$UCL_c = \bar{c} + 3\sqrt{\bar{c}} \quad LCL_c = \max(0, \bar{c} - 3\sqrt{\bar{c}})$$

26.4. U-charts (Unit)

Defects per unit (variable unit size)

$$UCL_u = \bar{u} + 3\sqrt{\frac{\bar{u}}{n_i}}$$

$$LCL_u = \bar{u} - 3\sqrt{\frac{\bar{u}}{n_i}}$$


Where:

- c_i : number of defects in sample i
- n_i : sample size of group i
- u_i : defects per unit

$$u_i = \frac{c_i}{n_i}$$

- \bar{u} :

$$\frac{\sum c_i}{\sum n_i}$$



26.5. \bar{X} -chart


Used to monitor the average of a continuous quality characteristic (e.g., weight, length, temperature), assuming constant sample size per subgroup.

$$UCL_{\bar{x}} = \bar{\bar{x}} + A_2 \cdot \bar{R}$$

$$LCL_{\bar{x}} = \bar{\bar{x}} - A_2 \cdot \bar{R}$$

Where:

- \bar{x}_i : average of subgroup i
- $\bar{\bar{x}}$: grand mean of all subgroup means
- \bar{R} : average range of all subgroups
- A_2 : constant dependent on sample size n (e.g., $A_2 = 0.577$ for $n = 5$)



26.6. R-chart


27. Process Capability Analysis

27.1. C_p (Process Capability Index)

Measure how well a process can produce outputs within specified limits

$$C_p = \frac{USL - LSL}{6\sigma}$$

Assumption: Process is **centered** within the specification limits



C_p Value	Meaning	Quality Impact
$C_p < 1$	Process variation exceeds specs	Defects likely, even if centered
$C_p = 1$	Process barely fits within specs	OK if perfectly centered
$C_p > 1$	Process well within specs	Robust, fewer defects
$C_p = 2$	Six Sigma level	World-class performance

A manufacturing process produces metal rods with a target diameter of 10.0 mm. The specification limits are:


- Target: 10 mm
- Lower Specification Limit (LSL): 9.7 mm
- Upper Specification Limit (USL): 10.3 mm
- Specification Width (Tolerance): 0.6 mm

Let’s analyze three different processes with varying levels of capability.

Process	Mean	Std Dev	C_p	Assessment
Process 1	9.998	0.099	1.009	Good
Process 2	10.001	0.147	0.682	Marginal
Process 3	9.994	0.255	0.393	Poor

Process 1	Process 2	Process 3	Six Sigma
-----------	-----------	-----------	-----------

$C_p = \frac{USL - LSL}{6\sigma}$	$C_p = \frac{USL - LSL}{6\sigma}$	$C_p = \frac{USL - LSL}{6\sigma}$	$C_p = \frac{USL - LSL}{6\sigma}$
$= \frac{10.3 - 9.7}{6 \times 0.099}$	$= \frac{10.3 - 9.7}{6 \times 0.147}$	$= \frac{10.3 - 9.7}{6 \times 0.255}$	$= \frac{10.3 - 9.7}{6 \times 0.05}$
$= \frac{0.6}{0.6}$	$= \frac{0.6}{0.9}$	$= \frac{0.6}{1.5}$	$= \frac{0.6}{0.30}$
$= 1.009$	$= 0.682$	$= 0.393$	$= 2$



Step 3: Interpretation

Process	C_p	Within Spec	Defective	DPMO
Process 1	1.009	96.64%	3.36%	33600
Process 2	0.682	84.13%	15.87%	159700
Process 3	0.393	50.00%	50.00%	500000
Six Sigma	2.0	99.9999998%	0.0000002%	0.002

Higher C_p means fewer defects and better process quality. and lower defects per million opportunities (DPMO)

```
cp_values = [2, 1, 0.1]

for cp in cp_values:
    z = 3 * cp # spec limit in standard deviations
    prob_defective = 2 * (1 - norm.cdf(z))
    within_specs = 1 - prob_defective
    dpmo = prob_defective * 1_000_000

    print({
        "Cp": cp,
        "Sigma limit (±z)": z,
        "Percent within specs": within_specs * 100,
        "Percent defective": prob_defective * 100,
        "DPMO": dpmo
    })
```

Output:

```
[{'Cp': 2,
  'Sigma limit (±z)': 6,
  'Percent within specs': 99.99999980268245,
  'Percent defective': 1.9731754008489588e-07,
  'DPMO': 0.001973175400848959},
 {'Cp': 1,
  'Sigma limit (±z)': 3,
  'Percent within specs': 99.73002039367398,
  'Percent defective': 0.2699796063260207,
  'DPMO': 2699.796063260207},
 {'Cp': 0.1,
  'Sigma limit (±z)': 0.30000000000000004,
  'Percent within specs': 23.582284437790534,
  'Percent defective': 76.41771556220947,
  'DPMO': 764177.1556220946}]
```

27.2. C_{pk} (Process Capability Index with Centering)

Minimum distance from mean to nearest spec limit divided by 3 standard deviations. Process capability considering centering and variation.

If process is centered:


$$C_{pk} = C_p = \frac{USL - LSL}{6\sigma}$$

If process mean is off center, the smaller side of the process-to-limit ratio determines C_{pk} :

$$C_{pk} = \text{Min} \left(\frac{USL - \bar{x}}{3\sigma}, \frac{\bar{x} - LSL}{3\sigma} \right)$$

If $C_{pk} < C_p$, process is not centered

$C_{\{pk\}}$ Value	Meaning	Quality Impact
$C_{\{pk\}} < 1$	Process not capable, output falls outside specs	Many defects, poor performance
$C_{\{pk\}} = 1$	Process just meets specification limits	Acceptable only if well centered
$C_{\{pk\}} > 1$	Process is capable, centered or nearly so	Fewer defects, reliable quality
$C_{\{pk\}} = 2$	Excellent capability, Six Sigma level	Near-perfect quality



Suppose a company manufactures metal rods, and the specification limits for the diameter of the rods are:

- Target: 10 mm
- Lower Specification Limit (**LSL**): 9.8 mm
- Upper Specification Limit (**USL**): 10.2 mm

The process has:

- A standard deviation σ of 0.05 mm.
- A process mean μ of 10.1 mm.

Step 1: Calculate the distance from the mean to the USL:

$$\frac{USL - \mu}{3\sigma} = \frac{10.2 \text{ mm} - 10.1 \text{ mm}}{3 \times 0.05 \text{ mm}} = \frac{0.1 \text{ mm}}{0.15 \text{ mm}} = 0.67$$

Step 2: Calculate the distance from the mean to the LSL:

$$\frac{\mu - LSL}{3\sigma} = \frac{10.1 \text{ mm} - 9.8 \text{ mm}}{3 \times 0.05 \text{ mm}} = \frac{0.3 \text{ mm}}{0.15 \text{ mm}} = 2.00$$

Step 3: Determine C_{pk} :

$$C_{pk} = \min(0.67, 2.00) = 0.67$$

Interpretation:


$C_p = 1.33$ means the process spread (6σ) fits 1.33 times within the tolerance range (the distance between the Upper Specification Limit and Lower Specification Limit).

- $C_p = 1.00$: Process mean is exactly at the midpoint of the specification limits, and the process variation fits exactly within these limits. 99.73% of the output will be within specifications, indicating a capable process (3 sigma process).
- $C_p > 1.00$: The higher the C_{pk} , the more capable and stable the process is, meaning it can consistently produce parts within tolerance with minimal risk of defects.
- $C_p < 1$: The process mean is off-center or the variation is wider than the specification limits, or both. A significant portion of the output may fall outside the specification limits.

27.3. C_{pm} (Taguchi Capability Index)

27.4. P_p (Process Performance Index)

$$P_p = \frac{USL - LSL}{6\sigma_{\text{overall}}}$$



Suppose a company manufactures metal rods, and the specification limits for the diameter of the rods are:

- Upper Specification Limit (**USL**): 10.2 mm
- Lower Specification Limit (**LSL**): 9.8 mm
- Overall standard deviation (σ_{overall}): 0.06 mm

Step 1: Determine the Specification Width

The specification width is the difference between the USL and LSL.

$$\text{Specification Width} = \text{USL} - \text{LSL} = 10.2 \text{ mm} - 9.8 \text{ mm} = 0.4$$

Step 2: Calculate the Process Performance Index P_p

The formula for P_p is:

$$P_p = \frac{\text{Specification Width}}{6\sigma_{\text{overall}}} = \frac{\text{USL} - \text{LSL}}{6\sigma_{\text{overall}}}$$

Substitute the values:

$$P_p = \frac{0.4 \text{ mm}}{6 \times 0.06 \text{ mm}} = \frac{0.4 \text{ mm}}{0.36 \text{ mm}} = 1.11$$

Interpretation:


- A P_p of 1.11 indicates that the process performance, considering all sources of variation over time, is capable but less so than the potential capability indicated by C_p . The value being slightly above 1 suggests that the process can generally produce rods within specifications, but there might be more variability in the process compared to the short-term capability measured by C_p .
- Decrease from the C_p value (1.33 to 1.11) reflects the impact of additional variability when evaluating the process over a longer time or under different conditions.

Use:

When you need to evaluate the **actual performance** of a process over a longer period, considering all sources of variation, including shifts, drifts, and other long-term factors.

27.5. P_{pk} (Process Performance Index with Centering)

$$P_{pk} = \text{Min} \left(\frac{\text{USL} - \mu_{\text{overall}}}{3\sigma}, \frac{\mu_{\text{overall}} - \text{LSL}}{3\sigma} \right)$$



Suppose a company manufactures metal rods, and the specification limits for the diameter of the rods are:

- Upper Specification Limit (**USL**): 10.2 mm
- Lower Specification Limit (**LSL**): 9.8 mm
- Overall standard deviation (σ_{overall}): 0.06 mm
- Overall process mean (μ_{overall}): 10.1 mm

Step 1: Calculate the Distance from the Process Mean to the Specification Limits

Calculate the distance from the overall process mean to both the USL and LSL:

$$\text{USL} - \mu_{\text{overall}} = 10.2 \text{ mm} - 10.1 \text{ mm} = 0.1 \text{ mm}$$

$$\mu_{\text{overall}} - \text{LSL} = 10.1 \text{ mm} - 9.8 \text{ mm} = 0.3 \text{ mm}$$

Step 2: Calculate the Process Performance Index P_k

The formula for P_{pk} is:

$$P_{pk} = \text{Min} \left(\frac{\text{USL} - \mu_{\text{overall}}}{3\sigma}, \frac{\mu_{\text{overall}} - \text{LSL}}{3\sigma} \right)$$

Substitute the values:

$$P_{pk} = \text{Min} \left(\frac{0.1 \text{ mm}}{3 \times 0.06 \text{ mm}}, \frac{0.3 \text{ mm}}{3 \times 0.06 \text{ mm}} \right)$$

$$P_{pk} = \text{Min} \left(\frac{0.1 \text{ mm}}{0.18 \text{ mm}}, \frac{0.3 \text{ mm}}{0.18 \text{ mm}} \right)$$

$$P_{pk} = \min(0.56, 1.67) = 0.56$$

Interpretation:

- A P_p of 1.11 indicates that the process performance, considering all sources of variation over time, is capable but less so than the potential capability indicated by C_p . The value being slightly above 1 suggests that the process can generally produce rods within

specifications, but there might be more variability in the process compared to the short-term capability measured by C_p

- Decrease from the C_p value (1.33 to 1.11) reflects the impact of additional variability when evaluating the process over a longer time or under different conditions.

Use:

When you need to evaluate the **actual performance** of a process over a longer period, considering all sources of variation, including shifts, drifts, and other long-term factors.

28. Inventory Management

28.1. Newsvendor

determine the optimal order quantity Q^* that minimizes the total expected cost or maximizes the expected profit, based on the trade-off between the overage and underage costs

Assumptions

- Products are separable
- Planning is done for a single period
- Demand is random
- Deliveries are made in advance of demand
- Costs of overage or underage are linear

1. Parameters

- P : Sale Price
- C : Purchase Cost
- S : Unsold Value
- μ : Mean Demand
- σ : Standard Deviation Demand

2. Calculate Underage and Overage Costs:

- Underage Cost (C_u): Profit lost for each unit of demand not met

$$C_u = P - C$$

- Overage Cost (C_o): This is the cost of holding an unsold newspaper.

$$C_o = C - S$$

3. Calculate Critical Ratio (CR)

$$CR = \frac{C_u}{C_u + C_o}$$

4. Find z-score: Find the number of standard deviations away from the mean corresponding to the critical ratio:

$$z^* = \Phi^{-1}(CR)$$

Where:

- Φ^{-1} : Inverse of the CDF of the standard normal distribution (PPF)

5. Calculate Optimal Order Quantity (Q^*)

$$Q^* = \mu + z^* \sigma$$

Where:

- z^* : z-score corresponding to the critical ratio CR from the standard normal distribution

Consider a newsvendor selling newspapers:

1. Parameters

- Sale Price (P): \$3 (per unit)
- Purchase Cost (C): \$1 (per unit)
- Unsold Value (S): \$0 (per unit)
- Mean Demand (μ): 100 (units)
- Standard Deviation Demand (σ): 20 (units)

2. Calculate Underage and Overage Costs:

- Underage Cost (C_u): Profit lost for each unit of demand not met

$$C_u = P - C = 3 - 1 = 2$$

- Overage Cost (C_o): This is the cost of holding an unsold newspaper.

$$C_o = C - S = 1 - 0 = 1$$

3. Calculate Critical Ratio (CR)

$$CR = \frac{C_u}{C_u + C_o} = \frac{2}{2 + 1} = \frac{2}{3} = 0.67$$

4. Find z-score: Find the number of standard deviations away from the mean corresponding to the critical ratio:

$$z^* = \Phi^{-1}(CR) = 0.44$$

5. Calculate Optimal Order Quantity (Q^*)

$$Q^* = \mu + z^* \sigma = 100 + 0.44 \cdot 20 = 108.8$$

```
import numpy as np
import scipy.stats as stats
import matplotlib.pyplot as plt

# Parameters for the newsvendor example
selling_price = 3 # Selling price per newspaper
purchase_cost = 1 # Purchase cost per newspaper
unsold_value = 0 # Value of unsold newspapers
mu = 100 # Mean of demand
sigma = 20 # Standard deviation of demand

# Calculate underage and overage costs
C_u = selling_price - purchase_cost # Underage cost
C_o = purchase_cost - unsold_value # Overage cost

# Calculate the critical ratio
CR = C_u / (C_u + C_o)
```



```
# Assume a normal distribution for demand
mu = 50 # Mean demand
sigma = 10 # Standard deviation of demand

# Calculate the optimal order quantity (Q*)
z_star = stats.norm.ppf(CR) # z-score corresponding to the critical ratio
Q_star = mu + z_star * sigma # Optimal order quantity
```

28.2. ABC Analysis

categorizes inventory items into three groups (A, B, and C) based on their importance

- A: Top 70-80% of the total annual consumption value
- B: Next 15-25% of the total annual consumption value
- C: Remaining 5-10% of the total annual consumption value

Item	Usage Quantity (Annual)	Unit Cost	Consumption Value (Annual)
I_1	50	\$100	\$5000
I_2	150	\$20	\$3000
I_3	300	\$10	\$3000
I_4	400	\$5	\$2000
I_5	500	\$1	\$500

Step 1: Calculate Annual Consumption Values

Step 2: Sort Items by Annual Consumption Value (Descending)

Step 3: Calculate Total Annual Consumption Value

$$\text{Total} = \$5,000(I_1) + \$3,000(I_2) + \$3,000(I_3) + \$2,000(I_4) + \$500(I_5) = \$13500$$

Step 4: Calculate Cumulative Consumption Value Percentages

- $I_1 : \frac{5000}{13500} \times 100\% = 37.04\%$
- $I_2 : \frac{3000}{13500} \times 100\% = 22.22\%$
- $I_3 : \frac{3000}{13500} \times 100\% = 22.22\%$
- $I_4 : \frac{2000}{13500} \times 100\% = 14.81\%$
- $I_5 : \frac{500}{13500} \times 100\% = 3.7\%$

Step 5: Cumulative Percentages

- I_1 : 37.04%
- $I_1 + I_2$: 59.26%
- $I_1 + I_2 + I_3$: 81.48%

- $I_1 + I_2 + I_3 + I_4$: 96.3%
- $I_1 + I_2 + I_3 + I_4 + I_5$: 100%

Step 6: Categorize Items

- A: I_1, I_2, I_3
- B: I_4
- C: I_5

28.3. Fill Rate

Percentage of customer demand that is satisfied from available inventory

$$F = \frac{U_f}{U_o} \times 100\%$$

Where:

- F : Fill Rate
- U_f : Number of units fulfilled
- U_o : Total number of units ordered

Customers order 100 units of a product and 90 units are fulfilled from stock:

$$F = \frac{90}{100} \times 100\% = 90\%$$

28.4. (OCT) Order Cycle Time

Measures the total time taken from when a customer places an order to when the order is delivered

$$\text{OCT} = T_{\text{order}} + T_{\text{processing}} + T_{\text{production}} + T_{\text{shipping}}$$

Where:

- T_{order} : Order Entry Time (time it takes to receive and log the order)
- $T_{\text{processing}}$: Order Processing Time (time to check inventory, verify details, and prepare for production or shipment)
- $T_{\text{production}}$: Production Time (time to manufacture or prepare the product)
- T_{shipping} : Shipping Time (time it takes to deliver the product from the warehouse to the customer)

28.5. ROP (Reorder Point)

The inventory level at which a new order should be placed to avoid stockouts

$$\text{ROP} = (\text{Average Demand per Period} \times \text{Lead Time})$$

Suppose your business sells 50 units per week, and the lead time for a new order is 2 weeks.
Using the formula:

$$\text{ROP} = 50(\text{units per week}) \times 2(\text{weeks}) = 100 \text{ units}$$

This means that when your inventory level drops to 100 units, you should place a new order to avoid running out of stock.

28.6. XYZ Analysis

Categorization based on variability

- X: Low variability
- Y: Moderate variability
- Z: High variability

Coefficient of Variation:

$$CV = \frac{\sigma}{\mu}$$

- X Items: Low CV
 - $CV < k_1$, where k_1 is the threshold value indicating low variability
- Y Items: Moderate CV
 - $CV < k_2$, where k_2 is the moderate value indicating low variability
- Z Items: High CV
 - $CV < k_3$, where k_3 is the threshold value indicating high variability

Step 1: Collect Historical Data (12 months)

- Product A: [100, 105, 98, 102, 101, 104, 103, 100, 99, 100, 101, 102]
- Product B: [150, 155, 145, 160, 140, 150, 155, 150, 165, 155, 150, 140]
- Product C: [200, 180, 220, 190, 210, 240, 180, 230, 220, 210, 250, 190]

Step 2: Calculate the Mean and Standard Deviation

- Product A:
 - $\mu = 101$
 - $\sigma = 2$
- Product B:
 - $\mu = 150$
 - $\sigma = 8$
- Product C:
 - $\mu = 210$
 - $\sigma = 25$

Step 3: Calculate the Coefficient of Variation (CV)

- $CV = \frac{2}{101} = 0.0198$
- $CV = \frac{8}{150} = 0.0533$
- $CV = \frac{25}{210} = 0.1190$

Step 4: Categorization

- X: Product A (Low variability)
- Y: Product B (Moderate variability)

- Z: Product C (High variability)

28.7. EOQ (Economic Order Quantity)

Optimal order quantity that **minimizes the total cost**, which includes both **holding costs** and **ordering costs**

$$EOQ = \sqrt{\frac{2DS}{H}}$$

Where:

- D : Demand per time period
- S : Ordering cost per order
- H : Holding cost per unit, per time period

A company sells widgets and wants to determine the optimal order quantity for inventory.

- The annual demand for widgets (D) is 12,000 units.
- The cost to place an order (S) is \$50.

The holding cost per unit per year (H) is \$2.

$$EOQ = \sqrt{\frac{2 \times 12000 \times 50}{2}} = 775$$

The company should order 775 widgets each time they place an order to minimize the total cost, which includes both ordering and holding costs

28.7.1. Perfect Order Rate

Measures the percentage of orders delivered to customers in full, on time, and without any damage

$$\text{Perfect Order Rate} = \frac{\text{Number of Perfect Orders}}{\text{Total Number of Orders}} \times 100\%$$

Where:

- **Number of Perfect Orders:** The number of orders that are delivered on time, complete, and undamaged.
- **Total Number of Orders:** The total number of orders fulfilled within a specific period.

Suppose you received 1,000 orders over a quarter, and 900 of those orders were delivered on time, complete, and without damage.

$$\text{Perfect Order Rate} = \frac{900}{1000} \times 100\% = 90\%$$

28.8. Safety Stock

Additional quantity of inventory kept on hand to protect against uncertainties in demand or supply. buffer to prevent stockouts due to unexpected variations in demand or delays in supply.

1. Constant Demand & Constant Lead Time

$$SS = Z \times \sigma_D$$

Where:

- Z : Z-score corresponding to the desired service level
- σ : standard deviation of demand

2. Variable Demand & Constant Lead Time

$$SS = Z \times \sigma_D \times \sqrt{L}$$

Where:

- Z : Z-score corresponding to the desired service level
- σ_D : Standard deviation of demand per unit of time
- L : Lead time

3. Variable Lead Time & Constant Demand

$$SS = Z \times \bar{D} \times \sigma_L$$

Where:

- Z : Z-score corresponding to the desired service level
- \bar{D} : Average demand
- σ_L : Standard deviation of lead time

4. Variable Demand & Variable Lead Time

$$SS = Z \times \sqrt{(\bar{D}^2 \times \sigma_L^2) + (L \times \sigma_D^2)}$$


Where:

- \bar{L} : Average Lead Time (average time it takes to receive inventory after placing an order)
- σ_D^2 : Demand Variance (variability in demand during the lead time)
- \bar{D} : Average Demand (mean quantity of demand per time period)
- σ_L^2 : Lead Time Variance (variability in lead time)

29. Queuing Theory


29.1. M/M/1

- Arrival rate (λ): Average number of customers arriving per unit of time
- Service rate (μ): Average number of customers served per unit of time



Exponential Distribution:
Inter-arrival Times

Rate parameter $\lambda = 2$ (same as above)



Poisson Distribution:
Number of Arrivals in Time Period $t = 2$

Rate parameter $\lambda = 2$, so $\lambda t = 4$

Key Relationship

Both distributions share the same rate parameter $\lambda = 2$

- If inter-arrival $\times \sim \text{Exponential}(\lambda)$, then arrivals in time $t \sim \text{Poisson}(\lambda t)$
- Mean inter-arrival time = $\frac{1}{\lambda} = 0.5$
- Mean arrivals per unit time = $\lambda = 2$
- Mean arrivals in time period 2 = $\lambda t = 4$

1. Utilization (ρ)

Fraction of time the server is busy

$$\rho = \frac{\lambda}{\mu}$$

2. Average Number of Customers in the System (L)

Average number of customers (both waiting and being served) in the system

$$L = \frac{\rho}{1 - \rho}$$

3. Average Number of Customers in the Queue (L_q)

Average number of customers waiting in the queue

$$L_q = \frac{\rho^2}{1 - \rho}$$

4. Average Time a Customer Spends in the System (W)

Average time a customer spends in the system (from arrival until they are done being served)

$$W = \frac{1}{\mu - \lambda}$$

5. Average Waiting Time in the Queue (W_q)

Average time a customer spends just waiting in line before being served

$$W_q = \frac{\rho}{\mu - \lambda}$$

6. Probability that the System is Empty (P_0)

Probability that there are zero customers in the system (no one is being served and no one is waiting)

$$P_0 = 1 - \rho$$

7. Probability that n Customers are in the System (P_n)

Probability that there are n customers in the system (either waiting or being served)

$$P_n = (1 - \rho) \cdot \rho^n$$

8. Probability the Queue is Full (if the Queue has Limited Capacity) ($P_{n_{\max}}$)

Probability that the system is at full capacity

$$P_{n_{\max}} = (1 - \rho) \cdot \rho^{n_{\max}}$$

9. System Throughput

Rate at which customers are served and leave the system

$$\text{Throughput} = \lambda$$

10. Expected Time in Service (W_s)

Average time a customer spends actually being served (not including waiting time)

$$W_s = \frac{1}{\mu}$$

11. Idle Time ($1 - \rho$)

Fraction of time that the server is idle (i.e., not serving any customers)

$$\text{Idle Time} = 1 - \rho$$

12. Probability of Having to Wait in the Queue (P_w)

Probability that an arriving customer will have to wait before being served, i.e., that the server is busy when the customer arrives

$$P_w = \rho$$

13. Variance of the Number of Customers in the System ($\text{Var}(L)$)

Variance of the number of customers in the system

$$\text{Var}(L) = \frac{\rho}{(1 - \rho)^2}$$

A bank with a single teller

- **Arrival rate (λ):** On average, 4 customers arrive every 10 minutes ($\lambda = 4$ customers per 10 minutes)
- **Service Rate (μ):** The teller can serve 6 customers every 10 minutes ($\mu = 6$ customers per 10 minutes)

1. Utilization (ρ)

$$\rho = \frac{\lambda}{\mu} = \frac{4}{6} = 0.67$$

The teller is busy 67% of the time. The remaining 33% of the time, the teller is idle, waiting for the next customer

2. Average Number of Customers in the System (L)

$$L = \frac{\rho}{1 - \rho} = \frac{0.67}{1 - 0.67} = 2$$

On average, there are 2 customers in the coffee shop at any given time, either being served or waiting in line

3. Average Number of Customers in the Queue (L_q)

$$L_q = \frac{\rho^2}{1 - \rho} = \frac{0.67^2}{1 - 0.67} = 1.33$$

On average, about 1.33 customers are waiting in line at any time

4. Average Time a Customer Spends in the System (W)

$$W = \frac{1}{\mu - \lambda} = \frac{1}{6 - 4} = 0.5$$

On average, a customer spends 5 minutes (0.5×10 minutes) in the shop (including both waiting in line and getting served)

5. Average Waiting Time in the Queue (W_q)

$$W_q = \frac{\rho}{\mu - \lambda} = \frac{0.67}{6 - 4} = 0.33$$

On average, a customer waits 3.3 minutes (0.33×10 minutes) in line before being served by the teller

6. Probability that the System is Empty (P_0)

$$P_0 = 1 - \rho = 1 - 0.67 = 0.33$$

There is a 33% chance that the coffee shop is empty, meaning there is no customer in the queue or being served

7. Probability that n Customers are in the System (P_n)

$$P_n = (1 - \rho) \cdot \rho^n = (1 - 0.67) \cdot 0.67^2 = 0.148$$

There is a 14.8% chance that exactly 2 customers are either in line or being served

8. Probability the Queue is Full (if the Queue has Limited Capacity) ($P_{n_{\max}}$)

$$P_{n_{\max}} = (1 - \rho) \cdot p^{n_{\max}} = (1 - 0.67) \cdot 0.67^5 = 0.028$$

There is a 2.8% chance that the system is full, and no new customers can enter

9. System Throughput

$$\text{Throughput} = \lambda = 4$$

The coffee shop serves 4 customers every 10 minutes, on average

10. Expected Time in Service (W_s)

$$W_s = \frac{1}{\mu} = \frac{1}{6} = 0.167$$

On average, a customer spends 1.67 minutes being served by the teller

11. Idle Time ($1 - \rho$)

$$\text{Idle Time} = 1 - \rho = 1 - 0.67 = 0.33$$

The teller is idle 33% of the time

12. Probability of Having to Wait in the Queue (P_w)

$$P_w = \rho = 0.67$$

There is a 67% chance that a customer will have to wait when they arrive

13. Variance of the Number of Customers in the System ($\text{Var}(L)$)

$$\text{Var}(L) = \frac{\rho}{(1 - \rho)^2} = \frac{0.67}{(1 - 0.67)^2} = 6.12^2$$

The queue length varies significantly, with a variance of 6.12 customers

Costs

- **Cost per Waiting Customer per Unit Time (C_w):** cost incurred per customer for each unit of time they spend waiting in the queue

$$\text{Total Waiting Cost} = L_q \times C_w \times \text{Unit Time}$$

- **Cost per Idle Server per Unit Time (C_s):** cost incurred per unit time when the server is not serving customers

$$\text{Total Idle Cost} = (1 - \rho) \times C_s \times \text{Unit Time}$$

Total Cost

$$\text{Total Cost} = \text{Total Waiting Cost} + \text{Total Idle Cost}$$

- Arrival Rate (λ): 4 customers per 10 minutes
- Service Rate (μ): 6 customers per 10 minutes
- Cost per Waiting Customer per Hour (C_w): \$10
- Cost per Idle Server per Hour (C_s): \$20

- Operational Time: 1 hour

1. Utilization

$$\rho = \frac{\lambda}{\mu} = \frac{4}{6} = 0.67$$

2. Average Number of Customers in Queue (L_q)

$$L_q = \frac{\rho^2}{1 - \rho} = \frac{0.67^2}{1 - 0.67} = 1.33 \text{ customers}$$

3. Total Waiting Cost

$$\text{Total Waiting Cost} = L_q \times C_w \times \text{Unit Time} = 1.33 \times 10 \times 1 = \$13.33$$

4. Idle Time

$$\text{Idle Time} = 1 - \rho = 1 - 0.67 = 0.33$$

5. Total Idle Cost

$$\text{Total Idle Cost} = (1 - \rho) \times C_s \times \text{Unit Time} = 0.33 \times 20 \times 1 = \$6.67$$

6. Total Cost

$$\text{Total Cost} = \text{Total Waiting Cost} + \text{Total Idle Cost} = 13.33 + 6.67 = \$20$$

30. Network Optimization

30.1. Shortest Path

30.2. Dijkstra's

Step 1: Initialize

- Mark all nodes as unvisited
- Set the distance to the source as 0
- Set the distance to all other nodes as ∞
- Create a priority queue (or a simple list) to keep track of unvisited nodes and their distances.

Graph nodes: A, B, C, D

Start node: A

Distances:

- $A = 0$
- $B = \infty$
- $C = \infty$
- $D = \infty$

Step 2: Select the Unvisited Node with the Smallest Distance

Among all unvisited nodes, pick the one with the smallest known distance. Let's call this the current node.

Step 3: Update Neighboring Distances

For each neighbor of the current node:

1. Calculate the tentative distance:

$$\text{distance}(\text{current}) + \text{weight}(\text{current} \rightarrow \text{neighbor})$$

2. If this tentative distance is less than the known distance, update it.

If $A \rightarrow B$ has weight 4 and distance to A is 0:

$$\text{Tentative distance to B} = 0 + 4 = 4$$

If current distance to B is ∞ , update to 4

Step 4: Mark the Current Node as Visited

- Once all neighbors are checked, mark the current node as visited
- Visited nodes are not checked again

Step 5: Repeat Until All Nodes Are Visited

- Go back to Step 2 and repeat.
- Stop when all nodes have been visited or the smallest tentative distance among the unvisited nodes is ∞ (meaning unreachable)


Step 6: Reconstruct the Shortest Paths (Optional)

- To find the actual shortest path, track the previous node for each visited node
- Starting from the destination, backtrack through the previous nodes

30.3. Maximum Flow**30.4. Network Flow Optimization**

Types of Problems:

- Maximum Flow
- Minimum Cost Flow
- Multi-Commodity Flow




Node i

- Sink node: $b_i > 0$
- Source node: $b_i < 0$
- Transshipment node: $b_i = 0$

Has demand of b_i units

Has supply of $-b_i$ units

Neither supply or demand



Units shipped from node i to node j :

$$x_{ij}$$

Minimize:

$$Z = \sum_{i,j} c_{ij} x_{ij}$$


s.t.

$$\sum_k x_{ki} - \sum_l x_{il} = \text{or } \leq \text{ or } \geq b_i$$

$$l_{ij} \leq x_{ij} \leq u_{ij}$$

Where:

- c_{ij} : Unit cost of flow from node i to node j
- b_i : Demand on node i
- l_{ij} : flow lower bound from i to j
- u_{ij} : Flow upper bound from i to j
- $\sum_k x_{ki}$: Inflow to i
- $\sum_l x_{il}$: Outflow from j




If:

- | | |
|---|--|
| <ul style="list-style-type: none"> • Total Supply = Total Demand • Total Supply > Total Demand • Total Supply < Total Demand | $[\text{Inflow to } i] - [\text{Outflow from } i] = b_i$
$[\text{Inflow to } i] - [\text{Outflow from } i] \geq b_i$
$[\text{Inflow to } i] - [\text{Outflow from } i] \leq b_i$ |
|---|--|

Important:

- One decision variable x_{ij} for each edge (i, j)

- One flow balancing constraint for each node i



Minimize

$$Z = 30x_{12} + 40x_{14} + 50x_{23} + 35x_{35} + 40x_{53} \\ + 30x_{54} + 35x_{56} + 25x_{65} + 50x_{74} + 45x_{75} + 50x_{76}$$

s.t.

$$x_{12} + x_{14} \leq 200 \quad (\text{Node 1})$$

$$x_{12} + x_{23} \geq 100 \quad (\text{Node 2})$$

$$x_{23} + x_{53} - x_{35} \geq 60 \quad (\text{Node 3})$$


$$x_{14} + x_{54} + x_{74} \geq 80 \quad (\text{Node 4})$$

$$x_{35} + x_{65} + x_{75} - x_{53} - x_{54} - x_{56} \geq 170 \quad (\text{Node 5})$$

$$x_{56} + x_{76} - x_{65} \geq 70 \quad (\text{Node 6})$$

$$x_{76} + x_{75} + x_{74} \leq 300 \quad (\text{Node 7})$$

$$x_{ij} \geq 0 \quad \forall (i, j) \in E$$



Minimize:

$$\sum_{ij} (c_{ij}^f \cdot y_{ij}) + \sum_{ij} (c_{ij}^v \cdot x_{ij}) + \sum_i (c_i^f \cdot y_i) + \sum_i (p_i \cdot s_i) +$$

Where:

- $\sum_{ij} (c_{ij}^f \cdot y_{ij})$: Edge fixed cost contribution
- $\sum_{ij} (c_{ij}^v \cdot x_{ij})$: Variable cost contribution
- $\sum_i (c_i^f \cdot y_i)$: Node fixed cost contribution
- $\sum_i (p_i \cdot s_i)$: Penalty contribution
- $\sum_{ij} (l_{ij} \cdot x_{ij})$: Edge lead time weighted by flow
- $\sum_i l_i \cdot \sum_j x_{ij}$: Node service time weighted by flow

s.t.

$\sum_j x_{ji} - \sum_j x_{ij} = \text{or } \leq \text{or } \geq d_i$	Flow Conservation
$b_{ij}^l \leq x_{ij} \leq b_{ij}^u$	Lower & Upper Flow Bound
$x_{ij} \leq M \cdot y_{ij}$	Fixed Cost Route
$\sum_i (x_{ji} + x_{ij}) \leq M \cdot y_i$	Fixed Cost Node
$\sum_i x_{ij} + s_j \geq d_j$	Unmet Demand Penalty
$x_{ij} \geq 0 \quad \forall (i, j) \in E$	Non Negative Flow

Node Properties:


- **Node Type:** Source, sink, or intermediary.
- **Supply (Source):** Flow capacity of source nodes.
- **Holding Cost (Intermediary):** Inventory cost for stored goods.
- **Service Time:** Processing time at the node.
- **Demand:** Required flow at sink nodes.
- **Storage Capacity (Intermediary):** Maximum amount of goods that can be held at a node
- **Penalty for Unfulfilled Demand (Sink):** Cost for unmet demand.
- **Disruption Risk (All Nodes):** Probability of a node being unavailable due to unforeseen circumstances

Edge Properties:

- **Edge Type:** Transport mode (air, water, road, rail).
- **Fixed Cost:** Cost incurred for using the edge, regardless of flow.
- **Reliability:** Probability of edge availability.
- **Flow Bounds:** Minimum and maximum allowable flow.
- **Unit Cost:** Cost per unit of flow.
- **Lead Time:** Time it takes for flow to travel along the edge.
- **Environmental Impact:** Account for the carbon footprint of using certain transport modes.

30.5. Ford-Fulkerson

Find augmenting paths in the network and increase the flow until no more augmenting paths can be found




- s : Source
- A & B : Intermediate nodes
- t : Sink

Capacities:

- $s \rightarrow t$: 10
- $s \rightarrow B$: 5
- $A \rightarrow B$: 15
- $A \rightarrow t$: 10
- $B \rightarrow t$: 10

Step 1: Initialize flow to 0

All flows through the edges are initially set to 0.



Step 2: Find an augmenting path

Find an augmenting path using Depth-First Search (DFS). Start from the source s


$$s \rightarrow A \rightarrow t$$

The minimum capacity along this path is 10 (bottleneck on edge $A \rightarrow t$)

We can push a flow of 10 units along this path.

Step 3: Update the residual graph

- $s \rightarrow A$: Capacity becomes $10 - 10 = 0$ (no residual capacity)
- $A \rightarrow t$: Capacity becomes $10 - 10 = 0$ (no residual capacity)



Step 4: Find another augmenting path

Find another augmenting path.

We cannot use $s \rightarrow A$ or $A \rightarrow t$ because their capacities are 0.


$$s \rightarrow B \rightarrow t$$

The minimum capacity along this path is 5 (bottleneck on edge $s \rightarrow B$)

We can push a flow of 5 units along this path

Step 5: Update the residual graph

- $s \rightarrow B$: Capacity becomes $5 - 5 = 0$
- $B \rightarrow t$: Capacity becomes $10 - 5 = 5$



Step 6: Termination


No more augmenting paths from s to t can be found, as all edges from s are fully saturated

Maximum flow:

- $s \rightarrow A \rightarrow t$: 10 units
- $s \rightarrow B \rightarrow t$: 5 units

Thus, the maximum flow from source s to sink t is 15 units

Step 7: Final Flow Distribution



31. Optimization

32. Linear Programming

Function Type	Method	Example
Maximization to Minimization	$\max f(x) \iff$ $\min -f(x)$	$\max 3x + 2y \iff$ $\min -3x - 2y$
\geq Constraint to \leq	$g_i(x) \geq b_i \iff$ $-g_i(x) \leq -b_i$	$x + y \geq 5 \iff$ $-x - y \leq -5$
Equality Constraint	$g_i(x) = b_i \iff$ $g_i(x) \geq b_i \wedge$ $g_i(x) \leq b_i$	$x + y = 4 \iff$ $x + y \geq 4 \wedge$ $x + y \leq 4$
Slack Variable		$x + y \leq 6$ Add slack variable $s \geq 0$: $x + y + s = 6$ s is how much less is being used than the maximum limit of 6
Surplus Variable		$x + y \geq 6$ Add slack variable $s \geq 0$: $x + y - s = 6$ s is the amount above the minimum required level 6

33. Integer Programming

Function Type	Method	Example
At least k of the items in S	$\sum_{i \in S} x_i \geq k$	Produce at least 3 products: $\sum_{i \in S} x_i \geq 3$
At most k of the items in S	$\sum_{i \in S} x_i \leq k$	Open at most 2 facilities: $\sum_{i \in S} x_i \leq 2$
Exactly k of the items in S	$\sum_{i \in S} x_i = k$	Choose exactly one location: $\sum_{i \in S} x_i = 1$
If $x_a = 1$, then $x_b = 1$	$x_a \leq x_b$	If project A is done, project B must also be done
If $x_a = 1$, then $x_b = 0$	$x_a + x_b \leq 1$	Projects A and B are mutually exclusive
If $x_a = 1$, then $x_b + x_c \leq 0$	$x_b + x_c \leq M(1 - x_a)$	If project A is selected, neither project B nor C can be selected

Function Type	Method	Example
If $x_j = 0$, then at least k items in S must be selected	$\sum_{i \in S} x_i \geq k(1 - x_j)$	You have a main server represented by x_j . If it is down or not used ($x_j = 0$), then at least k backup servers (in set S) must be activated
At least k of m constraints satisfied	$z_i = \begin{cases} 1 & \text{if } g_i(x) \leq b_i \text{ enforced} \\ 0 & \text{if } g_i(x) \leq b_i \text{ relaxed} \end{cases}$ <p>for $i = 1, \dots, m$</p> $g_i(x) - b_i \leq M_i(1 - z_i) \quad \forall i = 1, \dots, m$ $\sum_{i=1}^m z_i \geq k$ $M_i \geq \max_x (g_i(x) - b_i)$	Enforce any k of the m constraints by relaxing up to $m - k$ of them
Fixed (Setup) Costs	$y = \begin{cases} 1 & \text{if is used} \\ 0 & \text{if not used} \end{cases}$ $x \geq 0$ $x \leq M \cdot y$	If facility is used ($y = 1$), then allow production up to M ; else $x = 0$
Balanced Flow	$\sum_{\substack{i \in V \\ i \neq k}} x_{ik} = d_k$ $\sum_{\substack{i \in V \\ i \neq k}} x_{jk} = d_k$	Each node k receives and sends exactly d_k units of flow. If $d_k = 1$, this is a TSP or assignment
Unbalanced Flow	$\sum_{\substack{i \in V \\ i \neq k}} x_{ik} - \sum_{\substack{i \in V \\ i \neq k}} x_{jk} = b_k$	Each node k has net inflow minus outflow equal to b_k : <ul style="list-style-type: none"> $b_k > 0$: demand $b_k < 0$: supply $b_k = 0$: transshipment
Subtours (MTZ)	$1 \leq u_i \leq n - 1 \quad \forall i = 2, \dots, n$ $u_i - u_j + (n - 1)x_{ij} \leq n - 2$ $\forall i \neq j, i, j \in \{2, \dots, n\}$	
Subtours (SEC)	$\sum_{\substack{i \in S, j \in S \\ j \neq i}} x_{ij} \leq S - 1 \quad \forall S \subseteq V, S \geq 2$	For every subset S of the nodes V that has at least 2 nodes, the total number of arcs (or paths) within that subset – from node i to node j , where $i \neq j$ must be less than or equal to $ S - 1$

34. Non-Linear Programming

Function Type	Example	Linearization Method	Example
Binary × Binary Constraint	$z = x \cdot y$ $x, y \in \{0, 1\}$	$z \leq x$ $z \leq y$ $z \geq x + y - 1$ $x \in \{0, 1\}$	$z = 1 \iff$ $x = y = 1$
Binary × Continuous Constraint	$z = x \cdot y$ $x \in \{0, 1\}$ $y \in [L, U]$	$z \leq Ux$ $z \geq Lx$ $z \leq y - L(1 - x)$ $z \geq y - U(1 - x)$	
Continuous × Continuous Constraint	$z = x \cdot y$ $x \in [L_x, U_x]$ $y \in [L_y, U_y]$	$z \geq L_x y + L_y x - L_x L_y$ $z \geq U_x y + U_y x - U_x U_y$ $z \geq L_x y + U_y x - L_x U_y$ $z \geq U_x y + L_y x - U_x L_y$	
Absolute Value	$z = x $	$z \geq x$ $z \geq -x$	$z = x_1 - x_2 $ \iff $z \geq x_1 - x_2$ $z \geq x_2 - x_1$
Max Constraint	$y \geq \max(x_1, \dots, x_n)$	$y \geq x_i \quad \forall i = 1, \dots, n$	$y + x_1 + 3 \geq \max(x_1 - x_3, 2x_2 + 4)$ \iff $\begin{cases} y + x_1 + 3 \geq x_1 - x_3 \\ y + x_1 + 3 \geq 2x_2 + 4 \end{cases}$
Min Constraint	$y \leq \min(x_1, \dots, x_n)$	$y \leq x_i \quad \forall i = 1, \dots, n$	$y + x_1 \leq \min(x_1 - x_3, 2x_2 + 4, 0)$ \iff $\begin{cases} y + x_1 \leq x_1 - x_3 \\ y + x_1 \leq 2x_2 + 4 \\ y + x_1 \leq 0 \end{cases}$
Max Min Objective	$\max \min(x_1, \dots, x_n)$	$y = \min(x_1, \dots, x_n)$ $y \leq x_i \quad \forall i = 1, \dots, n$	
Min Max Objective	$\min \max(x_1, \dots, x_n)$	$y = \max(x_1, \dots, x_n)$ $y \geq x_i \quad \forall i = 1, \dots, n$	
Min Min Objective	$\min \min(x_1, \dots, x_n)$		
Max Max Objective	$\max \max(x_1, \dots, x_n)$		

34.1. Mathematical Programs

34.1.1. Formulation

$$\begin{array}{lll} \min & f(x_1, x_2, \dots, x_n) & \text{(objective function)} \\ s.t. & g_i(x_1, x_2, \dots, x_n) \leq b_i & \forall i \in 1, \dots, m \quad \text{(constraints)} \\ & x_j \in \mathbb{R} & \forall j \in 1, \dots, n \quad \text{(decision variables)} \end{array}$$

- m : Number of constraints
- n : Number of variables
- x_1, x_2, \dots, x_n : decision variables
- $f: \mathbb{R}^n \rightarrow \mathbb{R}$
- $g_i: \mathbb{R}^n \rightarrow \mathbb{R}$

$$\begin{array}{ll} \min & \sum_{j=1}^n c_j x_j \\ s.t. & \sum_{j=1}^n A_{ij} x_j \leq b_i \quad \forall i = 1, \dots, m \end{array}$$

- A_{ij} : constraint coefficients
- b_i : right-hand-side values (RHS)
- c_j : objective coefficients

Vector

$$\begin{array}{ll} \min & c^T x \\ s.t. & a_i^T x \leq b_i \quad \forall i = 1, \dots, m \end{array}$$

- $a_i \in \mathbb{R}^n$
- $b_i \in \mathbb{R}$
- $c \in \mathbb{R}^n$
- $x \in \mathbb{R}^n$

Matrix

$$\begin{array}{ll} \min & c^T x \\ s.t. & Ax \leq b \end{array}$$

- $A \in \mathbb{R}^{m \times n}$
- $b \in \mathbb{R}^m$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \quad c = \begin{bmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \end{bmatrix} \quad b = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$Ax = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11}x_1 & a_{12}x_2 & \dots & a_{1n}x_n \\ a_{21}x_1 & a_{22}x_2 & \dots & a_{2n}x_n \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}x_1 & a_{m2}x_2 & \dots & a_{mn}x_n \end{bmatrix} \leq \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

$$c^T x = [c_1, c_2, \dots, c_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \sum_{j=1}^n c_j x_j$$

34.1.2. Transformation

- **Min and Max**

$$\max f(x) \Leftrightarrow \min -f(x)$$

- **= and \geq**

$$g_i(x) \geq b_i \Leftrightarrow -g_i(x) \leq -b_i$$

$$g_i(x) = b_i \Leftrightarrow g_i(x) \geq b_i \wedge g_i(x) \leq b_i$$

- **E.g.,**

$$\begin{array}{llll} \max & x_1 & - & x_2 \\ s.t. & -2x_1 & + & x_2 \geq -3 \\ & x_1 & + & 4x_2 = 5 \end{array} \Leftrightarrow \begin{array}{llll} \min & -x_1 & + & x_2 \\ s.t. & 2x_1 & - & x_2 \leq 3 \\ & x_1 & + & 4x_2 \leq 5 \\ & -x_1 & + & 4x_2 \leq -5 \end{array}$$

34.1.3. Constraints

- **Sign Constraints**


$$x_i \geq 0 \text{ or } x_i \leq 0$$

- **Functional Constraints**

$$\sum_{i=1}^n c_i x_i \leq b_i$$

34.1.4. Feasible Solutions

- Feasible solution: Satisfies all constraints
- Infeasible solution: violates at least one constraint




34.1.5. Feasible Region and Optimal Solution

- Feasible Region: Set of feasible solutions
- Optimal Solution: Attains largest (maximization) or smallest (minimization) objective value

34.1.6. Binding Constraint

Let $g(\cdot) \leq b$ be an inequality constraint and $|(\cdot)$ be a solution. $g(\cdot) \leq b$ is binding at $|(\cdot)$ if $g(|(\cdot)) = b$

$2x_1 + x_2 \leq 100$	is binding (active) at the point	$(x_1, x_2) = (30, 40)$
$2x_1 + x_2 \leq 100$	is non-binding (inactive) at the point	$(x_1, x_2) = (20, 20)$




34.1.7. Strict and Weak Inequalities


- Strict: The two sides **cannot be equal** (e.g., $x_1 + x_2 > 5$)
- Weak: The two sides **can be equal** (e.g., $x_1 + x_2 \geq 5$)

34.1.8. Graphical Approach

$$\begin{aligned}
 \max \quad & 40x_1 + 30x_2 \\
 \text{s.t.} \quad & 2x_1 + x_2 \leq 100 \\
 & x_1 + 2x_2 \leq 80
 \end{aligned}$$



34.1.9. Types of LP



34.1.10. LP Formulation

Production

We produce desks and tables

- Producing a desk requires 3 units of wood, 1 hour of labor and 50 minutes of machine time
- Producing a table requires 5 units of wood, 2 hours of labor and 20 minutes of machine time

For each day, we have

- 200 workers that each work 8 hours
- 50 machines that each run for 16 hours
- A supply of 3600 units of wood

Desks and tables are sold \$700 and \$900 per unit, respectively

Everything that is produced is sold

1. Define Variables

x_1 : number of desks produced per day

x_2 : number of tables produced per day

2. Formulate Objective Function

$$\max \quad 700x_1 + 900x_2$$

3. Formulate Constraints

Resource	Consumption per		Total supply
	Desk	Table	
Wood	3 units	5 units	3600 units
Labor	1 hours	2 hours	200 workers \times 8 hours = 1600 hours
Machine time	50 minutes	20 minutes	50 machines \times 16 hours = 800 hours

$$\begin{aligned} 3x_1 + 5x_2 &\leq 3600 \\ x_1 + 2x_2 &\leq 1600 \\ 50x_1 + 20x_2 &\leq 48000 \end{aligned}$$

4. Complete Formulation

$$\begin{aligned} \max \quad & 700x_1 + 900x_2 \\ \text{s.t.} \quad & 3x_1 + 5x_2 \leq 3600 \quad (\text{wood}) \\ & x_1 + 2x_2 \leq 1600 \quad (\text{labor}) \\ & 50x_1 + 20x_2 \leq 48000 \quad (\text{machine}) \\ & x_1 \geq 0 \\ & x_2 \geq 0 \end{aligned}$$

Compact Formulation

$$\begin{aligned} \max \quad & \sum_{j=1}^n P_j x_j \\ \text{s.t.} \quad & \sum_{j=1}^n A_{ij} x_j \leq R_i \quad \forall i = 1, \dots, m \\ & x_j \geq 0 \quad \forall j = 1, \dots, n \end{aligned}$$

Where:

- n : number of products
- m : number of resources
- j : indices of products
- i : indices of resources

- P_j : unit sale price of product j
- R_i : supply limit of resource i
- A_{ij} : unit of resource i to produce one unit of product j
- $i = 1, \dots, m$
- $j = 1, \dots, n$
- x_j : production quantity for product $j, j = 1, \dots, n$

model.py

```

from pyomo.environ import *
from pyomo.dataportal import DataPortal

model = AbstractModel()

# Sets
model.Products = Set()
model.Resources = Set()

# Parameters
model.Profit = Param(model.Products)
model.Supply = Param(model.Resources)
model.Consumption = Param(model.Resources, model.Products)

# Variables
model.x = Var(model.Products, domain=NonNegativeReals)

# Objective: Maximize profit
def objective_rule(model):
    return sum(model.Profit[j] * model.x[j] for j in model.Products)
model.OBJ = Objective(rule=objective_rule, sense=maximize)

# Constraints: Do not exceed resource supply
def constraint_rule(model, i):
    return sum(model.Consumption[i, j] * model.x[j] for j in model.Products) <=
model.Supply[i]
model.ResourceConstraint = Constraint(model.Resources, rule=constraint_rule)

# Load data from .dat file
data = DataPortal()
data.load(filename='data.dat', model=model)

# Create an instance of the model
instance = model.create_instance(data)

# Create solver
solver = SolverFactory('glpk')
solver.options['tmlim'] = 60

# Solve with solver timeout (optional)
results = solver.solve(instance, tee=True)

# Display results
instance.display()

```

data.dat

```
set Products := Desk Table ;
set Resources := Wood Labor Machine ;

param Profit :=
Desk    700
Table   900 ;

param Supply :=
Wood    3600
Labor   1600
Machine 48000 ;

param Consumption:
      Desk Table :=
Wood    3      5
Labor   1      2
Machine 50     20 ;
```

Output:

```
Variables:
x : Size=2, Index=Products
  Key : Lower : Value          : Upper : Fixed : Stale : Domain
  Desk :      0 : 884.210526315789 : None : False : False :
NonNegativeReals
  Table :      0 : 189.473684210526 : None : False : False :
NonNegativeReals

Objectives:
OBJ : Size=1, Index=None, Active=True
  Key : Active : Value
  None : True : 789473.6842105257

Constraints:
ResourceConstraint : Size=3
  Key : Lower : Body          : Upper
  Labor : None : 1263.157894736841 : 1600.0
  Machine : None : 47999.99999999997 : 48000.0
  Wood : None : 3599.99999999997 : 3600.0
```

Multi Period (Inventory)

We produce and sell products

For the coming 4 days, the demand will be

- Days 1: 100
- Days 2: 150
- Days 3: 200
- Days 4: 170

The unit production costs are different for different days


- Days 1: \$9
- Days 2: \$12
- Days 3: \$10
- Days 4: \$12

Prices are all fixed. So maximizing profit is equivalent to minimizing costs.

We may store products and sell them later. Inventory cost is \$1 per unit per day.

E.g., producing 620 units on day 1 to fulfill all demands:

$$\$9 \times 620 + \$1 \times 150 + \$2 \times 200 + \$3 \times 170 = \$6,640$$



$$I_{t-1} + x_t - d_t = I_t$$


- I_{t-1} : inventory at the end of day $t-1$ (i.e., beginning inventory for day t)
- x_t : units produced on day t
- d_t : demand (or sales) on day t
- I_t : inventory at the end of day t

Inventory costs are calculated according to **ending inventory**

- x_t : production quantity of day t , $t = 1, \dots, 4$
- y_t : **ending** inventory of day t , $t = 1, \dots, 4$

Objective function:

$$\min \quad 9x_1 + 12x_2 + 10x_3 + 12x_4 + y_1 + y_2 + y_3 + y_4$$



Inventory balancing constraints:

- Day 1: $x_1 - 100 = y_1$
- Day 2: $y_1 + x_2 - 150 = y_2$
- Day 3: $y_2 + x_3 - 200 = y_3$
- Day 4: $y_3 + x_4 - 170 = y_4$

Demand fulfillment constraints:

- $x_1 \geq 100$
- $y_1 + x_2 \geq 150$
- $y_2 + x_3 \geq 200$
- $y_3 + x_4 \geq 170$

Non-negativity constraints:

- $y_t \geq 0$
- $x_t \geq 0$

Complete Formulation

$$\begin{aligned} \min \quad & 9x_1 + 12x_2 + 10x_3 + 12x_4 + y_1 + y_2 + y_3 + y_4 \\ \text{s.t.} \quad & x_1 - 100 = y_1 \\ & y_1 + x_2 - 150 = y_2 \\ & y_2 + x_3 - 200 = y_3 \\ & y_3 + x_4 - 170 = y_4 \\ & x_1 \geq 100 \\ & y_1 + x_2 \geq 150 \\ & y_2 + x_3 \geq 200 \\ & y_3 + x_4 \geq 170 \\ & x_t, y_t \geq 0 \quad \forall t = 1, \dots, 4 \end{aligned}$$

Simplification (Redundant Constraints)

- Inventory balancing & non-negativity imply fulfillment
 - E.g., in day 1, $x_1 - 100 = y_1$ and $y_1 \geq 0$ means $x_1 \geq 100$

$$\begin{aligned} \min \quad & 9x_1 + 12x_2 + 10x_3 + 12x_4 + y_1 + y_2 + y_3 + y_4 \\ \text{s.t.} \quad & x_1 - 100 = y_1 \\ & y_1 + x_2 - 150 = y_2 \\ & y_2 + x_3 - 200 = y_3 \\ & y_3 + x_4 - 170 = y_4 \\ & x_t, y_t \geq 0 \quad \forall t = 1, \dots, 4 \end{aligned}$$

- No need to have ending inventory in period 4 (costly but useless)

$$\begin{aligned} \min \quad & 9x_1 + 12x_2 + 10x_3 + 12x_4 + y_1 + y_2 + y_3 + y_4 \\ \text{s.t.} \quad & x_1 - 100 = y_1 \\ & y_1 + x_2 - 150 = y_2 \\ & y_2 + x_3 - 200 = y_3 \\ & y_3 + x_4 - 170 = 0 \\ & x_t \geq 0 \quad \forall t = 1, \dots, 4 \\ & y_t \geq 0 \quad \forall t = 1, \dots, 3 \end{aligned}$$

Compact Formulation

$$\begin{aligned} \min \quad & \sum_{t=1}^4 (C_t x_t + y_t) \\ \text{s.t.} \quad & y_t - 1 + x_t - D_t = y_t \quad \forall t = 1, \dots, 4 \\ & y_0 = 0 \\ & x_t, y_t \geq 0 \quad \forall t = 1, \dots, 4 \end{aligned}$$

Where:

- D_t : demand on day t
- y_t : ending inventory of day t
- C_t : unit production cost on day t

Personnel Scheduling

Scheduling employees

Each employee must work for 5 consecutive days and then take 2 consecutive rest days

Number of employees required for each day

Mon	Tue	Wed	Thu	Fri	Sat	Sun
110	80	150	30	70	160	120

Seven shifts

- Mon to Fri
- Tue to Sat
- ...

Minimize number of employees hired

Let x_i be the number of employees assigned to work from day i for 5 consecutive days

Objective function:

$$\min x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7$$

Demand Fulfilment Constraints

- 110 employees needed Monday

$$x_1 + x_4 + x_5 + x_6 + x_7 \geq 110$$

An employee that works on Tues (t_2) or Wed (t_3) does not work on Mon (t_1)

- 80 employees needed Tuesday

$$x_1 + x_2 + x_5 + x_6 + x_7 \geq 80$$

An employee that works on Wed (t_3) or Thu (t_4) does not work on Tue (t_2)

- 120 employees needed Sunday

$$x_3 + x_4 + x_5 + x_6 + x_7 \geq 120$$

An employee that works on Mon (t_1) or Tue (t_2) does not work on Sun (t_7)

Non-Negativity Constraints

$$x_i \geq 0 \quad \forall i = 1, \dots, 7$$

Complete Formulation

$$\begin{aligned} \min \quad & x_1 + x_2 + x_3 + x_4 + x_5 + x_6 + x_7 \\ \text{s.t.} \quad & x_1 + x_4 + x_5 + x_6 + x_7 \geq 110 \\ & x_1 + x_2 + x_5 + x_6 + x_7 \geq 80 \\ & x_1 + x_2 + x_3 + x_6 + x_7 \geq 150 \\ & x_1 + x_2 + x_3 + x_4 + x_7 \geq 30 \\ & x_1 + x_2 + x_3 + x_4 + x_5 \geq 70 \\ & x_2 + x_3 + x_4 + x_5 + x_6 \geq 160 \\ & x_3 + x_4 + x_5 + x_6 + x_7 \geq 120 \\ & x_i \geq 0 \quad \forall i = 1, \dots, 7 \end{aligned}$$

model.py

```
from pyomo.environ import *
from pyomo.dataportal import DataPortal

model = AbstractModel()

# Sets
model.Days = Set(ordered=True)          # Days 1..7 (Mon..Sun)
model.Shifts = Set(ordered=True)        # Shifts starting on days 1..7

# Parameters
model.Demand = Param(model.Days)        # Daily staffing requirements
model.Cover = Param(model.Days, model.Shifts, within=Binary) # Coverage matrix
# (1 if shift j covers day i)

# Variables
model.x = Var(model.Shifts, domain=NonNegativeReals)

# Objective: Minimize total employees hired
def obj_rule(model):
    return sum(model.x[s] for s in model.Shifts)
model.OBJ = Objective(rule=obj_rule, sense=minimize)

# Constraints: Cover daily demand
def demand_rule(model, d):
    return sum(model.Cover[d, s] * model.x[s] for s in model.Shifts) >=
model.Demand[d]
model.DemandConstraint = Constraint(model.Days, rule=demand_rule)

# Load data from .dat file
data = DataPortal()
data.load(filename='data.dat', model=model)
```



```

# Create an instance of the model
instance = model.create_instance(data)

# Create solver
solver = SolverFactory('glpk')
solver.options['tmlim'] = 60

# Solve with solver timeout (optional)
results = solver.solve(instance, tee=True)

# Display results
instance.display()

```

data.dat

```

set Days := Mon Tue Wed Thu Fri Sat Sun ;
set Shifts := s1 s2 s3 s4 s5 s6 s7 ;

param Demand :=
Mon 110
Tue 80
Wed 150
Thu 30
Fri 70
Sat 160
Sun 120 ;

# Each shift covers 5 consecutive days starting on the shift's day
# 1 if shift s_j covers day d_i, 0 otherwise

param Cover:
      s1 s2 s3 s4 s5 s6 s7 :=
Mon   1  0  0  1  1  1  1
Tue   1  1  0  0  1  1  1
Wed   1  1  1  0  0  1  1
Thu   1  1  1  1  0  0  1
Fri   1  1  1  1  1  0  0
Sat   0  1  1  1  1  1  0
Sun   0  0  1  1  1  1  1 ;

```

Output:

```

Variables:
x : Size=7, Index=Shifts
   Key : Lower : Value          : Upper : Fixed : Stale : Domain
   s1 :      0 : 3.333333333333333 : None : False : False : NonNegativeReals
   s2 :      0 : 40.0 : None : False : False : NonNegativeReals
   s3 :      0 : 13.3333333333333 : None : False : False : NonNegativeReals
   s4 :      0 : 0.0 : None : False : False : NonNegativeReals
   s5 :      0 : 13.3333333333333 : None : False : False : NonNegativeReals
   s6 :      0 : 93.3333333333333 : None : False : False : NonNegativeReals
   s7 :      0 : 0.0 : None : False : False : NonNegativeReals

```

```
Objectives:
  OBJ : Size=1, Index=None, Active=True
        Key   : Active : Value
        None  :   True  : 163.33333333333323
```

```
Constraints:
  DemandConstraint : Size=7
        Key : Lower : Body           : Upper
        Fri  :  70.0 : 69.99999999999993 : None
        Mon  : 110.0 : 109.99999999999993 : None
        Sat  : 160.0 : 159.99999999999999 : None
        Sun  : 120.0 : 119.99999999999999 : None
        Thu  :  30.0 : 56.66666666666663  : None
        Tue  :  80.0 : 149.99999999999994 : None
        Wed  : 150.0 : 149.99999999999994 : None
```

34.2. IP (Integer Programming)

34.3. Selection and Logical Relations on Binary Variables $x_i \in \{0, 1\}$

34.3.1. Cardinality Constraints

Given a subset $S \subseteq \{1, \dots, n\}$:

- At least k of the items in S

$$\sum_{i \in S} x_i \geq k$$

We want to select at least 2 of items 1, 2, 3

$$x_1 + x_2 + x_3 \geq 2$$

- At most k of the items in S

$$\sum_{i \in S} x_i \leq k$$

We want to select at most 2 of items 1, 2, 3

$$x_1 + x_2 + x_3 \leq 2$$

- Exactly k of the items in S

$$\sum_{i \in S} x_i = k$$

We want to select exactly 2 of items 1, 2, 3

$$x_1 + x_2 + x_3 = 2$$

34.3.2. Logical OR

Given a subset $S \subseteq \{1, \dots, n\}$:

- At least 1 item in S

$$\sum_{i \in S} x_i \geq 1$$

Select item 1 or item 2, or both

$$x_1 + x_2 \geq 1$$

34.3.3. Conditional Selection: “If X, then Y”

(Implication Constraints)

- If $x_a = 1$, then $x_b = 1$

$$x_a \leq x_b$$

If item 1 is selected, then item 2 must also be selected

$$x_1 \leq x_2$$

x_a	x_b	Valid?
0	0	✓
0	1	✓
1	0	✓
1	1	✗

- If $x_a = 1$, then $x_b = 0$

$$x_a + x_b \leq 1$$

If you choose project A, then you cannot choose project B (mutually exclusive).

x_a	x_b	Valid?
0	0	✓
0	1	✓
1	0	✓
1	1	✗

- If $x_a = 1$, then $x_b + x_c \leq 0$ (generalized)

$$x_b + x_c \leq M(1 - x_a)$$

For binary variables $M = 2$ works

If you choose project A, then you must not choose either project B or C.

x_a	x_b	x_c	Valid?
-------	-------	-------	--------

1	0	0	✓
1	1	0	✗
1	1	1	✗
0	1	1	✓
0	0	1	✓

34.3.4. Conditional OR: “If not X, then select Y and Z”

- If $x_j = 0$, then at least k items in S must be selected (generalized)

$$\sum_{i \in S} x_i \geq k(1 - x_j)$$

If you do not choose item 1, then you must choose both item 2 and item

If $x_1 = 0$, then $x_2 = x_3 = 1$

$$x_2 + x_3 \geq 2(1 - x_1)$$

x_1	x_2	x_3	LHS	RHS	Valid?
0	1	1	2	2	✓
0	1	0	1	2	✗
0	0	0	0	2	✗
1	0	0	0	0	✓
1	1	0	1	0	✓
1	1	1	2	0	✓

34.3.5. XOR / Exclusive Conditions

- Exactly one item in a set S

$$\sum_{i \in S} x_i = 1$$

- Select one or the other but not both (i.e., $x_i \oplus x_k$)

$$x_a + x_b = 1$$

You can select item 1 or item 2, but not both

x_1	x_2	Sum	Valid?
0	1	1	✓
1	0	1	✓
1	1	2	✗

0	0	0	✗
---	---	---	---

34.4. Constraint Activation and Relaxation via Binary Indicators $z_i \in \{1, 0\}$

34.4.1. Base Case: At Least One of Two Constraints Satisfied

You are given 2 constraints

$$\begin{aligned} g_1(x) &\leq b_1 \\ g_2(x) &\leq b_2 \end{aligned}$$

We want at least one of these two constraints to be satisfied.

Binary Indicator Variables

Define:

$$z_i = \begin{cases} 1 & \text{if } g_{i(x)} \leq b_i \text{ is enforced} \\ 0 & \text{if } g_{i(x)} \leq b_i \text{ may be violated} \end{cases} \quad \text{for } i = 1, 2$$

Let M_1 and M_2 be upper bounds on how much each constraint's LHS $g_{i(x)}$ can exceed b_i :

$$M_i \geq \max_x (g_{i(x)} - b_i)$$

Big-M Reformulation

We relax each constraint as follows:

$$\begin{aligned} g_1(x) - b_1 &\leq M_1(1 - z_1) \\ g_2(x) - b_2 &\leq M_2(1 - z_2) \end{aligned}$$

When $z_i = 1$, the original constraint is enforced

When $z_i = 0$, the original constraint can be violated by up to M_i

At Least One Constraint Enforced

We require:

$$z_1 + z_2 \geq 1$$

34.4.2. General Formulation for m Constraints with At Least k Satisfied

1. Introduce a binary variable $z_i \in \{0, 1\}$ for each constraint $g_{i(x)} \leq b_i$, where:

- $z_i = 1$: the constraint is enforced (i.e., $g_{i(x)} \leq b_i$)
- $z_i = 0$: the constraint may be violated

2. Use “big-M” constraint to relax $g_{i(x)} \leq b_i$ when $z_i = 0$, like:

$$g_{i(x)} - b_i \leq M_i(1 - z_i), \quad \forall i = 1, \dots, m$$

Where $M_i \geq \max_x (g_{i(x)} - b_i)$

3. Impose a condition that at least k of the m constraints must be satisfied:

$$\sum_{i=1}^m z_i \geq k$$

34.5. Fixed Charge Constraints (Setup Costs)

- n : Number of factories
- K_i : Capacity of factory i
- C_i : Unit production cost at factory i
- S_i : Fixed setup cost for factory i
- D : Total Demand

Decision Variables

- x_i : production quantity at factory i , $i = 1, \dots, n$
- y_i : Binary variable indicating whether factory i is used

$$y_i = \begin{cases} 1 & \text{if factory } i \text{ is used (i.e., } x_i > 0) \\ 0 & \text{if factory } i \text{ is not used (i.e., } x_i = 0) \end{cases}$$

Objective Function

$$\min \underbrace{\sum_{i=1}^n C_i x_i}_{\text{Production Cost}} + \underbrace{\sum_{i=1}^n S_i y_i}_{\text{Setup Cost}}$$

Capacity Constraints

$$x_i \leq K_i \quad \forall i = 1, \dots, n$$

Demand Fullfilment

$$\sum_{i=1}^n x_i \geq D$$

Logical Linking of x_i and y_i

To ensure $y_i = 1$ if any production occurs at factory i :

$$x_i \leq K_i y_i \quad \forall i = 1, \dots, n$$

- If $x_i > 0$, this forces $y_i = 1$
- If $x_i = 0$, y_i can be 0 or 1 (but to minimize cost, y_i will be 0)

Binary & Non-negative Constraints

$$y_i \in \{0, 1\}, \quad x_i \geq 0 \quad \forall i = 1, \dots, n$$

General Form (Big-M Formulation)

$$x \leq My$$

Where M must be set to the upper bound of x

34.6. Facility Location

Problem	Definition
Set Covering	Select the minimum number of facilities such that every demand point is within a specified coverage distance of at least one facility
Maximum Covering	Select a fixed number of facilities to maximize the number of demand points covered within a specified distance

Problem	Definition
Fixed Charge Location	Determine which facilities to open and how to assign demand to them to minimize the total cost

34.6.1. Set Covering Problem

Question: How to allocate as few facilities as possible to cover all demand nodes?

- I : Set of demand nodes
- J : Set of location nodes
- $d_{ij} > 0$: Distance between demand node $i \in I$ and location node $j \in J$
- s : Maximum allowable distance (a facility at j can cover demand node i if $d_{ij} < s$)
- Demand i is “covered” by location j if $d_{ij} < s$
- $w_j > 0$: Cost of building facility at location j (usually set to $w_j = 1$ for minimizing number of facilities, but can reflect other costs)
- $a_{ij} = \begin{cases} 1 & \text{if } d_{ij} < s \\ 0 & \text{otherwise} \end{cases}$: Indicator of whether location j can cover demand i
- $x_j = \begin{cases} 1 & \text{if facility is built at location } j \\ 0 & \text{otherwise} \end{cases}$

Complete Formulation

$$\begin{aligned}
 \min \quad & \sum_{j \in J} w_j x_j \\
 s.t. \quad & \sum_{j \in J} a_{ij} x_j \geq 1 \quad \forall i \in I \\
 & x_j \in \{0, 1\} \quad \forall j \in J
 \end{aligned}$$

34.6.2. Maximum Covering Problem

Question: How to allocate at most p facilities to cover as many demand nodes as possible?

- I : Set of demand nodes
- J : Set of location nodes
- $d_{ij} > 0$: Distance between demand node $i \in I$ and location node $j \in J$
- $p \in \mathbb{N}$: maximum number of facilities
- $w_j > 0$: Cost of building facility at location j (usually set to $w_j = 1$ for minimizing number of facilities, but can reflect other costs)
- $x_j = \begin{cases} 1 & \text{if facility is built at location } j \\ 0 & \text{otherwise} \end{cases}$
- $a_{ij} = \begin{cases} 1 & \text{if } d_{ij} < s \\ 0 & \text{otherwise} \end{cases}$: Indicator of whether location j can cover demand i
- $y_i = \begin{cases} 1 & \text{if demand } i \in I \text{ is covered by any facility} \\ 0 & \text{otherwise} \end{cases}$

Complete Formulation

$$\begin{aligned}
\min \quad & \sum_{i \in I} w_i y_i \\
s.t. \quad & \sum_{j \in J} a_{ij} x_j \geq y_i \quad \forall i \in I \\
& \sum_{j \in J} x_j \leq p \quad \forall j \in J \\
& x_j \in \{0, 1\} \quad \forall j \in J \\
& y_i \in \{0, 1\} \quad \forall i \in I
\end{aligned}$$

Fixed Charge Location Problems

Question: How to allocate some facilities to minimize the total shipping and construction costs?

Uncapacitated Facility Location Problem (UFL)

- I : Set of demand nodes
- J : Set of location nodes
- $h_i > 0$: Demand size at demand node i
- d_{ij} : Unit shipping cost from location node j to demand node i
- $f_j > 0$: Fixed construction cost at location j
- $x_j = \begin{cases} 1 & \text{if facility is built at location } j \\ 0 & \text{otherwise} \end{cases}$
- $y_{ij} = \begin{cases} 1 & \text{if demand } i \in I \text{ is served by facility at location } j \in J \\ 0 & \text{otherwise} \end{cases}$

Complete Formulation

$$\begin{aligned}
\min \quad & \underbrace{\sum_{i \in I} \sum_{j \in J} h_i d_{ij} y_{ij}}_{\text{Shipping Costs}} + \underbrace{\sum_{j \in J} f_j x_j}_{\text{Fixed Costs}} \\
s.t. \quad & y_{ij} \leq x_j \quad \forall i \in I, j \in J \\
& \sum_{j \in J} y_{ij} = 1 \quad \forall i \in I \\
& x_j \in \{0, 1\} \quad \forall j \in J \\
& y_{ij} \in \{0, 1\} \quad \forall i \in I
\end{aligned}$$

Capacitated Facility Location Problem (CFL)

If locations have a capacity, we may add constraint

$$\sum_{i \in I} h_i y_{ij} \leq K_j \quad \forall j \in J$$

Where:

- $K_j > 0$: Capacity of location j

$$\begin{aligned}
\min \quad & \underbrace{\sum_{j=1}^5 f_j x_j}_{\text{Fixed Costs}} + \underbrace{\sum_{i=1}^5 \sum_{j=1}^5 c_{ij} y_{ij}}_{\text{Shipping Costs}} \\
s.t. \quad & \sum_{i=1}^5 y_{ij} \leq K_j x_j \quad \forall j = 1, \dots, 5 & (\text{Capacity Constraint}) \\
& \sum_{j=1}^5 y_{ij} \geq D_i \quad \forall i = 1, \dots, 5 & (\text{Demand Constraint}) \\
& x_j \in \{0, 1\} \quad \forall j = 1, \dots, 5 & (\text{Binary Constraint}) \\
& y_{ij} \geq 0 \quad \forall i = 1, \dots, 5, j = 1, \dots, 5 & (\text{Positivity Constraint})
\end{aligned}$$

Parameters

- f_j : weekly operating cost of distribution center j
- c_{ij} : shipping cost per book from distribution center j to region i
- K_j : capacity of distribution center j
- D_i : book demand of region i

Decision variable

- x_{ij} : binary variable

$$x_{ij} = \begin{cases} 1 & \text{if a distribution center is build at location } j \\ 0 & \text{otherwise} \end{cases}$$

- y_{ij} : number of books shipped from distribution center j to region i

Supply

	WA	NV	NE	PA	FL	Demand (D_i)
NorthWest	y_{11}	y_{12}	y_{13}	y_{14}	y_{15}	8000
SouthWest	y_{21}	y_{22}	y_{23}	y_{24}	y_{25}	12000
MidWest	y_{31}	y_{32}	y_{33}	y_{34}	y_{35}	9000
SouthEast	y_{41}	y_{42}	y_{43}	y_{44}	y_{45}	14000
NorthEast	y_{51}	y_{52}	y_{53}	y_{54}	y_{55}	17000

Fixed Costs (f_j) and Capacity (K_j)

	x_1	x_2	x_3	x_4	x_5
Operation Cost (f_i)	40000	30000	25000	40000	30000
Capacity (K_i)	20000	20000	15000	25000	15000

Shipping Cost (c_j)

	WA	NV	NE	PA	FL
NorthWest	2.4	3.25	4.05	5.25	6.95
SouthWest	3.5	2.3	3.25	6.05	5.85

	WA	NV	NE	PA	FL
MidWest	4.8	3.4	2.85	4.3	4.8
SouthEast	6.8	5.25	4.3	3.25	2.1
NorthEast	5.75	6	4.75	2.75	3.5

Binary Decision Variable

WA	x_1
NV	x_2
NE	x_3
PA	x_4
FL	x_5

model.py

```

from pyomo.environ import *
from pyomo.dataportal import DataPortal

model = AbstractModel()

# Sets
model.I = Set(doc='Regions (demand points)')
model.J = Set(doc='Candidate distribution centers')

# Parameters
model.D = Param(model.I, within=NonNegativeReals, doc='Demand at region i')
model.K = Param(model.J, within=NonNegativeReals, doc='Capacity at distribution center j')
model.f = Param(model.J, within=NonNegativeReals, doc='Fixed cost to open center j')
model.c = Param(model.I, model.J, within=NonNegativeReals, doc='Shipping cost from j to i')

# Decision Variables
model.x = Var(model.J, within=Binary, doc='1 if center j is opened')
model.y = Var(model.I, model.J, within=NonNegativeReals, doc='Units shipped from j to i')

# Objective Function: Minimize total cost
def total_cost_rule(model):
    fixed = sum(model.f[j] * model.x[j] for j in model.J)
    shipping = sum(model.c[i, j] * model.y[i, j] for i in model.I for j in model.J)
    return fixed + shipping
model.TotalCost = Objective(rule=total_cost_rule, sense=minimize)

# Constraints

# Capacity Constraint: Total shipped from center ≤ capacity if opened
def capacity_rule(model, j):
    return sum(model.y[i, j] for i in model.I) <= model.K[j] * model.x[j]

```

```

model.CapacityConstraint = Constraint(model.J, rule=capacity_rule)

# Demand Constraint: Total received by region ≥ its demand
def demand_rule(model, i):
    return sum(model.y[i, j] for j in model.J) >= model.D[i]
model.DemandConstraint = Constraint(model.I, rule=demand_rule)

# Load data from .dat file
data = DataPortal()
data.load(filename='data.dat', model=model)

# Create an instance of the model
instance = model.create_instance(data)

# Create solver
solver = SolverFactory('glpk')
solver.options['tmlim'] = 60

# Solve with solver timeout (optional)
results = solver.solve(instance, tee=True)

# Display results
instance.display()

```

data.dat

```

set I := NorthWest SouthWest MidWest SouthEast NorthEast;
set J := WA NV NE PA FL;

param: D :=
NorthWest 8000
SouthWest 12000
MidWest 9000
SouthEast 14000
NorthEast 17000;

param: f :=
WA 40000
NV 30000
NE 25000
PA 40000
FL 30000;

param: K :=
WA 20000
NV 20000
NE 15000
PA 25000
FL 15000;

param c:
      WA      NV      NE      PA      FL :=
NorthWest 2.4  3.25  4.05  5.25  6.95

```

SouthWest	3.5	2.3	3.25	6.05	5.85
MidWest	4.8	3.4	2.85	4.3	4.8
SouthEast	6.8	5.25	4.3	3.25	2.1
NorthEast	5.75	6	4.75	2.75	3.5 ;

Output:

Variables:

x : 1 if center j is opened

Size=5, Index=J

Key : Lower : Value : Upper : Fixed : Stale : Domain

FL : 0 : 1.0 : 1 : False : False : Binary

NE : 0 : 0.0 : 1 : False : False : Binary

NV : 0 : 1.0 : 1 : False : False : Binary

PA : 0 : 1.0 : 1 : False : False : Binary

WA : 0 : 0.0 : 1 : False : False : Binary

y : Units shipped from j to i

Size=25, Index=I*J

Key : Lower : Value : Upper : Fixed : Stale : Domain

('MidWest', 'FL') : 0 : 1000.0 : None : False : False :

NonNegativeReals

('MidWest', 'NE') : 0 : 0.0 : None : False : False :

NonNegativeReals

('MidWest', 'NV') : 0 : 0.0 : None : False : False :

NonNegativeReals

('MidWest', 'PA') : 0 : 8000.0 : None : False : False :

NonNegativeReals

('MidWest', 'WA') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthEast', 'FL') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthEast', 'NE') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthEast', 'NV') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthEast', 'PA') : 0 : 17000.0 : None : False : False :

NonNegativeReals

('NorthEast', 'WA') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthWest', 'FL') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthWest', 'NE') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthWest', 'NV') : 0 : 8000.0 : None : False : False :

NonNegativeReals

('NorthWest', 'PA') : 0 : 0.0 : None : False : False :

NonNegativeReals

('NorthWest', 'WA') : 0 : 0.0 : None : False : False :

NonNegativeReals

('SouthEast', 'FL') : 0 : 14000.0 : None : False : False :

NonNegativeReals

('SouthEast', 'NE') : 0 : 0.0 : None : False : False :

NonNegativeReals

```

('SouthEast', 'NV') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthEast', 'PA') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthEast', 'WA') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthWest', 'FL') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthWest', 'NE') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthWest', 'NV') :      0 : 12000.0 : None : False : False :
NonNegativeReals
('SouthWest', 'PA') :      0 :      0.0 : None : False : False :
NonNegativeReals
('SouthWest', 'WA') :      0 :      0.0 : None : False : False :
NonNegativeReals

Objectives:
  TotalCost : Size=1, Index=None, Active=True
    Key : Active : Value
      None :   True : 268950.0

Constraints:
  CapacityConstraint : Size=5
    Key : Lower : Body : Upper
    FL :  None :  0.0 :  0.0
    NE :  None :  0.0 :  0.0
    NV :  None :  0.0 :  0.0
    PA :  None :  0.0 :  0.0
    WA :  None :  0.0 :  0.0
  DemandConstraint : Size=5
    Key      : Lower : Body : Upper
    MidWest : 9000.0 : 9000.0 : None
    NorthEast : 17000.0 : 17000.0 : None
    NorthWest : 8000.0 : 8000.0 : None
    SouthEast : 14000.0 : 14000.0 : None
    SouthWest : 12000.0 : 12000.0 : None

```

34.7. Machine Scheduling

	Problem	Definition
One Machine	Single Machine Serial Production	Scheduling jobs one after another on a single machine
Multiple Machines	Multiple Parallel Machines	Scheduling jobs on two or more identical machines working in parallel; each job can be assigned to any machine
	Flow Shop	All jobs follow the same sequence of machines; each job visits every machine in the same order
	Job Shop	Each job has its own specific sequence of machines to follow


- Job Splitting
 - Non-preemptive problems

Once a job starts processing on a machine, it must continue without interruption until completion



- Preemptive problems

A job can be interrupted and resumed later, possibly on a different machine, without losing progress



- Performance Metrics

- **Makespan:** The sum of completion times of all jobs, possibly weighted by job importance
- **(Weighted) total completion time:** The sum of completion times of all jobs, possibly weighted by job importance (measures overall throughput)
- **(Weighted) number of delayed jobs:** The total number of jobs completed after they are due, possibly weighted by job priority
- **(Weighted) total lateness:** The sum of differences between each job's completion time and due date (can be negative or positive), possibly weighted
- **(Weighted) total tardiness:** The sum of positive delays (i.e., lateness when a job finishes after its due date), possibly weighted. Negative values (early completions) are treated as zero.
- ...

34.7.1. Single Machine Serial Production



Notation

- n : Total number of jobs
- $J = \{1, 2, \dots, n\}$: Set of jobs
- $j \in J$: A job
- p_j : processing time of job j

- x_j : completion time of job j
- $z_{ij} \in \{0, 1\}$: Binary variable

$$z_{ij} = \begin{cases} 1 & \text{if job } j \text{ is before job } i \\ 0 & \text{otherwise} \end{cases}$$

Job Order Constraints

We must ensure that either job i comes before job j or vice versa, but not both

If job i is before job j ($z_{ij} = 0$)

$$x_j \geq x_i + p_j$$

If job j is before job i ($z_{ij} = 1$):


$$x_i \geq x_j + p_i$$

These two cases can be rewritten using a big- M formulation:

$$\begin{aligned} x_i + p_j - x_j &\leq Mz_{ij} \\ x_j + p_i - x_i &\leq M(1 - z_{ij}) \end{aligned}$$

Where:

- $M = \sum_{j \in J} p_j$



Complete Formulation

$$\begin{aligned} \min \quad & \sum_{j \in J} x_j \\ \text{s.t.} \quad & x_i + p_j - x_j \leq Mz_{ij} & \forall i \in J, j \in J, i < j \\ & x_j + p_i - x_i \leq M(1 - z_{ij}) & \forall i \in J, j \in J, i < j \\ & x_j \geq p_j & \forall j \in J \\ & x_j \geq 0 & \forall j \in J \\ & z_{ij} \in \{0, 1\} & \forall i \in J, j \in J, i < j \end{aligned}$$

Interpretation

If jobs are run in order 1, 2, ..., n , the completion times are:

$$\begin{aligned} x_1 &= p_1 \\ x_2 &= p_1 + p_2 \\ x_3 &= p_1 + p_2 + p_3 \\ &\vdots \\ x_n &= \sum_{i=1}^n p_i \end{aligned}$$


So, the completion time of a job depends on the sum of processing times of all jobs before it in the schedule

Optional: Release Times

If jobs have release times R_j , meaning each job j cannot start before R_j , then we add:

$$x_j \geq R_j + p_j \quad \forall j \in J$$

34.7.2. Multiple Parallel Machines



We aim to assign jobs to machines such that:

- Each job is assigned to exactly one machine.
- A job can be processed on any machine.
- The objective is to minimize the makespan, which is the time when the last machine finishes processing.

Notation

- $J = \{1, 2, \dots, n\}$: set of jobs
- $I = \{1, 2, \dots, m\}$: set of machines
- p_j : processing time of job $j \in J$
- x_{ij} : a binary variable that indicates whether job j is assigned to machine i :

$$x_{ij} = \begin{cases} 1 & \text{if job } j \text{ is assigned to machine } i \\ 0 & \text{otherwise} \end{cases}$$

- The completion time of machine i is the total processing time of all jobs assigned to it

$$C_i = \sum_{j \in J} p_j x_{ij}$$

- The makespan w is the maximum completion time across all machines:

$$w \geq \sum_{j \in J} p_j x_{ij} \quad \forall i \in I$$

Objective

Minimize makespan w

Complete Formulation

$$\begin{aligned}
\min \quad & w \\
s.t. \quad & \sum_{j \in J} p_j x_{ij} \leq w \quad \forall i \in I \\
& \sum_{i \in I} x_{ij} = 1 \quad \forall j \in J \\
& x_{ij} \in \{0, 1\} \quad \forall i \in I, j \in J
\end{aligned}$$

- The first constraint ensures the makespan is at least as large as each machine's workload
- The second ensures each job is assigned to one and only one machine

```

from pyomo.environ import *

model = AbstractModel()

model.MACHINES = Set()
model.JOBS = Set()

model.p = Param(model.JOBS, within=PositiveReals)

model.x = Var(model.MACHINES, model.JOBS, domain=Binary)
model.w = Var(domain=NonNegativeReals)


def obj_rule(m):
    return m.w
model.Obj = Objective(rule=obj_rule, sense=minimize)

def job_assignment_rule(m, j):
    return sum(m.x[i, j] for i in m.MACHINES) == 1
model.JobAssignment = Constraint(model.JOBS, rule=job_assignment_rule)

def machine_completion_rule(m, i):
    return sum(m.p[j] * m.x[i, j] for j in m.JOBS) <= m.w
model.MachineCompletion = Constraint(model.MACHINES, rule=machine_completion_rule)

```

34.7.3. Flow Shop



34.7.4. Job Shop

34.8. Vehicle Routing

34.8.1. Traveling Salesperson

- Let $G = (V, E)$ be a directed complete graph
- Let d_{ij} be the distance (cost) from node i to j
- Let x_{ij} be a binary decision variable:

$$x_{ij} = \begin{cases} 1 & \text{if } (i, j) \text{ is selected} \\ 0 & \text{otherwise} \end{cases}$$

Objective

Minimize the total travel cost

$$\min \sum_{(i,j) \in E} d_{ij} x_{ij}$$

Constraints

1. Flow Balancing

For node $k \in V$

- One **incoming** edge:

$$\sum_{i \in V, i \neq k} = 1$$

- One **outgoing** edge:

$$\sum_{j \in V, j \neq k} = 1$$

2. Subtours

a. Miller-Tucker-Zemlin (MTZ)

- Let u_i be the position of node i in the tour (e.g., $u_i = k$ if node i is the k th node to be passed on the tour)

$$u_1 = 1$$

$$2 \leq u_i \leq n \quad \forall i \in V \setminus \{1\}$$

$$u_i - u_j + 1 \leq (n-1)(1 - x_{ij}) \quad \forall (i, j) \in E, i \neq 1, j \neq 1$$

b. Subtour Elimination Constraint (SEC)

For every subset $S \subset V$ with $|S| \geq 2$:

$$\sum_{\substack{i \in S, j \in S \\ i \neq j}} x_{ij} \leq |S| - 1 \quad \forall S \subset V, |S| \geq 2$$

This ensures that no subset of nodes forms a closed tour independent of the main tour.

Complete Formulation

$$\begin{aligned} \min \quad & \sum_{(i,j) \in E} d_{ij} x_{ij} \\ \text{s.t.} \quad & \sum_{i \in V, i \neq k} = 1 \quad \forall k \in V \\ & \sum_{j \in V, j \neq k} = 1 \quad \forall k \in V \\ & x_{ij} \in \{0, 1\} \quad \forall (i, j) \in E \\ & \text{MTZ or SEC} \end{aligned}$$

34.9. MIP (Mixed Integer Programming)

Optimizing (maximizing or minimizing) a linear **objective function** subject to linear equality or inequality **constraints**. **Decision variables** can take any continuous real or integer values

Minimize (or Maximize):

$$c^T x$$

Subject to:

- $Ax \leq b$
- $x_i \in \mathbb{Z}$ for some i
- $x_j \in \mathbb{R}$ for the remaining j

Where:

- x is the vector of decision variables
- c is the vector of objective function coefficients
- A is the constraint coefficient matrix
- b is the vector of constraint right-hand side values
- x_i represents integer variables
- x_j represents continuous variables

Objective Function

Maximize

$$c^T x$$
$$(3 \ 2 \ -5 \ -4) \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}$$

Constraints

$$Ax \leq b$$
$$\begin{pmatrix} 1 & 2 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ 1 & 0 & -5 & 0 \\ 0 & 1 & 0 & -5 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix} \leq \begin{pmatrix} 8 \\ 12 \\ 0 \\ 0 \end{pmatrix}$$

34.10. Non-Linear Programming

34.10.1. EOQ

Parameters


- D : annual demand (units)
- K : unit ordering cost (\$)
- h : unit holding cost per year (\$)
- p : unit purchasing cost (\$)

Decision Variable

- Q : order quantity per order (units)

Objective

Minimize annual total cost



- $\frac{q}{2}$: average inventory level
- $h \times \frac{q}{2} = \frac{hq}{2}$: annual holding cost
- pD : annual purchasing cost
- $\frac{D}{q}$: number of orders in a year
- $K \times \frac{D}{q} = \frac{KD}{q}$: annual ordering cost


Objective Function

$$\min_{q \geq 0} \underbrace{\frac{KD}{q}}_{\text{Ordering Cost}} + \underbrace{pD}_{\text{Purchasing Cost}} + \underbrace{\frac{hq}{2}}_{\text{Inventory Cost}}$$

pD is just a constant, so:

$$TC(q) = \underbrace{\frac{KD}{q}}_{\text{Ordering Cost}} + \underbrace{\frac{hq}{2}}_{\text{Inventory Cost}}$$

- As q **increases**:
 - Inventory costs increase
 - Ordering cost decrease
- As q **decreases**:
 - Inventory costs decrease
 - Ordering cost increase



34.10.2. Portfolio Optimization

Objective

Invest B dollars in n stocks while managing risk and meeting required return

Notation

- B : budget (\$)
- R : required minimum expected return
- For each stock $i \in \{1, \dots, n\}$
 - p_i : current price
 - μ_i : expected future price
 - σ_i^2 : variance of return per share
 - σ_{ij} : Covariance between assets i and j
 - x_i : number of shares to buy (decision variable)

Model

$$\begin{aligned}
 \min \quad & \sum_{i=1}^n \sigma_i^2 x_i^2 + 2 \sum_{i=1}^n \sum_{j=i+1}^n \sigma_{ij} x_i x_j && \text{(Total risk)} \\
 \text{s.t.} \quad & \sum_{i=1}^n p_i x_i \leq B && \text{(Budget constraint)} \\
 & \sum_{i=1}^n \mu_i x_i \geq R && \text{(Return constraint)} \\
 & x_i \geq 0 \quad \forall i = 1, \dots, n && \text{(No short-selling)}
 \end{aligned}$$

34.11. Linearizing Maximum/Minimum Functions

1. When the **maximum** function is on the **smaller side** of inequality

General Form

$$y \geq \max(x_1, x_2) \quad \Longleftrightarrow \quad y \geq x_1 \wedge y \geq x_2$$

- This rule holds regardless of whether x_1, x_2, y are variables, constants, or expressions
- You're ensuring that y is greater than or equal to all values in the maximum function, so it must be greater than or equal to each term individually

Generalized form with more than two elements

$$y \geq \max(x_1, x_2, \dots, x_n) \iff y \geq x_i \quad \forall i = 1, \dots, n$$

Example

$$y + x_1 + 3 \geq \max(x_1 - x_3, 2x_2 + 4) \iff \begin{cases} y + x_1 + 3 \geq x_1 - x_3 \\ y + x_1 + 3 \geq 2x_2 + 4 \end{cases}$$

2. When the **minimum** function is on the **larger side** of inequality

General Form

$$y \leq \min(x_1, x_2) \iff y \leq x_1 \wedge y \leq x_2$$

- This rule holds regardless of whether x_1, x_2, y are **variables**, **constants**, or **expressions**
- You're ensuring that y is smaller than or equal to all values in the minimum function, so it must be smaller than or equal to each term individually

Generalized form with more than two elements

$$y \leq \min(x_1, x_2, \dots, x_n) \iff y \leq x_i \quad \forall i = 1, \dots, n$$

Example

$$y + x_1 \leq \min(x_1 - x_3, 2x_2 + 4, 0) \iff \begin{cases} y + x_1 \leq x_1 - x_3 \\ y + x_1 \leq 2x_2 + 4 \\ y + x_1 \leq 0 \end{cases}$$

Cases Where Linearization Does Not Apply

- $y \leq \max(x_1, x_2) \not\iff y \leq x_1 \wedge y \leq x_2$
- $y \geq \max(x_1, x_2) \not\iff y \geq x_1 \wedge y \geq x_2$
- Maximum or minimum function in an equality

$$y = \max(x_1, x_2) \quad \text{or} \quad y = \min(x_1, x_2)$$

34.12. Linearize Objective Function

1. Minimize a Maximum Function

$$\min \max(x_1, x_2) \iff \begin{array}{ll} \min & w \\ \text{s.t.} & w \geq x_1 \\ & w \geq x_2 \end{array}$$

2. Maximize a Minimum Function

$$\begin{array}{ll} \max & \min(x_1, x_2, 2x_3 + 5) + x_4 \\ \text{s.t.} & 2x_1 + x_2 - x_4 \leq x_3 \end{array} \iff \begin{array}{ll} \max & w + x_4 \\ \text{s.t.} & w \leq x_1 \\ & w \leq x_2 \\ & w \leq 2x_3 + 5 \\ & 2x_1 + x_2 - x_4 \leq x_3 \end{array}$$

Cases Where Linearization Does Not Apply

- $\max \max(x_1, x_2)$
- $\min \min(x_1, x_2)$

Absolute Function

The absolute value is equivalent to a maximum function:

$$|x| = \max(x, -x)$$

Thus, it can be linearized when it appears on the smaller side of an inequality:

$$|x| \leq y \iff \begin{cases} x \leq y \\ -x \leq y \end{cases}$$


We want to allocate \$1000 to 2 people in a fair way

- **Fairness criterion:** Minimize the difference between the amounts each person receives.
- Let x_i be amount to allocate to person i for $i = 1, 2$

We write the problem as:

$$\begin{aligned} \min \quad & |x_2 - x_1| \\ \text{s.t.} \quad & x_1 + x_2 = 1000 \\ & x_i \geq 0 \quad \forall i = 1, 2 \end{aligned}$$

- The absolute value makes the objective **nonlinear**



Linearizing the Problem

We now reformulate the problem to make it linear

Step 1: Introduce a new variable


Let w be the absolute difference: $w = |x_2 - x_1|$

$$\begin{aligned} \min \quad & w \\ \text{s.t.} \quad & x_1 + x_2 = 1000 \\ & w = |x_2 - x_1| \\ & x_i \geq 0 \quad \forall i = 1, 2 \end{aligned}$$

Step 2: Relax the equality

We replace the equality with an inequality:


$$\begin{aligned} \min \quad & w \\ \text{s.t.} \quad & x_1 + x_2 = 1000 \\ & w \geq |x_2 - x_1| \\ & x_i \geq 0 \quad \forall i = 1, 2 \end{aligned}$$



Step 3. Replace absolute value with maximum

Recall:

$$|x_2 - x_1| = \max(x_2 - x_1, x_1 - x_2)$$



Therefore:

$$w \geq |x_2 - x_1| \quad \Leftrightarrow \quad w \geq \max(x_2 - x_1, x_1 - x_2) \quad \Leftrightarrow \quad w \geq x_2 - x_1 \wedge w \geq x_1 - x_2$$


Now the problem is fully linear:

$$\begin{aligned} \min \quad & w \\ \text{s.t.} \quad & x_1 + x_2 = 1000 \\ & w \geq x_2 - x_1 \\ & w \geq x_1 - x_2 \\ & x_i \geq 0 \quad \forall i = 1, 2 \end{aligned}$$

Reformulating in One Variable

using $x_2 = 1000 - x_1$, we can express everything in terms of x_1 :

$$\begin{aligned} \min \quad & w \\ \text{s.t.} \quad & w \geq 1000 - 2x_1 \\ & w \geq 2x_1 - 1000 \\ & x_1 \geq 0 \end{aligned}$$



We are minimizing w , which represents the difference between the two allocations. The solution occurs when the two inequalities intersect — that is, when both are equal:

$$1000 - 2x_1 = 2x_1 - 1000 \Rightarrow x_1 = 500$$

So:

- $x_1 = 500$
- $x_2 = 1000 - x_1 = 500$
- $w = |500 - 500| = 0$

This is the fair allocation: each person receives \$500, and the difference between the two amounts is 0.

34.13. Linearizing Products of Decision Variables

Products of decision variables can be linearized if:

- A binary and a continuous variable
- Two binary variables

Cannot be linearized if:

- Two continuous variables

34.14. Simplex Method


34.14.1. Linear Algebra Review

$$x + y = 5$$

$$2x - y = 4$$

Row View


Solution: Intersection of n lines or (hyper)planes



Column View

Solution: Combination of LHS column vectors to form the RHS vector

$$x \begin{bmatrix} 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 5 \\ 4 \end{bmatrix}$$



A linear system is **singular** if there is no unique solution

- Row view: the n (hyper)planes do not intersect at exactly one point
- Column view: the n vectors do not span a complete n -dimensional space

34.14.2. Gaussian Elimination

Given a system $Ax = b$, or augmented matrix $[A | b]$, use the following 3 rules:

- Row Swapping: $R_i \leftrightarrow R_j$
- Row Scaling: $k \cdot R_i \rightarrow R_i$
- Row Replacement: $R_i - k \cdot R_j \rightarrow R_i$

To solve the system:

1. Transform the augmented matrix into row echelon form (REF) using these operations:
 - The first nonzero entry (pivot) in each row is to the right of the pivot in the row above
 - All entries below a pivot are zero
 - Rows of all zeros (if any) are at the bottom
2. Back-substitute starting from the last nonzero row to find the solution

Non-singular Case

System of equations

$$\begin{aligned}2y &= 4 \\ 1x + 1y &= 3\end{aligned}$$

Augmented matrix

$$\left[\begin{array}{cc|c} 0 & 2 & 4 \\ 1 & 1 & 3 \end{array} \right]$$

Swapping ($R_1 \leftrightarrow R_2$)

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 0 & 2 & 4 \end{array} \right]$$

Scaling ($\frac{1}{2}R_2 \rightarrow R_2$)

Row Echelon Form (REF)

$$\left[\begin{array}{cc|c} 1 & 1 & 3 \\ 0 & 1 & 2 \end{array} \right]$$

Replacement ($R_1 - R_2 \rightarrow R_1$)

Reduced Row Echelon Form (RREF)

$$\left[\begin{array}{cc|c} 1 & 0 & 1 \\ 0 & 1 & 2 \end{array} \right]$$

Result

$$x = 1, \quad y = 2$$

Singular Case

System of equations

$$\begin{aligned}x + y &= 2 \\ 2x + 2y &= 4\end{aligned}$$

Augmented matrix

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 4 \end{array} \right]$$

Replacement ($R_2 - 2R_1 \rightarrow R_2$)

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 0 & 0 & 0 \end{array} \right]$$

Infinitely many solutions E.g.

- $x = 0, \quad y = 2$
- $x = 1, \quad y = 1$
- $x = 2, \quad y = 0$
- etc.

Rows are **linearly dependent**

The coefficient matrix

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$$

A has **determinant** 0, so it's **not invertible**

34.14.3. Inverse

$$A^{-1}A = I \quad \wedge \quad AA^{-1} = I$$

A^{-1} is **unique**

Finding A^{-1} : Gauss-Jordan Elimination

A square matrix is **nonsingular** iff it is **invertible**

$$[A | I] \longrightarrow [I | A^{-1}]$$

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

$$[A | I] = \left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 2 & 1 & 0 & 1 \end{array} \right]$$

Replacement ($R_2 - 2R_1 \longrightarrow R_2$)

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & -3 & -2 & 1 \end{array} \right]$$

Scale ($-\frac{1}{3}R_2 \longrightarrow R_2$)

$$\left[\begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & 1 & \frac{2}{3} & -\frac{1}{3} \end{array} \right]$$

Replacement ($R_1 - 2R_2 \longrightarrow R_1$)

$$\left[\begin{array}{cc|cc} 1 & 0 & -\frac{1}{3} & \frac{2}{3} \\ 0 & 1 & \frac{2}{3} & -\frac{1}{3} \end{array} \right]$$

Result

$$A^{-1} = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} \end{bmatrix}$$

34.14.4. Linear Dependence and Independence

A set of m n -dimensional vectors x_1, x_2, \dots, x_m are **linearly dependent** if there exists a non-zero vector $w \in \mathbb{R}^m$ such that:

$$w_1x_1 + w_2x_2 + \dots + w_mx_m = 0$$

That is, at least one of the vectors can be written as a linear combination of the others.

Dependent

Let:

$$x_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

Augmented:

$$\left[\begin{array}{cc|c} 1 & 2 & 0 \\ 2 & 4 & 0 \end{array} \right]$$

Elimination ($R_2 - 2R_1 \rightarrow R_2$)

$$\left[\begin{array}{cc|c} 1 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Independent

Let:

$$x_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad x_2 = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

Augmented:

$$\left[\begin{array}{cc|c} 1 & 3 & 0 \\ 2 & 1 & 0 \end{array} \right]$$

Elimination ($R_2 - 2R_1 \rightarrow R_2$)

$$\left[\begin{array}{cc|c} 1 & 2 & 0 \\ 0 & -5 & 0 \end{array} \right]$$

Back-substitute

- $-5w_2 = 0 \Rightarrow w_2 = 0$
- $w_1 + 3w_2 = 0 \Rightarrow w_1 = 0$

Only solution is $w_1 = w_2 = 0$




In Gaussian elimination, a **row of all zeros**

1. **Homogeneous** System (RHS = 0)

$$\left[\begin{array}{cc|c} 1 & 2 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

$$x_1 + 2x_2 = 0$$

$$0 = 0$$



-  Represents a **redundant** equation (i.e., no new information) **dependent** set of vectors
-  Does not affect consistency: the system is always **consistent**
-  The system has fewer pivots than variables, hence **infinitely many solutions**

2. **Inconsistent** System (Nonzero RHS)

$$\left[\begin{array}{cc|c} 1 & 2 & 3 \\ 0 & 0 & 5 \end{array} \right]$$

$$x_1 + 2x_2 = 3$$

$$0 = 5$$

-  The second row leads to a **contradiction**
-  The system is **inconsistent** (no solution)

34.14.5. Extreme Points

Given that x , x_1 , and x_2 are points in the set S :


- If you can find two different points x_1 and x_2 in S such that x lies somewhere strictly between them on the straight line connecting x_1 and x_2 , then x is not an extreme point
- If no such pair of points exists—meaning you cannot place x anywhere strictly between two other points in S on a straight line—then x is an extreme point

Formal Definition

For a set $S \subseteq \mathbb{R}^n$, a point x is an extreme point if there does not exist a three-tuple (x_1, x_2, λ) such that $x_1 \in S$, $x_2 \in S$, $\lambda \in (0, 1)$, and

$$x = \lambda x_1 + (1 - \lambda)x_2$$

A point x in set $S \subseteq \mathbb{R}^n$ is extreme if it cannot be written as a strict **convex combination** of two different points $x_1, x_2 \in S$.



Convex Combination

Let $x_1, x_2, \dots, x_k \in \mathbb{R}^n$. A convex combination of these vectors is any point of the form:

$$x = \lambda_1 x_1 + \lambda_2 x_2 + \dots + \lambda_k x_k$$

where:

- $\lambda_i \geq 0 \quad \forall i$
- $\sum_{i=1}^k \lambda_i = 1$

x lies strictly between x_1 and x_2 on the line segment connecting them, not equal to either one

34.14.6. Slack and Suplus

1. Slack

Unused capacity


Given a constraint:

$$x_1 + x_2 \leq 5$$

The total **must not exceed** 5. To convert this into an equation, we **add** a slack variable $s \geq 0$:

$$x_1 + x_2 + s = 5$$

- If $x_1 + x_2 = 3$, then $s = 2$
- If $x_1 + x_2 = 5$, then $s = 0$



2. Excess (surplus)

Excess above the required amount


Given a constraint:

$$x_1 + x_2 \geq 5$$

The total **must be at least** 5. To convert this into an equation, we **subtract** a excess variable $s \geq 5$:

$$x_1 + x_2 - e = 5$$

- If $x_1 + x_2 = 7$, then $e = 2$
- If $x_1 + x_2 = 5$, then $e = 0$



Objective Function

Max	✓	$\max z = c^T x$
-----	---	------------------

Min \rightarrow Max	$-z$	$\min z = c^T x$ \Leftrightarrow $\max z = -c^T x$
Constraint Forms		
\leq	Slack	$a_1 x_1 + a_2 x_2 \leq b$ $a_1 x_1 + a_2 x_2 + s = b$
\geq	Excess + Artificial	$a_1 x_1 + a_2 x_2 \geq b$ $a_1 x_1 + a_2 x_2 - e + a = b$
$=$	Artificial	$a_1 x_1 + a_2 x_2 = b$ $a_1 x_1 + a_2 x_2 + a = b$
Negative b	Multiply by -1	$a_1 x_1 + a_2 x_2 \leq -b$ $-a_1 x_1 - a_2 x_2 \geq b$
Variable Bounds		
Non-negativity		$x_i, s_i, a_i \geq 0 \quad \forall i$
Lower Bound $\neq 0$	Substitution	$x_1 \geq 5$ $y_1 = x_1 - 5$ $x_1 = y_1 + 5$ $y_1 \geq 0$
Unrestricted Variables	Replace with difference	$x_j \text{ unrestricted}$ $x_j = x_j' - x_j''$ $x_j', x_j'' \geq 0$
Variable Roles		
Basic Variables	Usually non-zero	Part of the current solution (solved from the constraints)
Non-Basic Variables	Always zero	Temporarily set to 0 so basic variables can be solved
Simplex		

<p>Selection:</p> <ul style="list-style-type: none"> • Entering variable • Pivot column selection 	<p>For maximization problems:</p> <ul style="list-style-type: none"> • Select the column with the most negative value in the objective row • This corresponds to the variable that will increase the objective function the most <p>For minimization problems:</p> <ul style="list-style-type: none"> • Select the column with the most positive value in the objective row • This corresponds to the variable that will decrease the objective function the most 	
<p>Selection:</p> <ul style="list-style-type: none"> • Leaving variable • Pivot row 	<ol style="list-style-type: none"> 1. Calculate ratios for each row (except the objective row) <ul style="list-style-type: none"> • For each row: b_i/p_i • Only consider rows where the pivot column element $p_i > 0$ • Ignore rows where the pivot column element $p_i \leq 0$ 2. Select the minimum ratio: <ul style="list-style-type: none"> • The row with the smallest non-negative ratio becomes the pivot row • The basic variable in this row is the leaving variable (exits the basis) 	

,

$$\begin{aligned} \max \quad & z = 3x_1 + 2x_2 \\ \text{s.t.} \quad & x_1 + x_2 \leq 4 \\ & x_1 \leq 2 \\ & x_2 \leq 3 \\ & x_1, x_2 \geq 0 \end{aligned}$$

Step 1: Convert to Standard Form

$$\begin{aligned} \max \quad & z = 3x_1 + 2x_2 \\ \text{s.t.} \quad & x_1 + x_2 + s_1 = 4 \\ & x_1 + s_2 = 2 \\ & x_2 + s_3 = 3 \\ & x_1, x_2, s_1, s_2, s_3 \geq 0 \end{aligned}$$


Objective

$$z - 3x_1 - 2x_2 = 0$$

Step 2: Initial Tableau

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	4
s_2	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	0	0	0

- Basic Variables: $s_1 = 4, s_2 = 2, s_3 = 3$
- Non-basic variables: $x_1 = 0, x_2 = 0$
 - $V_0 = (0, 0)$
 - Objective: $z = 3x_1 + 2x_2 = 3(0) + 2(0) = 0$



Step 3: Choose entering variable

- Look at the bottom row (z):
 - Coefficients of $x_1 = -3$, $x_2 = -2$
- Most negative: $x_1 \rightarrow$ Enter the basis

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	4
s_2	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	0	0	0

Step 4: Determine leaving variable

Look at ratios of b/x_1 :

Basis	x_1	x_2	s_1	s_2	s_3	b	b/x_1
s_1	1	1	1	0	0	4	$4/1 = 4$
s_2	1	0	0	1	0	2	$2/1 = 2$
s_3	0	1	0	0	1	3	$3/0 = \infty$
z	-3	-2	0	0	0	0	

- Minimum Ratio
 - s_2 leaves, x_1 enters

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	4
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3

z	-3	-2	0	3	0	0
-----	------	------	-----	-----	-----	-----

Step 5: Pivot on row 2, column x_1

Now eliminate x_1 from other rows:

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	4
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	3	0	0

$$R_{s_1} := R_{s_1} - R_{x_1}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	4
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	3	0	0

$$R_{s_1} := R_{s_1} - R_{x_1} \implies \begin{array}{lcl} x_1 : & 1 - 1 = & 0 \\ x_2 : & 1 - 0 = & 1 \\ s_1 : & 1 - 0 = & 1 \\ s_2 : & 0 - 1 = & -1 \\ s_3 : & 0 - 0 = & 0 \\ b : & 4 - 2 = & 2 \end{array}$$

Update R_{s_1}

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	3	0	0

$$R_z := R_z - 3R_{x_1}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	-3	-2	0	3	0	0


$$R_z := R_z - 3R_{x_1} \implies \begin{array}{lcl} x_1 : & -3 + 3(1) & = 0 \\ x_2 : & -2 + 3(0) & = -2 \\ s_1 : & 0 + 3(0) & = 0 \\ s_2 : & 0 + 3(1) & = 3 \\ s_3 : & 0 + 3(0) & = 0 \\ b : & 0 + 3(2) & = 6 \end{array}$$

Update R_z

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

- Basic Variables: $x_1 = 2, s_1 = 2, s_3 = 3$
- Non-basic variables: $x_2 = 0$
 - $V_1 = (2, 0)$
 - Objective: $z = 3x_1 + 2x_2 = 3(2) + 2(0) = 6$

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6



Step 3: Choose entering variable

- Look at the bottom row (z):
 - Coefficients of $x_2 = -2$

- Most negative: $x_2 \rightarrow$ Enter the basis

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

Step 4: Determine leaving variable

Look at ratios of b/x_2 :

Basis	x_1	x_2	s_1	s_2	s_3	b	b/x_2
s_1	0	1	1	-1	0	2	$2/1 = 2$
x_1	1	0	0	1	0	2	$2/0 = \infty$
s_3	0	1	0	0	1	3	$3/1 = 3$
z	0	-2	0	3	0	6	

- s_1 leaves, x_2 enters

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

Step 5: Pivot on row 1, column x_2

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

Now eliminate x_2 from other rows:

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

$$R_{s_3} := R_{s_3} - R_{x_2}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	1	0	0	1	3
z	0	-2	0	3	0	6

$$R_{s_3} := R_{s_3} - R_{x_2} \implies \begin{array}{l} x_1 : 0 - 0 = 0 \\ x_2 : 1 - 1 = 0 \\ s_1 : 0 - 1 = -1 \\ s_2 : 0 - (-1) = 1 \\ s_3 : 1 - 0 = 1 \\ b : 3 - 2 = 1 \end{array}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	0	-1	1	1	1
z	0	-2	0	3	0	6


$$R_z = R_z - (-2)R_{x_2} = R_z + 2R_{x_2}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	0	-1	1	1	1
z	0	-2	0	3	0	6

$$R_z = R_z + 2R_{x_2} \implies \begin{array}{l} x_1 : 0 + 2(0) = 0 \\ x_2 : -2 + 2(1) = 0 \\ s_1 : 0 + 2(1) = 2 \\ s_2 : 3 + 2(-1) = 1 \\ s_3 : 0 + 2(0) = 0 \\ b : 6 + 2(2) = 10 \end{array}$$

Basis	x_1	x_2	s_1	s_2	s_3	b
x_2	0	1	1	-1	0	2
x_1	1	0	0	1	0	2
s_3	0	0	-1	1	1	1
z	0	0	2	1	0	10

- Basic Variables: $x_1 = 2, x_2 = 2, s_3 = 1$
- Non-basic variables: $s_1 = 0, s_2 = 0$
 - $V_2 = (2, 2)$
 - Objective: $z = 3x_1 + 2x_2 = 3(2) + 2(2) = 10$



Problem

$$\begin{array}{ll}
 \min & z = x_1 + 2x_2 \\
 \text{s.t.} & x_1 + x_2 \leq 1 \quad (\text{slack}) \\
 & 2x_1 - x_2 \geq 2 \quad (\text{surplus} + \text{artificial}) \\
 & -x_1 + 3x_2 \leq -1 \quad (\text{negative } b) \\
 & x_1 - x_2 = 1 \quad (\text{artificial}) \\
 & x_1 \geq 0 \\
 & x_2 \quad (\text{unrestricted})
 \end{array}$$

Standard Form

Step 1: Handle Unrestricted Variable x_2

Since x_2 is unrestricted:

$$\begin{aligned}
 x_2 &= x'_2 - x''_2 \\
 x'_2, x''_2 &\geq 0
 \end{aligned}$$

Rewrite the objective:

$$\begin{aligned}
 z &= x_1 + 2(x'_2 - x''_2) \\
 &= x_1 + 2x'_2 - 2x''_2
 \end{aligned}$$

Step 2: Slack Constraint (\leq)

$$x_1 + x_2 \leq 1$$

Rewrite with $x'_2 - x''_2$:

$$x_1 + (x'_2 - x''_2) \leq 1$$

Add slack variable:

$$\begin{aligned} x_1 + x'_2 - x''_2 + s_1 &= 1 \\ s_1 &\geq 0 \end{aligned}$$

Step 3: Surplus + Artificial Constraint (\geq)

$$2x_1 - x_2 \geq 2$$

Rewrite with $x'_2 - x''_2$:

$$2x_1 - (x'_2 - x''_2) \geq 2 \implies 2x_1 - x'_2 + x''_2 \geq 2$$

Convert to equality by subtracting a surplus variable and adding an artificial variable:

$$\begin{aligned} 2x_1 - x'_2 + x''_2 - s_2 + a_1 &= 2 \\ s_2, a_1 &\geq 0 \end{aligned}$$

Step 4: Negative RHS Constraint

$$-x_1 + 3x_2 \leq -1$$

Rewrite with $x'_2 - x''_2$:

$$-x_1 + 3(x'_2 - x''_2) \leq -1 \implies -x_1 + 3x'_2 - 3x''_2 \leq -1$$

Multiply both sides by -1 to get a positive RHS and flip inequality:

$$x_1 - 3x'_2 + 3x''_2 \geq 1$$

Convert to equality by subtracting a surplus variable and adding an artificial variable:

$$\begin{aligned} x_1 - 3x'_2 + 3x''_2 - s_3 + a_2 &= 1 \\ s_3, a_2 &\geq 0 \end{aligned}$$

Step 5: Artificial Equality Constraint ($=$)

$$x_1 - x_2 = 1$$

Rewrite with $x'_2 - x''_2$:

$$x_1 - (x'_2 - x''_2) = 1 \implies x_1 - x'_2 + x''_2 = 1$$

Add an artificial variable:

$$\begin{aligned} x_1 - x'_2 + x''_2 + a_3 &= 1 \\ a_3 &\geq 0 \end{aligned}$$

Final Standard Form

$$\begin{aligned}
\min \quad & z = x_1 + 2x'_2 - 2x''_2 \\
s.t. \quad & x_1 + x'_2 - x''_2 + s_1 = 1 \\
& 2x_1 - x'_2 + x''_2 - s_2 + a_1 = 2 \\
& x_1 - 3x'_2 + 3x''_2 - s_3 + a_2 = 1 \\
& x_1 - x'_2 + x''_2 + a_3 = 1 \\
& x_1, x'_2, x''_2, s_1, s_2, s_3, a_1, a_2, a_3 \geq 0
\end{aligned}$$

Phase I

Objective

$$\begin{aligned}
\min \quad & w = a_1 + a_2 + a_3 \\
s.t. \quad & x_1 + x'_2 - x''_2 + s_1 = 1 && \text{(slack)} \\
& 2x_1 - x'_2 + x''_2 - s_2 + a_1 = 2 && \text{(surplus + artificial)} \\
& x_1 - 3x'_2 + 3x''_2 - s_3 + a_2 = 1 && \text{(negative } b) \\
& x_1 - x'_2 + x''_2 + a_3 = 1 && \text{(artificial)} \\
& x_1, x'_2, x''_2, s_1, s_2, s_3, a_1, a_2, a_3 \geq 0 && \text{(unrestricted)}
\end{aligned}$$

Deriving Phase I Objective Row R_w

- Objective function before substitution

$$w = a_1 + a_2 + a_3$$

- Express each artificial variable in terms of other variables from their constraints

- R_{a_1}

$$\begin{aligned}
2x_1 - x'_2 + x''_2 - s_2 + a_1 &= 2 \\
a_1 &= -2x_1 + x'_2 - x''_2 + s_2 + 2
\end{aligned}$$

- R_{a_2}

$$\begin{aligned}
x_1 - 3x'_2 + 3x''_2 - s_3 + a_2 &= 1 \\
a_2 &= -x_1 + 3x'_2 - 3x''_2 + s_3 + 1
\end{aligned}$$

- R_{a_3}

$$\begin{aligned}
x_1 - x'_2 + x''_2 + a_3 &= 1 \\
a_3 &= -x_1 + x'_2 - x''_2 + 1
\end{aligned}$$

- Substitute into w :

$$\begin{aligned}
w &= a_1 + a_2 + a_3 \\
&= (-2x_1 + x'_2 - x''_2 + s_2 + 2) + (-x_1 + 3x'_2 - 3x''_2 + s_3 + 1) + (-x_1 + x'_2 - x''_2 + 1) \\
&= (-2x_1 - x_1 - x_1) + (x'_2 + 3x'_2 + x'_2) + (-x''_2 - 3x''_2 - x''_2) + (s_2 + s_3) + (2 + 1 + 1) \\
&= -4x_1 + 5x'_2 - 5x''_2 + s_2 + s_3 + 4
\end{aligned}$$

- Rewrite as a constraint (bring all variables to the left):

$$w + 4x_1 - 5x'_2 + 5x''_2 - s_2 - s_3 = 4$$

$$w = -4x_1 + 5x'_2 - 5x''_2 + s_2 + s_3 + 4$$

So in tableau row R_w :

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	1
a_1	2	-1	1	0	-1	0	1	0	0	2
a_2	1	-3	3	0	0	-1	0	1	0	1
a_3	1	-1	1	0	0	0	0	0	1	1
w	4	-5	5	0	-1	-1	0	0	0	4

Perform First Pivot

1. Pivot Column

Look at the w row (objective row) for the most negative coefficient:

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	1
a_1	2	-1	1	0	-1	0	1	0	0	2
a_2	1	-3	3	0	0	-1	0	1	0	1
a_3	1	-1	1	0	0	0	0	0	1	1
w	4	-5	5	0	-1	-1	0	0	0	4

The most negative is $-5 \rightarrow$ pivot column is x'_2

2. Ratio

- Compute ratios (b/x'_2) for rows with positive pivot entries:

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b	b/x'_2
s_1	1	1	-1	1	0	0	0	0	0	1	1
a_1	2	-1	1	0	-1	0	1	0	0	2	×
a_2	1	-3	3	0	0	-1	0	1	0	1	×
a_3	1	-1	1	0	0	0	0	0	1	1	×
w	4	-5	5	0	-1	-1	0	0	0	4	

- Minimum ratio is 1, so we pivot on row s_1 :

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b	b/x'_2
s_1	1	1	-1	1	0	0	0	0	0	1	1
a_1	2	-1	1	0	-1	0	1	0	0	2	×
a_2	1	-3	3	0	0	-1	0	1	0	1	×

a_3	1	-1	1	0	0	0	0	0	1	1	✗
w	4	-5	5	0	-1	-1	0	0	0	4	

3. Pivot $s_1 \rightarrow x'_2$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
x'_2	1	1	-1	1	0	0	0	0	0	1
a_1	2	-1	1	0	-1	0	1	0	0	2
a_2	1	-3	3	0	0	-1	0	1	0	1
a_3	1	-1	1	0	0	0	0	0	1	1
w	4	-5	5	0	-1	-1	0	0	0	4

4. Gauss-Jordan Elimination

Step 3: Pivot $a_1 \rightarrow x_1$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	5
x_1	3	-1	1	0	-1	0	1	0	0	4
a_2	1	-2	2	0	0	-1	0	1	0	6
a_3	1	-1	1	0	0	0	0	0	1	2
w	-5	4	-4	0	1	1	0	0	0	12

Gauss-Jordan Elimination

$$R_{x_1} = \frac{1}{3}R_{x_1}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	5
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	1	-2	2	0	0	-1	0	1	0	6
a_3	1	-1	1	0	0	0	0	0	1	2
w	-5	4	-4	0	1	1	0	0	0	12

$$R_{a_2} = R_{a_2} - 1R_{x_1}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	5
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
a_3	1	-1	1	0	0	0	0	0	1	2

w	-5	4	-4	0	1	1	0	0	0	12
-----	----	---	----	---	---	---	---	---	---	----

$$R_{a_3} = R_{a_3} - 1R_{x_1}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	5
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
a_3	0	-2/3	2/3	0	1/3	0	-1/3	0	1	2/3
w	-5	4	-4	0	1	1	0	0	0	12

$$R_w = R_w + 5R_{x_1}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	1	1	-1	1	0	0	0	0	0	5
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
a_3	0	-2/3	2/3	0	1/3	0	0-1/3	0	1	2/3
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

$$R_{s_1} = R_{s_1} - 1R_{x_1}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
a_3	0	-2/3	2/3	0	1/3	0	-1/3	0	1	2/3
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

Step 3: Perform the Second Pivot

1. Look at the w row (objective row) for the most negative coefficient:

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
a_3	0	-2/3	2/3	0	1/3	0	0-1/3	0	1	2/3
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

- The most negative is $-\frac{7}{3}$ for x''_2
 - Pivot column is x''_2

2. Ratio

- Compute ratios (b/x_2'') for rows with positive pivot entries:

Basis	x_1	x_2'	x_2''	s_1	s_2	s_3	a_1	a_2	a_3	b	b/x_2''
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3	×
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3	$\frac{4}{3} \div \frac{1}{3} = 4$
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3	$\frac{14}{3} \div \frac{5}{3} = 2.8$
a_3	0	-2/3	2/3	0	1/3	0	0-1/3	0	1	2/3	$\frac{2}{3} \div \frac{2}{3} = 1$
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3	

- Minimum ratio is 1, so we pivot on row a_3 :

Basis	x_1	x_2'	x_2''	s_1	s_2	s_3	a_1	a_2	a_3	b	b/x_2''
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3	×
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3	$\frac{4}{3} \div \frac{1}{3} = 4$
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3	$\frac{14}{3} \div \frac{5}{3} = 2.8$
a_3	0	-2/3	2/3	0	1/3	0	0-1/3	0	1	2/3	$\frac{2}{3} \div \frac{2}{3} = 1$
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3	

Step 4: Pivot $a_3 \rightarrow x_2''$

Basis	x_1	x_2'	x_2''	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
x_2''	0	-2/3	2/3	0	1/3	0	-1/3	0	1	2/3
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

Gauss-Jordan Elimination

$$R_{x_2''} = R_{x_2''} / (2/3)$$

Basis	x_1	x_2'	x_2''	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	4/3	-4/3	1	1/3	0	-1/3	0	0	11/3
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
x_2''	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

$$R_{s_1} = R_{s_1} - \left(-\frac{4}{3}\right)R_{x_2''} = R_{s_1} + \frac{4}{3}R_{x_2''}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	0	0	1	1	0	-1	0	2	5
x_1	1	-1/3	1/3	0	-1/3	0	1/3	0	0	4/3
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
x''_2	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

$$R_{x_1} = R_{x_1} - \frac{1}{3}R_{x''_2}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	0	0	0	-1/2	0	1/2	0	-1/2	1
x_1	1	0	0	0	-1/2	0	1/2	0	-1/2	1
a_2	0	-5/3	5/3	0	1/3	-1	-1/3	1	0	14/3
x''_2	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

$$R_{a_2} = R_{a_2} - \frac{5}{3}R_{x''_2}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	0	0	0	-1/2	0	1/2	0	-1/2	1
x_1	1	0	0	0	-1/2	0	1/2	0	-1/2	1
a_2	0	0	0	0	-1/2	-1	1/2	1	-5/2	3
x''_2	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	7/3	-7/3	0	-2/3	1	5/3	0	0	56/3

$$R_w = R_w - \left(-\frac{7}{3}\right)R_{x''_2} = R_w + \frac{7}{3}R_{x''_2}$$

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
s_1	0	0	0	0	-1/2	0	1/2	0	-1/2	1
x_1	1	0	0	0	-1/2	0	1/2	0	-1/2	1
a_2	0	0	0	0	-1/2	-1	1/2	1	-5/2	3
x''_2	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	0	0	0	1/2	1	1/2	0	7/2	21

Step 5: Perform the Third Pivot

1. Look at the w row (objective row) for the most negative coefficient:

Basis	x_1	x'_2	x''_2	s_1	s_2	s_3	a_1	a_2	a_3	b
-------	-------	--------	---------	-------	-------	-------	-------	-------	-------	-----

s_1	0	0	0	0	-1/2	0	1/2	0	-1/2	1
x_1	1	0	0	0	-1/2	0	1/2	0	-1/2	1
a_2	0	0	0	0	-1/2	-1	1/2	1	-5/2	3
x_2''	0	-1	1	0	1/2	0	-1/2	0	3/2	1
w	0	0	0	0	-1/6	1	1/2	0	7/2	21

- The most negative is $-\frac{1}{6}$ for s_2
 - Pivot column is s_2

2. Ratio

- Compute ratios (b/s_2) for rows with positive pivot entries:


Basis	x_1	x_2'	x_2''	s_1	s_2	s_3	a_1	a_2	a_3	b	b/s_2
s_1	0	0	0	0	-1/2	0	1/2	0	-1/2	1	×
x_1	1	0	0	0	-1/2	0	1/2	0	-1/2	1	×
a_2	0	0	0	0	-1/2	-1	1/2	1	-5/2	3	×
x_2''	0	-1	1	0	1/2	0	-1/2	0	3/2	1	2
w	0	0	0	0	-1/6	1	1/2	0	7/2	21	

Two-Phase

$$\begin{aligned}
 \max \quad & z = 2x_1 + x_2 \\
 s.t. \quad & x_1 + x_2 \geq 2 \\
 & x_1 \geq 1 \\
 & x_1 + x_2 \leq 5 \\
 & x_1, x_2 \geq 0
 \end{aligned}$$

$$\begin{aligned}
 \max \quad & z = 2x_1 + x_2 + 0e_1 + 0e_2 + 0s_1 \\
 s.t. \quad & x_1 + x_2 - e_1 \geq 2 \\
 & x_1 - e_2 \geq 1 \\
 & x_1 + x_2 + s_1 \leq 5 \\
 & z - 2x_1 - x_2 = 0
 \end{aligned}$$

Basis	x_1	x_2	e_1	e_2	s_1	b
???	1	1	-1	0	0	2
???	1	0	0	-1	0	1
s_1	1	1	0	0	1	5
z	-2	-1	0	0	0	0



$$\begin{array}{llllllllll}
 \min & w = & 0x_1 & + & 0x_2 & + & 0e_1 & + & 0e_2 & + & 0s_1 \\
 s.t. & & x_1 & + & x_2 & - & e_1 & & & & \geq & 2 \\
 & & x_1 & & & & & - & e_2 & & \geq & 1 \\
 & & x_1 & + & x_2 & & & & & + & s_1 & \leq & 5
 \end{array}$$

Degeneracy

Degeneracy occurs when a basic variable takes the value zero in a basic feasible solution. This leads to ties in the minimum ratio test, which determines the leaving variable during a pivot.

Problem

$$\begin{array}{ll}
 \max & z = x_1 + 2x_2 \\
 s.t. & x_1 + x_2 \leq 3 \\
 & x_2 \leq 2 \\
 & \frac{1}{2}x_1 + x_2 \leq 2.5 \\
 & x_1, x_2 \geq 0
 \end{array}$$

Standard Form


$$\begin{array}{l}
 z - x_1 - 2x_2 = 0 \\
 x_1 + x_2 + s_1 = 3 \\
 x_2 + s_2 = 2 \\
 \frac{1}{2}x_1 + x_2 + s_3 = 2.5 \\
 x_1, x_2, s_1, s_2, s_3 \geq 0
 \end{array}$$

Initialize Tableau

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	3
s_2	0	1	0	1	0	2
s_3	1/2	1	0	0	1	2.5
w	-1	-1	0	0	0	0

$$x_1 = 0$$

$$x_2 = 0$$



Minimum Test

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	3
s_2	0	1	0	1	0	2
s_3	1/2	1	0	0	1	2.5
w	-1	-1	0	0	0	0

Ratio Test

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	3
s_2	0	1	0	1	0	2
s_3	1/2	1	0	0	1	2.5
w	-1	-1	0	0	0	0

Pivot


Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	1	1	0	0	3
x_2	0	1	0	1	0	2
s_3	1/2	1	0	0	1	2.5
w	-1	-1	0	0	0	0

Gaussian Elimination

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	0	1	-1	0	1
x_2	0	1	0	1	0	2
s_3	1/2	0	0	-1	1	0.5
w	-1	0	0	2	0	4

$$x_1 = 0$$

$$x_2 = 2$$



Minimum Test

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	0	1	-1	0	1
x_2	0	1	0	1	0	2
s_3	1/2	0	0	-1	1	0.5
w	-1	0	0	2	0	4

Ratio Test

Basis	x_1	x_2	s_1	s_2	s_3	b
s_1	1	0	1	-1	0	1
x_2	0	1	0	1	0	2
s_3	1/2	0	0	-1	1	0.5
w	-1	0	0	2	0	4

$$\left. \begin{array}{l} \frac{1}{1} = 1 \\ \frac{0.5}{0.5} = 1 \end{array} \right\} \text{Tie}$$


Pivot

Basis	x_1	x_2	s_1	s_2	s_3	b
x_1	1	0	1	-1	0	1
x_2	0	1	0	1	0	2
s_3	1/2	0	0	-1	1	0.5
w	-1	0	0	2	0	4

Gaussian Elimination

Basis	x_1	x_2	s_1	s_2	s_3	b
x_1	1	0	1	-1	0	1
s_2	0	1	0	0	0	2
s_3	0	0	-1/2	-1/2	1	0
w	0	0	1	1	0	5

If after elimination a basic variable has a value of 0, it indicates degeneracy.



Degeneracy comes from redundant constraints

Bland's Rule: Choose the variable with the lowest index in the basis

- x_1 over x_2
- x_2 over s_1

Unbounded Solution

When identifying the entering variable, look at its column in the tableau. If all entries in that column (above the objective row) are ≤ 0 , the problem is unbounded

Can't perform ratio test \rightarrow no constraint limits the entering variable \rightarrow objective function increases without bound

Problem

$$\begin{array}{ll}\max & x_1 + x_2 \\s.t. & x_1 - x_2 \leq 1 \\ & x_1, x_2 \geq 0\end{array}$$

Standard Form

$$\begin{array}{ll}\max & z - x_1 - x_2 = 0 \\s.t. & x_1 - 2x_2 + s_1 = 1 \\ & x_1, x_2, s_1 \geq 0\end{array}$$

Initialize Tableau

Basis	x_1	x_2	s_1	b
s_1	1	-1	1	1
z	-1	-1	0	0

Minimum and Ratio Test

Basis	x_1	x_2	s_1	b
s_1	1	-1	1	1
z	-1	-1	0	0

Pivot

Basis	x_1	x_2	s_1	b
x_1	1	-1	1	1
z	-1	-1	0	0

Gaussian Elimination

Basis	x_1	x_2	s_1	b
x_1	1	-1	1	1

z	0	-2	1	1
-----	---	----	---	---

Minimum and Ratio Test

Basis	x_1	x_2	s_1	b
x_1	1	-1	1	1
z	0	-2	1	1

Check for Optimality

Objective row has -2 for $x_2 \rightarrow$ not optimal

Entering variable: x_2 (most negative coefficient)

Unboundedness Detection

Looking at the x_2 column:

Entry in constraint row: -1 (which is ≤ 0)

All entries in x_2 column are ≤ 0

Conclusion: Cannot perform ratio test \rightarrow Problem is unbounded

Alternative Solutions

Problem

$$\begin{aligned}
 \max \quad & 2x_1 + 4x_2 \\
 s.t. \quad & x_1 + x_2 \geq 3 \\
 & \frac{1}{2}x_1 + x_2 \geq 2.5 \\
 & x_1, x_2 \geq 0
 \end{aligned}$$

Standard Form

$$\begin{aligned}
 z - 2x_1 - x_2 &= 0 \\
 x_1 + x_2 + s_1 &= 3 \\
 \frac{1}{2}x_1 + x_2 + s_2 &= 2.5 \\
 x_1, x_2, x_3 &\geq 0
 \end{aligned}$$

Initialize Tableau

Basis	x_1	x_2	s_1	s_2	b
s_1	1	1	1	0	3
s_2	1/2	1	0	1	2.5
z	-2	-4	0	0	0

Minimum Test

Basis	x_1	x_2	s_1	s_2	b
s_1	1	1	1	0	3
s_2	1/2	1	0	1	2.5
z	-2	-4	0	0	0

Ratio Test

Basis	x_1	x_2	s_1	s_2	b
s_1	1	1	1	0	3
s_2	1/2	1	0	1	2.5
z	-2	-4	0	0	0


Pivot

Basis	x_1	x_2	s_1	s_2	b
s_1	1	1	1	0	3
x_2	1/2	1	0	1	2.5
z	-2	-4	0	0	0

Gaussian Elimination

Basis	x_1	x_2	s_1	s_2	b
s_1	1/2	0	1	-1	0.5
x_2	1/2	1	0	1	2.5
z	0	0	0	4	10

In a maximization or minimization linear programming problem, if there is a 0 in the z row (objective function row) of the final (optimal) simplex tableau in a non-basic column (i.e. a variable not currently in the solution), then there are multiple optimal solutions.



34.14.7. Standard Form

- **Equalities:** Convert inequalities (using slack, surplus, or artificial variables)
- **Non-negative variables:** If a variable is unrestricted or can be negative, replace it with the difference of two non-negative variables

$$2x_1 + 3x_2 \leq -4$$

$$\Leftrightarrow$$

$$-2x_1 - 3x_2 \geq 4$$

Nonpositive

$$2x_1 + 3x_2 \leq 4, \quad x_1 \leq 0$$

$$\Leftrightarrow$$

$$-2x_1 + 3x_2 \leq 4, \quad x_1 \geq 0$$

Free (Unrestricted)

$$2x_1 + 3x_2 \leq 6, \quad x_1 \text{ urs}$$

$$\Leftrightarrow$$

$$2x'_1 - 2x''_1 + 3x_2 \leq 6, \quad x'_1, x''_1 \geq 0$$

34.14.8. Simplex Method

$$\begin{aligned} \max \quad & 7x_1 + 6x_2 \\ \text{s.t.} \quad & 2x_1 + 4x_2 \leq 16 \\ & 3x_1 + 2x_2 \leq 12 \\ & x_1, x_2 \geq 0 \end{aligned}$$

$$\begin{aligned} \max \quad & 7x_1 + 6x_2 + 0s_1 + 0s_2 \\ \text{s.t.} \quad & 2x_1 + 4x_2 + s_1 = 16 \\ & 3x_1 + 2x_2 + s_2 = 12 \\ & x_1, x_2, s_1, s_2 \geq 0 \end{aligned}$$

Basis	c_B	x_1	x_2	s_1	s_2	b
s_1	0	2	4	1	0	16
s_2	0	3	2	0	1	12
<hr/>						
		0×2				
		+				
z_j		0×3				
		$= 0$				0
$c_j - z_j$		7	6	0	0	

$$x_1 = 0, x_2 = 0, s_1 = 16, s_2 = 12$$

Basis	c_B	x_1	x_2	s_1	s_2	b	Ratio
s_1	0	2	4	1	0	16	$16/2 = 8$
s_2	0	3	2	0	1	12	$12/3 = 4$
<hr/>							
z_j		0	0	0	0	0	
$c_j - z_j$		7	6	0	0		

1. Formulation

Objective Function

$$\begin{aligned} Z &= c_1x_1 + c_2x_2 + \dots + c_nx_n \\ &= c^T x \end{aligned}$$

Constraints

$$\begin{aligned} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n &\leq b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n &\geq b_2 \\ &\vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n &= b_m \\ x_1, x_2, \dots, x_n &\geq 0 \end{aligned}$$

Or,

$$Ax \leq, =, \geq b$$

$$x \geq 0$$

- \leq for maximization
- \geq for minimization

2. Convert to Canonical Form

In slack variable form

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n + s_1 = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n + s_2 = b_2$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n + s_m = b_m$$

$$x_1, x_2, \dots, x_n \geq 0$$

$$s_1, s_2, \dots, s_m \geq 0$$

Or matrix notation,

$$\begin{aligned} \max \quad & z = c^T x \\ \text{s.t.} \quad & Ax + Is = b_i \\ & x \geq 0 \\ & s \geq 0 \end{aligned}$$

$$\begin{aligned} \max \quad z &= c^T x \\ &= [c_1, c_2, \dots, c_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \end{aligned}$$

$$A \quad x \quad + \quad I \quad s \quad = \quad b$$

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{bmatrix} \begin{bmatrix} s_1 \\ s_2 \\ \vdots \\ s_m \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

3. Set up Simplex Tableau

	x_1	x_2	\dots	x_n	s_1	s_2	\dots	s_m	RHS
Constraint 1	a_{11}	a_{12}	\dots	a_{1n}	1	0	\dots	0	b_1
Constraint 2	a_{21}	a_{22}	\dots	a_{2n}	0	1	\dots	0	b_2
\vdots	\vdots	\vdots	\ddots	\vdots	\vdots	\vdots	\ddots	\vdots	\vdots
Constraint n	a_{m1}	a_{m2}	\dots	a_{mn}	0	0	\dots	1	b_m
Objective Function	$-c_1$	$-c_2$	\dots	$-c_n$	0	0	\dots	0	0

- $-$ for maximization
- $+$ for minimization

4. Perform Pivot Operation

- **Identify the Pivot Column:** Choose the column with the most:
 - Maximization: Negative coefficient
 - Minimization: Positive coefficient (or multiply the objective by -1 and treat it as a maximization)

in the objective function row (this indicates which variable will enter the basis).

- **Identify the Pivot Row:** Calculate the ratio of RHS to the pivot column's coefficients (only where coefficients are positive). The row with the smallest ratio determines the variable to leave the basis.
- **Pivot:** Perform row operations to change the pivot element to 1 and other elements in the pivot column to 0. This involves dividing the pivot row by the pivot element and then adjusting the other rows to zero out the pivot column.

5. Iterate

Repeat the pivot operations until there are no more negative coefficients in the objective function row, indicating that the current solution is optimal.

6. Read the Solution

The final tableau will give the values of the variables at the optimal solution. The basic variables are the variables corresponding to the columns with the identity matrix in the final tableau, while non-basic variables are set to zero.

Maximize : $Z = 3x_1 + 2x_2$

s.t.

$$x_1 + x_2 \leq 4$$

$$2x_1 + x_2 \leq 5$$

$$x_1, x_2 \geq 0$$

1. Convert to Canonical Form

Introduce slack variables s_1 and s_2 to convert inequalities into equalities:

$$x_1 + x_2 + s_1 = 4$$

$$2x_1 + x_2 + s_2 = 5$$

$$x_1, x_2, s_1, s_2 \geq 0$$

Initialize Simplex Tableau

	x_1	x_2	s_1	s_2	RHS
Constraint 1	1	1	1	0	4
Constraint 2	2	1	0	1	5
Objective Function	-3	-2	0	0	0

2. Identify the Pivot Column

In the objective function row, the coefficients are $[-3, -2, 0, 0]$. The most negative coefficient is -3 , which is in the x_1 column. Therefore, x_1 will enter the basis.

3. Identify the Pivot Row

Calculate the ratio of RHS to the pivot column's coefficients (where the coefficients are positive):

- For Row 1: $\frac{4}{1} = 4$ (pivot column coefficient is 1)
- For Row 2: $\frac{5}{2} = 2.5$ (pivot column coefficient is 2)

The smallest ratio is 2.5, so Row 2 will be the pivot row. This means that s_2 will leave the basis.

4. Pivot

Perform row operations to make the pivot element 1 and zero out other elements in the pivot column.

Pivot Element: The element at the intersection of Row 2 and x_1 column is 2.

Steps:

1. Make Pivot Element 1: Divide all elements in Row 2 by the pivot element (2):

$$\begin{aligned}\text{New Row 2} &= \frac{1}{2} \times \text{Old Row 2} \\ &= [1, 0.5, 0, 0.5, 2.5]\end{aligned}$$

2. Zero Out Pivot Column in Other Rows:

- For Row 1: Subtract 1 times New Row 2 from Row 1:

$$\begin{aligned}\text{New Row 1} &= \text{Old Row 1} - [1, 0.5, 0, 0.5, 2.5] \\ &= [0, 0.5, 1, -0.5, 1.5]\end{aligned}$$

- For Objective Function: Add 3 times New Row 2 to the Objective Function row


$$\begin{aligned}\text{New Objective Function} &= \text{Old Objective Function} + 3 \times [1, 0.5, 0, 0.5, 2.5] \\ &= [0, -0.5, 0, 1.5, 7.5]\end{aligned}$$

Updated Simplex Tableau

	x_1	x_2	s_1	s_2	RHS
Constraint 1	0	0.5	1	-0.5	1.5
Constraint 2	1	0.5	0	0.5	2.5
Objective Function	0	-0.5	0	1.5	7.5

34.15. Branch-and-Bound

Integer Program



Linear Relaxation:

IP \longrightarrow LP

Decompose an IP into multiple LPs

Definition 1: Linear Relaxation

For a Given LP, its linear relaxation is the resulting LP after removing all integer constraints

Integer Problem:

$$\begin{array}{ll} \max & 3x_1 + x_2 \\ \text{s.t.} & 4x_1 + 2x_2 \leq 11 \\ & x_i \in \mathbb{Z}_+ \quad \forall i = 1, 2 \end{array}$$

Linear Relaxation:

$$\begin{array}{ll} \max & 3x_1 + x_2 \\ \text{s.t.} & 4x_1 + 2x_2 \leq 11 \\ & x_i \geq 0 \quad \forall i = 1, 2 \end{array}$$

Integer Program:

$$\begin{array}{ll} \max & 16x_1 + 22x_2 + 12x_3 + 8x_4 \\ \text{s.t.} & 5x_1 + 7x_2 + 4x_3 + 3x_4 \leq 10 \\ & x_i \in \{0, 1\} \quad \forall i = 1, \dots, 4 \end{array}$$

Linear Relaxation:

$$\begin{array}{ll} \max & 16x_1 + 22x_2 + 12x_3 + 8x_4 \\ \text{s.t.} & 5x_1 + 7x_2 + 4x_3 + 3x_4 \leq 10 \\ & x_i \in [0, 1] \quad \forall i = 1, \dots, 4 \end{array}$$

$$x_i \in [0, 1] \iff x_i \geq 0 \wedge x_i \leq 1$$

Linear relaxation provides a bound


- z^* : Optimal value of the Integer Program (IP)
- z' : Optimal value of the Linear Relaxation (LP)

1. Minimization

- For **minimization** IP, linear relaxation provides a **lower bound**
- The feasible region of the LP contains all feasible solutions of the IP (i.e., superset)
- Therefore, the LP can achieve a value that is at most as large as the IP
- Conclusion:

$$z' \leq z^*$$

So, linear relaxation provides a lower bound on the optimal value of the integer program




2. Maximization

- For **maximization** IP, linear relaxation provides a **upper bound**
- The LP may find a solution with an objective value greater than or equal to that of any feasible integer solution.
- Conclusion:

$$z' \geq z^*$$

- So, linear relaxation provides an upper bound on the optimal value of the integer program



Problem Type	Linear Relaxation Provides	Inequality Between LP and IP
Minimization IP	Lower Bound	$z' \leq z^*$
Maximization IP	Upper Bound	$z' \geq z^*$








- If the linear relaxation is infeasible or unbounded then the IP is infeasible or unbounded
- If an optimal solution to the linear relaxation is feasible,

Let x' be an optimal solutions to the linear relaxation of an IP. If x' is feasible to the IP, it is the optimal to the IP

$$\begin{array}{ll}
 \max & z = 3x_1 + 4x_2 \\
 s.t. & 2x_1 + x_2 \leq 6 \\
 & 2x_1 + 3x_2 \leq 9 \\
 & x_i \in \mathbb{N}_+ \quad \forall i = 1, 2
 \end{array}$$

$$\begin{array}{ll}
 \max & z = 3x_1 + 4x_2 \\
 s.t. & 2x_1 + x_2 \leq 6 \\
 & 2x_1 + 3x_2 \leq 9 \\
 & x_i \geq 0 \quad \forall i = 1, 2
 \end{array}$$

$$\begin{array}{l}
 x_1 = \frac{9}{4}, \quad x_2 = \frac{3}{2} \\
 z = 12.75
 \end{array}$$



34.16. Gradient Descent

34.16.1. Gradient

The gradient tells us the direction of steepest increase

For a function

$$f(x_1, x_2, \dots, x_n)$$

the gradient of f , written ∇f , is:

$$\nabla f(x) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)$$

- The **magnitude** of the gradient vector tells you how fast the function is increasing (steepness)
- The **direction** of the gradient vector points to where the function increases most rapidly (direction)

Function

$$f(x_1, x_2) = x_1^2 + x_2^2$$

Compute $\frac{\partial f}{\partial x_1}$

- Differentiate f w.r.t. x_1 (treat x_2 as constant):

$$\frac{\partial}{\partial x_1}(x_1^2 + x_2^2) = \frac{\partial}{\partial x_1}(x_1^2) + \frac{\partial}{\partial x_1}(x_2^2)$$

- Since x_2^2 is constant w.r.t. x_1 , its derivative is 0:

$$\frac{\partial}{\partial x_1}(x_2^2) = 0$$

- The derivative of x_1^2 w.r.t. x_1 is:

$$\frac{\partial}{\partial x_1}(x_1^2) = 2x_1$$

- So,

$$\frac{\partial f}{\partial x_1} = 2x_1$$

Compute $\frac{\partial f}{\partial x_2}$

- Differentiate f w.r.t. x_2 (treat x_1 as constant):

$$\frac{\partial}{\partial x_2}(x_1^2 + x_2^2) = \frac{\partial}{\partial x_2}(x_1^2) + \frac{\partial}{\partial x_2}(x_2^2)$$

- Since x_1^2 is constant w.r.t. x_2 , its derivative is 0:

$$\frac{\partial}{\partial x_2}(x_1^2) = 0$$

- The derivative of x_2^2 w.r.t. x_2 is:

$$\frac{\partial}{\partial x_2}(x_2^2) = 2x_2$$

So,

$$\frac{\partial f}{\partial x_2} = 2x_2$$

Combine to form gradient

$$\nabla f(x_1, x_2) = (2x_1, 2x_2)$$

Evaluate at a point

$$x = (1, 2)$$

$$\nabla f(1, 2) = (2 \times 1, 2 \times 2) = (2, 4)$$

Interpretation

- The gradient vector $(2, 4)$ at $(1, 2)$ points in the **direction** where the function f **increases most rapidly**
- Its **magnitude** $\sqrt{2^2 + 4^2} = \sqrt{20} \approx 4.47$ measures the **steepness** of that increase
- **Moving one unit along the x_1 axis alone would increase the function by the partial derivative with respect to x_1 , which is 2**
- **Moving one unit along the x_2 axis alone would increase the function by the partial derivative with respect to x_2 , which is 4**
- **Moving from $(1, 2)$ in the direction of $(2, 4)$, the function value will rise fastest and at a rate roughly proportional to 4.47 per unit distance moved**

34.16.2. Algorithm

Find the minimum of a function

In **gradient descent**, we move in the **opposite direction** of the gradient to minimize a function

Direction of **maximum increase**:

$$\nabla f(x)$$

Direction of **maximum decrease**:

$$-\nabla f(x)$$

1. **Initialize**: Start with an initial guess for the parameters
2. **Compute Gradient**: Find the gradient of the function at the current parameters

3. **Update Parameters:** Adjust the parameters by moving in the opposite direction of the gradient, scaled by the learning rate
4. **Repeat:** Continue the process until the parameters converge to a minimum or the changes are minimal

Update Rule:

$$x^{k+1} \leftarrow x^k - \alpha_k \nabla f(x^k)$$

Where:

- x^k : current iterate (point in domain of f)
- α_k : **step size** or **learning rate** at iteration k
- $\nabla f(x^k)$: **gradient** of the function f with respect to θ
- x^{k+1} : next iterate (point in domain of f)

```
import numpy as np

def gradient_descent(
    f,                # Function to minimize, f(x)
    grad_f,           # Gradient of f, grad_f(x)
    x0,               # Initial guess for x
    alpha=0.01,       # Learning rate
    max_iter=1000,    # Maximum number of iterations
    tol=1e-6          # Tolerance for stopping
):
    x = x0.copy()
    for _ in range(max_iter):
        grad = grad_f(x)
        if np.linalg.norm(grad) < tol:
            break
        x -= alpha * grad
    return x


def f(x):
    """Example: minimize f(x) = x^2 + 2x + 1"""
    return x**2 + 2*x + 1

def grad_f(x):
    """Gradient of f(x) = x^2 + 2x + 1 is f'(x) = 2x + 2"""
    return 2*x + 2

gradient_descent(
    f,
    grad_f,
    x_initial,
    alpha=0.1,
    max_iter=100
)
```

Function

$$x_1^2 + x_2^2$$



Gradient

$$\nabla f(x) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2} \right) = (2x_1, 2x_2)$$

Learning Rate

$$\alpha = 0.1$$

Iteration 1

- Initial Parameter Values:

$$x_1 = 1 \quad x_2 = 2$$

- Gradient Calculation

$$\nabla f(x) = (2 \times 1, 2 \times 2) = (2, 4)$$

- Parameter Update:

$$x \leftarrow x - \alpha \nabla f(x)$$

$$x_1 \leftarrow 1 - 0.1 \times 2 = 0.8$$

$$x_2 \leftarrow 2 - 0.1 \times 4 = 1.6$$

- Updated Parameter Values:

$$x_1 = 0.8 \quad x_2 = 1.6$$

Iteration 2

- Current Parameter Values

$$x_1 = 0.8 \quad x_2 = 1.6$$

- Gradient Calculation:

$$\nabla f(x) = (2 \times 0.8, 2 \times 1.6) = (1.6, 3.2)$$

- Parameter Update:

$$x \leftarrow x - \alpha \nabla f(x)$$

$$x_1 \leftarrow 0.8 - 0.1 \times 1.6 = 0.64$$

$$x_2 \leftarrow 1.6 - 0.1 \times 3.2 = 1.28$$

- Updated Parameter Values:

$$x_1 = 0.64 \quad x_2 = 1.28$$

Iteration 3

- Current Values

$$x_1 = 0.64 \quad x_2 = 1.28$$

- Gradient Calculation:

$$\nabla f(x) = (2 \times 0.64, 2 \times 1.28) = (1.28, 2.56)$$

- Parameter Update:

$$x \leftarrow x - \alpha \nabla f(x)$$

$$x_1 \leftarrow 0.64 - 0.1 \times 1.28 = 0.512$$

$$x_2 \leftarrow 1.28 - 0.1 \times 2.56 = 1.024$$


- Updated Parameter Values:

$$x_1 = 0.512 \quad x_2 = 1.024$$

Problem Setup

Minimize the quadratic function:

$$\min_{x \in \mathbb{R}^2} f(x) = 4x_1^2 - 4x_1x_2 + 2x_2^2 \quad \text{where} \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$



Gradient of $f(x)$

The gradient is:

$$\nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \end{bmatrix}$$

Step 1: Compute $\frac{\partial f}{\partial x_1}$

Take the derivative of each term with respect to x_1 (treat x_2 as a constant):

- $\frac{\partial}{\partial x_1}(4x_1^2) = 8x_1$
- $\frac{\partial}{\partial x_1}(-4x_1x_2) = -4x_2$
- $\frac{\partial}{\partial x_1}(2x_2^2) = 0$

So:

$$\frac{\partial f}{\partial x_1} = 8x_1 - 4x_2$$

Step 2: Find $\frac{\partial f}{\partial x_2}$

Take the derivative of each term with respect to x_2 (treat x_1 as a constant):

- $\frac{\partial}{\partial x_2}(4x_1^2) = 0$
- $\frac{\partial}{\partial x_2}(-4x_1x_2) = -4x_1$
- $\frac{\partial}{\partial x_2}(2x_2^2) = 4x_2$

So:

$$\frac{\partial f}{\partial x_2} = -4x_1 + 4x_2$$

So the gradient of $f(x)$ is:

$$\nabla f(x) = \begin{bmatrix} 8x_1 - 4x_2 \\ -4x_1 + 4x_2 \end{bmatrix}$$

Optimal Solution

$$x^* = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$
$$f(x^*) = 0$$

Gradient Descent Iterations

Iteration 1

- Initial guess:

$$x^0 = \begin{bmatrix} 2 \\ 3 \end{bmatrix} \Rightarrow f(x^0) = 4(2^2) - 4(2)(3) + 2(3^2) = 10$$

- Gradient at x^0 :

$$\nabla f(x^0) = \begin{bmatrix} 8(2) - 4(3) \\ -4(2) + 4(3) \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \end{bmatrix}$$

- Line Search:

$$x(\alpha_0) = x^0 - \alpha_0 \nabla f(x^0) = \begin{bmatrix} 2 \\ 3 \end{bmatrix} - \alpha_0 \begin{bmatrix} 4 \\ 4 \end{bmatrix}$$
$$= \begin{bmatrix} 2 - 4\alpha_0 \\ 3 - 4\alpha_0 \end{bmatrix}$$

$$f(x(\alpha_0)) = 4x_1^2 - 4x_1x_2 + 2x_2^2$$
$$= 4(2 - 4\alpha_0)^2 - 4(2 - 4\alpha_0)(3 - 4\alpha_0) + 2(3 - 4\alpha_0)^2$$
$$= 32\alpha_0^2 - 32\alpha_0 + 10$$

- Minimizing:

$$\alpha_0 = \operatorname{argmin} f(x(\alpha))$$

$$\frac{\partial}{\partial \alpha} f(x(\alpha_0)) = \frac{\partial}{\partial \alpha} (32\alpha_0^2 - 32\alpha_0 + 10) = 0$$

$$= 64\alpha_0 - 32 = 0$$

$$\alpha_0 = \frac{1}{2}$$

- Update Step:

$$x^1 = x^0 - a_0 \nabla f(x^0) = \begin{bmatrix} 2 \\ 3 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 4 \\ 4 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Check Progress:

1. New point:

$$x^1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

2. Function value decreases $f(x^{k+1}) < f(x^k)$

$$f(x^1) = 2 < f(x^0) = 10$$

3. Gradient at new point:

$$\nabla f(x^1) = \begin{bmatrix} 8(0) - 4(1) \\ -4(0) + 4(1) \end{bmatrix} = \begin{bmatrix} -4 \\ 4 \end{bmatrix}$$

4. Gradient magnitude $\|\nabla f(x^{k+1})\| < \|\nabla f(x^k)\|$

$$\|\nabla f(x^1)\| = 4\sqrt{2} = \|\nabla f(x^0)\|$$

Note: Gradient magnitude stays the same in this iteration due to the specific structure of this quadratic function

Iteration 2

- New Point:

$$x^1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

- Gradient at x_1 :

$$\nabla f(x^1) = \begin{bmatrix} 8(0) - 4(1) \\ -4(0) + 4(1) \end{bmatrix} = \begin{bmatrix} -4 \\ 4 \end{bmatrix}$$

$$a_1 = \operatorname{argmin} f(x(\alpha_1))$$

- Line search:

$$\begin{aligned} x(\alpha_1) &= x^1 - \alpha_1 \nabla f(x^1) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} - \alpha_1 \begin{bmatrix} -4 \\ 4 \end{bmatrix} \\ &= \begin{bmatrix} 4\alpha_1 \\ 1 - 4\alpha_1 \end{bmatrix} \end{aligned}$$

$$\begin{aligned} f(x(\alpha_1)) &= 4x_1^2 - 4x_1x_2 + 2x_2^2 \\ &= 4(4\alpha_1)^2 - 4(4\alpha_1)(1 - 4\alpha_1) + 2(1 - 4\alpha_1)^2 \\ &= 16\alpha_1^2 - 32\alpha_1 + 2 \end{aligned}$$

- Minimize:

$$\alpha_1 = \operatorname{argmin} f(x(\alpha_1)) = \frac{1}{10}$$

$$\begin{aligned}\frac{\partial}{\partial \alpha} f(x(\alpha_1)) &= \frac{\partial}{\partial \alpha} (160\alpha_1^2 - 32\alpha_1 + 2) = 0 \\ &= 320\alpha_1 - 32 = 0\end{aligned}$$

$$\alpha_1 = \frac{1}{10}$$

- Update Step:

$$x^2 = x^1 - \alpha_1 \nabla f(x^1) = \begin{bmatrix} 0 \\ 1 \end{bmatrix} - \frac{1}{10} \begin{bmatrix} -4 \\ 4 \end{bmatrix} = \begin{bmatrix} 0.4 \\ 0.6 \end{bmatrix}$$

- Improvement:

1. New point:

$$x_2 = \begin{bmatrix} 0.4 \\ 0.6 \end{bmatrix}$$

2. Function value decreases $f(x^{k+1}) < f(x^k)$

$$f(x^2) = 0.4 < f(x^1) = 2$$

3. Gradient at new point:

$$\nabla f(x_2) = \begin{bmatrix} 8(0.4) - 4(0.6) \\ -4(0.4) + 4(0.6) \end{bmatrix} = \begin{bmatrix} 0.8 \\ 0.8 \end{bmatrix}$$


4. Gradient magnitude $\|\nabla f(x^{k+1})\| < \|\nabla f(x^k)\|$

$$\|\nabla f(x^2)\| = \|(0.8, 0.8)\| = \frac{4\sqrt{2}}{5}$$

Problem Setup


Minimize the quadratic function:

$$\min_{x \in \mathbb{R}^2} f(x) = x_1^2 - 2x_1x_2 + 2x_2^2 + 2x_1 \quad \text{where} \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$



Gradient of $f(x)$

$$\nabla f(x_1, x_2) = \begin{bmatrix} 2x_1 - 2x_2 + 2 \\ -2x_1 + 4x_2 \end{bmatrix}$$



34.17. Newton's Method

Iterative numerical technique to find local minima or maxima of a function

Taylor Series (Expansion)

Approximate a complicated function near a point a by a polynomial made from the function's derivatives at that point

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

Where:

- $f^{(n)}(a)$: n -th derivative of f evaluated at a

Setup

$$f(x) = x^2$$

Expand it around the point $a = 1$

Step 1: Compute Derivatives

- $f(x) = x^2$
- $f'(x) = 2x$
- $f''(x) = 2$

Step 2: Evaluate at $x = 1$

- $f(1) = 1^2 = 1$
- $f'(1) = 2 \cdot 1 = 2$
- $f''(1) = 2$


Step 3: Write the Taylor expansion

$$\begin{aligned} f(x) &= f(1) + f'(1)(x-1) + \frac{f''(1)}{2!}(x-1)^2 \\ &= 1 + 2(x-1) + \frac{2}{2}(x-1)^2 \\ &= 1 + 2(x-1) + (x-1)^2 \end{aligned}$$

Step 3: Expand

$$1 + 2(x-1) + (x-1)^2 = 1 + 2x - 2 + x^2 + 1 = x^2$$

It gives back the original function exactly



34.18. Monte Carlo