
a **Z80 competitions: transmitter share and byte-motif prevalence (L = 32)**

50:50 (10 worlds) · 1% R in T (5 worlds) · 1% T in R (5 worlds) · Archive: light/open · C1: dark/filled

Regenerator R | Transmitter T | World medians; C1 = corrected Z80 instructions



Endpoint medians: C1 | archive


Start	T share	Motif share
50:50	0.05 | 0.06	0.85 | 0.84
1% R in T	0.09 | 0.07	0.88 | 0.80
1% T in R	0.08 | 0.08	0.76 | 0.87

Endpoint medians: C1 | archive

Start	T share	Motif share
50:50	0.76 | 0.77	0.90 | 0.92
1% R in T	0.81 | 0.76	0.91 | 0.88
1% T in R	0.77 | 0.78	0.88 | 0.93

b **Final world distributions at 300,000 world updates**

Open: archive / filled: C1 implementation



20 worlds/group; C1 benign ratio n = 19 (one world has R + T = 0)

c **Random-code dial: L = 16; 10 worlds per implementation per p**

Each entry: C1 | archive

p: probability a zero instruction halts	0	0.01	0.03	0.1	0.3	1
First copier open (out of 10)	10 | 10	10 | 10	10 | 10	9 | 10	4 | 2	0 | 0
T >50% heritable cells at 300k (out of 10)	0 | 1	1 | 1	1 | 1	3 | 1	3 | 6	8 | 6
Kaplan-Meier median to first confined copier (thousand world updates)	32.5 | 18	42.5 | 42.5	60 | 60	80 | 75	212.5 | 57.5	115 | 52.5

Open = partner opcode fetch; confined = no partner fetch