
a **Z80 competitions: full-tape T share and motif prevalence (L = 32)**




Endpoint medians: Corrected | Historical

Start	T share	Motif share
50:50	0.05 | 0.06	0.85 | 0.84
1% R in T	0.09 | 0.07	0.88 | 0.80
1% T in R	0.08 | 0.08	0.76 | 0.87

Endpoint medians: Corrected | Historical

Start	T share	Motif share
50:50	0.76 | 0.77	0.90 | 0.92
1% R in T	0.81 | 0.76	0.91 | 0.88
1% T in R	0.77 | 0.78	0.88 | 0.93

b **Final world distributions at 300,000 updates**



20 worlds/group; Corrected benign ratio n = 19 (one world has R + T = 0)

Historical: open; Corrected: filled

c **Random-code dial: 10 Corrected + 10 Historical worlds at each p; L = 16**

Entries: Corrected | Historical

p: probability fetching a zero opcode halts	0	0.01	0.03	0.1	0.3	1
First copier open (out of 10)	10 | 10	10 | 10	10 | 10	9 | 10	4 | 2	0 | 0
Confined T >50% of heritable cells at 300,000 updates (T: ≥5 bytes retain introduced values; out of 10)	0 | 1	1 | 1	1 | 1	3 | 1	3 | 6	8 | 6
Kaplan–Meier median to first confined copier (thousand world updates)	32.5 | 18	42.5 | 42.5	60 | 60	80 | 75	212.5 | 57.5	115 | 52.5

Open: partner instruction-stream byte fetch in ≥8/16 traces; confined: none in 16/16.