a Genealogy cohorts: worlds and nested lines

Cohort	Worlds	Confined worlds	Lines from confined worlds	Open-first lines
N3-v6	20	17	1088	1088
N6	20	20	1280	1280

64 sampled-cell ancestries / confined world; world = simulation

N3-v6 = discovery; N6 = prospective confirmation; Z80 L16

Copier: ≥75% tape bytes copied in ≥32/64 uniform partners

Counted trace = fetch of any partner opcode or operand byte

Genealogy fetch class: open ≥32/64 counted traces; confined <32/64

Open-first: earliest copier on each sampled line is classified open

Sampled lines: 64 ancestries per world with a confined founder
Worlds column counts all worlds; genealogy classes use 64 partner traces

b First founders versus later conversions

Cohort	First confined founders	Later confined founder events	Events
N3-v6	17	15	32
N6	20	14	34

20 worlds/cohort; at most one first confined founder/world

Founder: first confined copier after latest open copier on sampled line

Transition: open ancestry to confined copier; sampled classifier as a

Later events follow the first founder in that world; events can repeat

Later events can reuse existing confined-cell byte sources

First founders and later confined-founder events stay separate
Worlds, ancestral lines and founding events are different units

c N6 first-founder mechanisms

Recorded mechanism	Founders /20
novel	18/20
damage	2/20

novel = rewrite, <75% byte match to either parent (18)

damage = partial overwrite, ≥75% parent byte match (2)

Separate N6-1 source rule: copy/rewrite 18/20, mutation 0/20
Recorded event categories and causal source classes differ

d N6 founding executors: two copier scopes

Executors, n	Whole-tape	Executed-byte
Non-copier	20	11
Copier	0	9

N6-4: 11/20 executed-byte non-copiers; requires 12/20; not met

Whole-tape: 20/20 non-copiers; requires ≥16/20

Soup-abundance null: P(all 20 whole-tape non-copiers) = 0.062

Null draws executors by pre-founder soup composition

One founding executor/first confined founder: 20 N6 worlds
Copier: ≥75% scoped bytes copied in ≥50% partner encounters
Scopes: entire tape / only executed instruction-stream bytes

e N6: all first-founder chain lengths



N6: one point per first founder, 20 worlds
Records = contributing chain events; intervening whole-tape
donors are non-copiers (executed-byte function kept separate)

f Registered N6: complete component ledger

ID	Requirement	Observation	Verdict
N6-1	Copy/rewrite ≥16/20; mutation ≤2/20 founders	Copy/rewrite 18/20 Mutation 0/20	met
N6-2	Whole-tape non-copier chains ≥15/20	20/20	met
N6-3	Median cell carrier share ≥5× random-tape baseline	Observed 0.00190 Baseline 0.00024 Ratio 7.9×	met
N6-4	Non-copier executors whole ≥16/20; exec ≥12/20	Whole 20/20 Executed-byte 11/20 Null P = 0.062	**not met**
N6-5′	Expected mutation founders <0.05 in 20-world cohort	0.0064 expected 0/20 observed founders	met

Input: 55 point-mutation events on 20 pre-founder chains
N6-3: cells carrying a word / all soup cells; 2-byte word at executed start
Median across 99 word/founder rows in pre-founder snapshots
Reject if point mutations complete ≥5/20 first founders; observed 0/20