

Real analysis

Zijie Xia

2024-08-01

Contents

1	Prerequisites	5
2	Measure Theory	7
2.1	Preliminaries	7
2.2	The exterior measure	8
2.3	Measurable sets and the Lebesgue measure	10
2.4	Measurable functions	13
2.5	The Brunn-Minkowski inequality	15
2.6	Exercise	16
2.7	Problem	16
3	Integration theory	17
3.1	The lebesgue integral: basic properties and convergence theorems	17
4	Methods	21
4.1	math example	21
5	Applications	23
5.1	Example one	23
5.2	Example two	23
6	Final Words	25

Chapter 1

Prerequisites

This is a *sample* book written in **Markdown**. You can use anything that Pandoc's Markdown supports, e.g., a math equation $a^2 + b^2 = c^2$.

The **bookdown** package can be installed from CRAN or Github:

```
install.packages("bookdown")  
# or the development version  
# devtools::install_github("rstudio/bookdown")
```

Remember each Rmd file contains one and only one chapter, and a chapter is defined by the first-level heading #.

To compile this example to PDF, you need XeLaTeX. You are recommended to install TinyTeX (which includes XeLaTeX): <https://yihui.name/tinytex/>.

Chapter 2

Measure Theory

2.1 Preliminaries

The **open ball** in \mathbb{R}^d centered at x and of radius r is defined by

$$B_r(x) = \{y \in \mathbb{R}^d : |y - x| < r\}.$$

A subset $E \subset \mathbb{R}^d$ is **open** if for every $x \in E$ there exists $r > 0$ with $B_r(x) \subset E$. By definition, a set is **closed** if its complement is open.

We note that any (not necessarily countable) union of open sets is open, while in general the intersection of only finitely many open sets is open. A similar statement holds for the class of closed sets, if one interchanges the roles of unions and intersections.

A set E is **bounded** if it is contained in some ball of finite radius. A bounded set is **compact** if it is also closed. Compact sets enjoy the **Heine-Borel covering property**:

- Assume E is compact, $E \subset \bigcup_{\alpha} O_{\alpha}$, and each O_{α} is open. Then there are finitely many of the open sets, $O_{\alpha_1}, O_{\alpha_2}, \dots, O_{\alpha_N}$, such that $E \subset \bigcup_{j=1}^N O_{\alpha_j}$.

In words, any covering of a compact set by a collection of open sets contains a finite subcovering.

A point $x \in \mathbb{R}^d$ is a **limit point** of the set E if for every $r > 0$, the ball $B_r(x)$ contains points of E . This means that there are points in E which are arbitrarily close to x . An **isolated point** of E is a point $x \in E$ such that there exists an $r > 0$ where $B_r(x) \cap E$ is equal to $\{x\}$.

A point $x \in E$ is an **interior point** of E if there exists $r > 0$ such that $B_r(x) \subset E$. The set of all interior points of E is called the **interior** of E . Also,

the **closure** \overline{E} of the E consists of the union of E and all its *limit points*. The **boundary** of a set E , denoted by ∂E , is the set of points which are in the closure of E but not in the interior of E .

Note that the closure of a set is a closed set; every point in E is a limit point of E ; and a set is closed if and only if it contains all its limit points. Finally, a closed set E is **perfect** if E does not have any isolated points.

Lemma 2.1. *If a rectangle is the almost disjoint union of finitely many other rectangles, say $R = \bigcup_{k=1}^N R_k$, then*

$$|R| = \sum_{k=1}^N |R_k|$$

Lemma 2.2. *If R, R_1, \dots, R_N are rectangles, and $R \subset \bigcup_{k=1}^N R_k$, then*

$$|R| \leq \sum_{k=1}^N |R_k|$$

Proof.

Every open subset O of \mathbb{R} can be written uniquely as a countable union of disjoint open intervals.

Every open subset O of \mathbb{R}^d , $d \geq 1$, can be written uniquely as a countable union of almost disjoint closed cubes.

The **Cantor set** C is by definition the intersection of all C_k 's:

$$C = \bigcap_{k=0}^{\infty} C_k.$$

The set C is not empty, since all end-points of the intervals in C_k (all k) belong to C . Despite its simple construction, the Cantor set enjoys many interesting topological and analytical properties. For instance, C is closed and bounded, hence compact. Also, C is totally disconnected: given any $x, y \in C$ there exists $z \notin C$ that lies between x and y . Finally, C is perfect: it has no isolated points.

In terms of cardinality the Cantor set is rather large: it is not countable. Since it can be mapped to the interval $[0, 1]$, the Cantor set has the cardinality of the continuum. However, from the point view of “length” the size of C is small. Roughly speaking, it has length zero.

2.2 The exterior measure

Definition 2.1. If E is any subset of \mathbb{R}^d , the **exterior measure** of E is

$$m_*(E) = \inf \sum_{j=1}^{\infty} |Q_j|,$$

where the infimum is taken over all countable coverings $E \subset \bigcup_{j=1}^{\infty} Q_j$ by closed cubes. In general we have $0 \leq m_*(E) \leq \infty$.

- It would not suffice to allow finite sums in the definition of $m_*(E)$. The quantity that would be obtained if one considered only coverings of E by finite unions of cubes is in general larger than $m_*(E)$.
- One can, however, replace the coverings by cubes, with coverings by rectangles; or with coverings by balls.

Example 2.1. The exterior measure of a closed cube is equal to its volume.

Proof.

We consider an arbitrary covering $Q \subset \bigcup_{j=1}^{\infty} Q_j$ by cubes, where Q_j is closed for $j = 1, 2, \dots$. Then it suffices to prove that

$$|Q| \leq \sum_{j=1}^{\infty} |Q_j|.$$

Example 2.2. The exterior measure of an open cube is equal to its volume.

Example 2.3. The exterior measure of a rectangle is equal to its volume.

Proof.

Arguing as in the first example, we see that $|R| \leq m_*(R)$. To obtain the reverse inequality, consider a grid in \mathbb{R}^d formed by cubes of side length $1/k$, then let k tend to infinity yields $m_*(R) \leq |R|$.

Example 2.4. The Cantor set C has exterior measure 0.

2.2.1 Properties of the exterior measure

From the definition we know that: - For every $\epsilon > 0$, there exists a covering $E \subset \bigcup_{j=1}^{\infty} Q_j$ with

$$\sum_{j=1}^{\infty} m_*(Q_j) \leq m_*(E) + \epsilon.$$

Proposition 2.1.

1. (Monotonicity) If $E_1 \subset E_2$, then $m_*(E_1) \leq m_*(E_2)$.
2. (Countable sub-additivity) If $E = \bigcup_{j=1}^{\infty} E_j$, then $m_*(E) \leq \sum_{j=1}^{\infty} m_*(E_j)$.
3. If $E \subset \mathbb{R}^d$, then $m_*(E) = \inf m_*(O)$, where the infimum is taken over all open sets O containing E .

Proof.

It suffices to prove that $m_*(E) \geq \inf m_*(O)$. Notice that for any closed cube Q_j , there exists an open set S_j which contains Q_j and such that $|S_j| \leq (1 + \epsilon)|Q_j|$ for a fixed $\epsilon > 0$.

4. If $E = E_1 \cup E_2$, and $d(E_1, E_2) > 0$, then

$$m_*(E) = m_*(E_1) + m_*(E_2).$$

5. If a set E is the countable union of almost disjoint cubes $E = \bigcup_{j=1}^{\infty} Q_j$, then

$$m_*(E) = \sum_{j=1}^{\infty} |Q_j|.$$

Proof.

Consider constructing a family of cubes that are at a finite distance from one another, so we can use proposition 4.

2.3 Measurable sets and the Lebesgue measure

The notion of measurability isolates a collection of subsets in \mathbb{R}^d for which the exterior measure satisfies all our desired properties, including additivity (and in fact countable additivity) for disjoint unions of sets.

There are a number of different ways of defining measurability, but these all turn out to be equivalent. Probably the simplest and most intuitive is the following: A subset E of \mathbb{R}^d is **Lebesgue measurable**, or simply **measurable**, if for any $\epsilon > 0$ there exists an open set O with $E \subset O$ and

$$m_*(O - E) \leq \epsilon.$$

This should be compared to Proposition 2.1 3, which holds for all sets E . If E is measurable, we define its **Lebesgue measure** (or **measure**) $m(E)$ by

$$m(E) = m_*(E).$$

Immediately from the definition, we find:

Proposition 2.2.

1. Every open set in \mathbb{R}^d is measurable.
2. If $m_*(E) = 0$, then E is measurable. In particular, if F is a subset of a set of exterior measure 0, then F is measurable.
3. A countable union of measurable sets is measurable.
4. Closed sets are measurable.

Proof.

Since any closed set F can be written as the union of compact sets, say $F = \bigcup_{k=1}^{\infty} F \cap B_k$, where B_k denotes the closed ball of radius k centered at the origin, it suffices to prove that compact sets are measurable.

5. The complement of a measurable set is measurable.

Proof.

For every positive integer n we choose an open set O_n with $E \subset O_n$ and $m_*(O_n - E) \leq 1/n$. Notice that

$$(E^c - \bigcup_{n=1}^{\infty} O_n^c) \subset (O_n - E)$$

then E^c is measurable since $E^c = (E^c - \bigcup_{n=1}^{\infty} O_n^c) \cup \bigcup_{n=1}^{\infty} O_n^c$.

6. A countable intersection of measurable sets is measurable.

To prove the fourth proposition, we need the following lemma.

Lemma 2.3. *If F is closed, K is compact, and these sets are disjoint, then $d(F, K) > 0$.*

Theorem 2.1. *If E_1, E_2, \dots , are disjoint measurable sets, and $E = \bigcup_{j=1}^{\infty} E_j$, then*

$$m(E) = \sum_{j=1}^{\infty} m(E_j).$$

Proof.

If F_1, F_2, \dots, F_N , are compact and disjoint, then obviously for any $j, k, j \neq k$, $d(F_j, F_k) > 0$, so $m\left(\bigcup_{j=1}^N F_j\right) = \sum_{j=1}^N m(F_j)$. If each E_j is bounded, we can choose a closed subset F_j for E_j with $m_*(E_j - F_j) \leq \epsilon/2^j$ for each j . Then

$$m(E) \geq \sum_{j=1}^N m(F_j) \geq \sum_{j=1}^N m(E_j) - \epsilon$$

Letting N tend to infinity, since ϵ is arbitrary we find that

$$m(E) \geq \sum_{j=1}^{\infty} m(E_j).$$

In the general case, we select any sequence of cubes $\{Q_k\}_{k=1}^{\infty}$ that increases to \mathbb{R}^d , in the sense that $Q_k \subset Q_{k+1}$ for all $k \geq 1$ and $\bigcup_{k=1}^{\infty} Q_k = \mathbb{R}^d$. We then let $S_1 = Q_1$ and $S_k = Q_k - Q_{k-1}$ for $k \geq 2$. If we define measurable sets by $E_{j,k} = E_j \cap S_k$, then

$$m(E) = \sum_{j,k} m(E_{j,k}) = \sum_j \sum_k m(E_{j,k}) = \sum_j m(E_j).$$

Corollary 2.1. *Suppose E_1, E_2, \dots are measurable subsets of \mathbb{R}^d .*

- If $E_k \nearrow E$, then $m(E) = \lim_{N \rightarrow \infty} m(E_N)$.

- If $E_k \searrow E$ and $m(E_k) < \infty$ for some k , then $m(E) = \lim_{N \rightarrow \infty} m(E_N)$.

Theorem 2.2. Suppose E is measurable subset of \mathbb{R}^d . Then for every $\epsilon > 0$:

1. There exists an open set O with $E \subset O$ and $m(O - E) \leq \epsilon$.
2. There exists a closed set F with $F \subset E$ and $m(E - F) \leq \epsilon$.
3. If $m(E)$ is finite, there exists a compact set K with $K \subset E$ and $m(E - K) \leq \epsilon$.
4. If $m(E)$ is finite, there exists a finite union $F = \bigcup_{j=1}^N Q_j$ of closed cubes such that

$$m(E \triangle F) \leq \epsilon.$$

Proof.

Choose a family of closed cubes $\{Q_j\}_{j=1}^\infty$ so that

$$E \subset \bigcup_{j=1}^\infty Q_j \text{ and } \sum_{j=1}^\infty |Q_j| \leq m(E) + \epsilon/2.$$

Since $m(E) < \infty$, then series converges and there exists $N > 0$ such that $\sum_{j=n+1}^\infty |Q_j| \leq \epsilon/2$. If $F = \bigcup_{j=1}^N Q_j$, then

$$\begin{aligned} m(E \triangle F) &= m(E - F) + m(F - E) \\ &\leq m\left(\bigcup_{j=1}^\infty Q_j - F\right) + m\left(\bigcup_{j=1}^\infty Q_j - E\right) \\ &\leq m\left(\bigcup_{j=n+1}^\infty Q_j\right) + m\left(\bigcup_{j=n+1}^\infty Q_j\right) - m(E) \\ &\leq \sum_{j=n+1}^\infty |Q_j| + \sum_{j=1}^\infty |Q_j| - m(E) \\ &\leq \epsilon. \end{aligned}$$

2.3.1 Invariance properties of Lebesgue measure

2.3.2 σ -algebra and Borel sets

Borel σ -algebra in \mathbb{R}^d , denoted by $B_{\mathbb{R}^d}$, is the smallest σ -algebra in \mathbb{R}^d that contains all open sets. Elements of this σ -algebra are called **Borel sets**. Since we observe that any intersection (not necessarily countable) of σ -algebra is again a σ -algebra, we may define $B_{\mathbb{R}^d}$ as the intersection of all σ -algebras that contain the open sets. This shows the existence and uniqueness of the Borel σ -algebra.

Remark. There exists Lebesgue measurable sets that are not Borel sets. (See Exercise @ref(exr: #35))

2.3.3 Construction of a non-measurable set

2.3.4 Axiom of choice

2.4 Measurable functions

2.4.1 Definition and basic properties

The starting point is the notion of a **characteristic function** of a set E , which is defined by

$$\chi_E(x) = \begin{cases} 1 & \text{if } x \in E, \\ 0 & \text{if } x \notin E. \end{cases}$$

For the Riemann integral it is in effect the class of **step functions** that build the blocks of integration theory, with each given as a finite sum

$$f = \sum_{k=1}^N a_k \chi_{R_k}$$

where each R_k is a rectangle, and the a_k are constants.

For the Lebesgue integral we need a more general notion. A **simple function** is a finite sum

$$f = \sum_{k=1}^N a_k \chi_{E_k}$$

where each E_k is a measurable set of finite measure, and the a_k are constants.

A function f defined on a measurable subset E of \mathbb{R}^d is **measurable**, if for all $a \in \mathbb{R}$, the set

$$f^{-1}([-\infty, a)) = \{x \in E : f(x) < a\}$$

is measurable. Note that this definition applies to extended-valued functions, so we use $f^{-1}([-\infty, a))$ instead of $f^{-1}((-\infty, a))$.

Proposition 2.3.

1. The finite-valued function f is measurable iff $f^{-1}(O)$ is measurable for every open set O , and iff $f^{-1}(F)$ is measurable for every closed set F . (Note that this property also applies to extended-valued functions, if we make the additional hypothesis that both $f^{-1}(\infty)$ and $f^{-1}(-\infty)$ are measurable sets since $[-\infty, a) = \{-\infty\} \cup (\bigcup_{n=1}^{\infty} (-n, a))$.)
2. If f is continuous on \mathbb{R}^d , then f is measurable. If f is measurable and finite-valued, and Φ is continuous, then $\Phi \circ f$ is measurable. (Note that it is not true that $f \circ \Phi$ is measurable. See exercise 35.)

Proof.

Φ is continuous, so $\Phi^{-1}((-\infty, a))$ is open set O .

3. Suppose $\{f_n\}_{n=1}^\infty$ is a sequence of measurable functions. Then

$$\sup_n f_n(x), \quad \inf_n f_n(x), \quad \limsup_{n \rightarrow \infty} f_n(x) \quad \text{and} \quad \liminf_{n \rightarrow \infty} f_n(x)$$

are measurable.

4. If $\{f_n\}_{n=1}^\infty$ is a collection of measurable functions, and

$$\lim_{n \rightarrow \infty} f_n(x) = f(x),$$

then f is measurable.

5. If f and g are measurable, then

- (i) The integer powers f^k , $k \geq 1$ are measurable.
- (ii) $f + g$ and fg are measurable if both f and g are finite-valued.

6. Suppose f is measurable, and $f(x) = g(x)$ for a.e. x . Then g is measurable.

2.4.2 Approximation by simple functions or step functions

Theorem 2.3.

Suppose f is a non-negative measurable function on \mathbb{R}^d . Then there exists an increasing sequence of non-negative simple functions $\{\varphi_n\}_{n=1}^\infty$ that converges pointwise to f , namely,

$$\varphi_k(x) \leq \varphi_{k+1}(x) \quad \text{and} \quad \lim_{k \rightarrow \infty} \varphi_k(x) = f(x), \quad \text{for all } x.$$

Proof.

Φ is continuous, so $\Phi^{-1}((-\infty, a))$ is open set O .

Theorem 2.4.

Suppose f is a measurable function on \mathbb{R}^d . Then there exists a sequence of simple functions $\{\varphi_n\}_{n=1}^\infty$ that satisfies

$$|\varphi_k(x)| \leq |\varphi_{k+1}(x)| \quad \text{and} \quad \lim_{k \rightarrow \infty} \varphi_k(x) = f(x), \quad \text{for all } x.$$

Proof.

We use the decomposition of f : $f(x) = f^+(x) - f^-(x)$.

Theorem 2.5.

Suppose f is measurable on \mathbb{R}^d . Then there exists an sequence of step functions $\{\psi_n\}_{n=1}^\infty$ that converges pointwise to $f(x)$ for almost every x ,

$$\varphi_k(x) \leq \varphi_{k+1}(x) \quad \text{and} \quad \lim_{k \rightarrow \infty} \varphi_k(x) = f(x), \quad \text{for all } x.$$

Proof.

By Theorem 2.4, it suffices to show that if E is a measurable set with finite measure, then $f = \chi_E$ can be approximated by step functions. This can be proven by split E into cubes and then rectangles with Theorem 2.2.

2.4.3 Littlewood's three principles

Theorem 2.6 (Egorov). Suppose $\{f_k\}_{k=1}^\infty$ is a sequence of measurable functions defined on a measurable set E with $m(E) < \infty$, and assume that $f_k \rightarrow f$ a.e. on E and f_1, f_2, \dots, f_k, f are finite valued a.e. on E . Given $\epsilon > 0$, we can find a closed set $A_\epsilon \subset E$ such that $m(E - A_\epsilon) \leq \epsilon$ and $f_k \rightarrow f$ uniformly on A_ϵ .

Remark. Note that f_1, f_2, \dots, f_k, f are finite valued a.e. on E . Indeed, if it is not satisfied, then we cannot construct $\{E_k^n\}_{k=1}^\infty$ such that $E_k^n \nearrow E$. A counterexample is that

$$f_k(x) = \begin{cases} k & |x| \leq k, \\ \infty & |x| > k \end{cases}$$

and $f(x) = \infty$ on \mathbb{R} .

Remark. Note that $m(E) < \infty$ and it is easy to construct counterexamples when $m(E) = \infty$. Indeed, if $m(E) = \infty$, then we cannot find k_n such that $m(E - E_{k_n}^n) < 1/2^n$ since $m(E - E_{k_n}^n) = m(E) - m(E_{k_n}^n)$.

Theorem 2.7 (Lusin). Suppose f is measurable and finite valued a.e. on E with E of finite measure. Then for every $\epsilon > 0$ there exists a closed set F_ϵ , with

$$F_\epsilon \subset E, \text{ and } m(E - F_\epsilon) \leq \epsilon$$

and such that $f|_{F_\epsilon}$ is continuous.

2.5 The Brunn-Minkowski inequality

Remark.

Let $a, b \geq 0$, then

$$\begin{aligned} (a+b)^\gamma &\geq a^\gamma + b^\gamma \text{ if } \gamma \geq 1, \\ (a+b)^\gamma &\leq a^\gamma + b^\gamma \text{ if } 0 < \gamma < 1 \end{aligned}$$

Proof.

Let $f(\gamma) = (1+x)^\gamma - (1+x^\gamma)$, where $x > 0$. Then

$$\begin{aligned} f'(\gamma) &= (1+x)^\gamma \ln(1+x) - x^\gamma \ln x \\ &= [(1+x)^\gamma - x^\gamma] \ln(1+x) + x^\gamma \ln(1+1/x) > 0, \end{aligned}$$

notice that $f(1) = 0$, so when $\gamma \geq 1$, $(1+x)^\gamma \geq (1+x^\gamma)$ and when $0 < \gamma < 1$, $(1+x)^\gamma < (1+x^\gamma)$. With this result, the original inequality is obvious.

??

Theorem 2.8. *Suppose A and B are measurable sets in \mathbb{R}^d and their sum $A+B$ is also measurable. Then*

$$m(A+B)^{1/d} \geq m(A)^{1/d} + m(B)^{1/d}.$$

2.6 Exercise

2.7 Problem

Chapter 3

Integration theory

3.1 The lebesgue integral: basic properties and convergence theorems

3.1.1 Stage one: simple functions

Proposition 3.1. *The integral of simple functions defined bve satisfies the following properties:*

1. *Independence of the representation. If $\varphi = \sum_{k=1}^N a_k \chi_{E_k}$ is any representation of φ , then*

$$\int \varphi = \sum_{k=1}^N a_k m(E_k).$$

2. *Linearity.*
3. *Additivity.*
4. *Monotonicity.*
5. *Triangle inequality. If φ is a simple function, then so is $|\varphi|$, and*

$$\left| \int \varphi \right| \leq \int |\varphi|.$$

3.1.2 Stage two: bounded functions supported on a set of finite measure

Lemma 3.1. *Let f be a bounded function supported on a set E of finite measure. If $\{\varphi_n\}_{n=1}^\infty$ is any sequence of simple functions bounded by M , supported on E , and with $\varphi_n(x) \rightarrow f(x)$ for a.e. x , then:*

1. The limit $\lim_{n \rightarrow \infty} \int \varphi_n$ exists.
2. If $f = 0$ a.e., then the limit $\lim_{n \rightarrow \infty} \int \varphi_n$ equals 0.

Proof.

Setting $I_n = \int \varphi_n$ and applying Egorov's theorem which is proven in Chapter 2 we have that for and large n and m

$$\begin{aligned}
 |I_n - I_m| &\leq \int_E |\varphi_n - \varphi_m| \\
 &= \int_{A_\epsilon} |\varphi_n - \varphi_m| + \int_{E-A_\epsilon} |\varphi_n - \varphi_m| \\
 &\leq \int_{A_\epsilon} \epsilon \, dx + \int_{E-A_\epsilon} 2M \, dx \\
 &\leq m(E)\epsilon + 2M\epsilon.
 \end{aligned}$$

given any $\epsilon > 0$. This proves that $\{I_n\}$ is a Cauchy sequence and hence converges. If $f = 0$, letting m tend to infinity we have $|I_n - f| = |I_n| \leq m(E)\epsilon + 2M\epsilon$, which yields $\lim_{n \rightarrow \infty} I_n = 0$.

For a bounded function f that is supported on sets of finite measure, we define its **Lebesgue integral** by

$$\int f = \lim_{n \rightarrow \infty} \int \varphi_n.$$

where $\{\varphi_n\}$ is any sequence of simple functions satisfying: $|\varphi_n| \leq M$, each φ_n is supported on the support of f , and $\varphi_n(x) \rightarrow f(x)$ for a.e. x as n tends to infinity.

Next, we must show that $\int f$ is independent of the limiting sequence $\{\varphi_n\}$ used, in order for the integral to be well-defined. Suppose that $\{\psi_n\}$ is another sequence of simple functions that satisfies the properties above. Then, if $\eta_n = \varphi_n - \psi_n$, the sequence $\{\eta_n\}$ consists of simple functions bounded by $2M$, supported on a set of finite measure, and such that $\eta_n \rightarrow 0$ a.e. as n tends to infinity. Applying the lemma we find

$$\lim_{n \rightarrow \infty} \int \varphi_n = \lim_{n \rightarrow \infty} \int \psi_n + \lim_{n \rightarrow \infty} \int \eta_n = \lim_{n \rightarrow \infty} \int \psi_n$$

as desired.

Proposition 3.2. *Suppose f and g are bounded functions supported on sets of finite measure. Then the following properties hold.*

1. *Linearity*
2. *Additivity*
3. *Monotonicity*

4. Triangle inequality

Theorem 3.1 (Bounded convergence theorem).

Suppose that $\{f_n\}$ is a sequence of measurable functions that are all bounded by M , are supported on a set E of finite measure, and $f_n(x) \rightarrow f(x)$ a.e. x as $n \rightarrow \infty$. Then f is measurable, bounded, supported on E for a.e. x , and

$$\int |f_n - f| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Consequently,

$$\int f_n \rightarrow \int f \text{ as } n \rightarrow \infty.$$

Proof.

The proof is a reprise of the argument in Lemma 3.1. Given $\epsilon > 0$, we may find, by Egorov's theorem,

$$\begin{aligned} \int |f_n - f| &\leq \int_{A_\epsilon} |f_n - f| + \int_{E-A_\epsilon} |f_n - f| \\ &\leq m(E)\epsilon + 2M\epsilon. \end{aligned}$$

for all large n .

3.1.3 Return to Riemann integrable functions

3.1.4 Stage three: non-negative functions

3.1.5 General form

Chapter 4

Methods

We describe our methods in this chapter.

Math can be added in body using usual syntax like this

4.1 math example

p is unknown but expected to be around $1/3$. Standard error will be approximated

$$SE = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{1/3(1-1/3)}{300}} = 0.027$$

You can also use math in footnotes like this¹.

We will approximate standard error to 0.027^2

¹where we mention $p = \frac{a}{b}$

² p is unknown but expected to be around $1/3$. Standard error will be approximated

$$SE = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{1/3(1-1/3)}{300}} = 0.027$$

Chapter 5

Applications

Some *significant* applications are demonstrated in this chapter.

5.1 Example one

5.2 Example two

Chapter 6

Final Words

We have finished a nice book.