

# Assignment 01

## A posteriori probability

Let  $P(X|v = u)$  denote the probability of drawing out  $X$  blacks in a total of  $N = 10$  draws from urn  $v$ . Obviously, the probability of drawing out a black each time from urn  $v$  is

$$p_v \equiv P(\text{color} = \text{black}|v = u) = \frac{u}{10}$$

Considering the process of drawing 10 balls to be a Bernoulli distribution, we have the probability

$$\begin{aligned} P(X = 3|v = u) &= C_{10}^3 \times p_v^3 \times (1 - p_v)^7 \\ &= 120 \times \left(\frac{u}{10}\right)^3 \times \left(1 - \frac{u}{10}\right)^7 \end{aligned}$$

Applying Bayesian law we have

$$P(v = u|X = 3) = \frac{P(X = 3|v = u) \times P(v = u)}{P(X = 3)}$$

where  $P(v = u) = \frac{1}{11}$ , and  $P(X = 3)$  can be calculated as

$$P(X = 3) = \frac{1}{11} \sum_{i=0}^{10} P(X = 3|v = i)$$

Applying the figures we have results as follows:

$u$	$P(v = u X = 3)$
0	0
1	0.063073
2	0.221240
3	<b>0.293220</b>
4	0.236256
5	0.128779
6	0.046668
7	0.009892
8	0.000864
9	0.000010
10	0

where maximum probability occurs when  $u = 3$ .

## Computer Generation of Random Variables

(a)

Since  $Y = G(X)$ , we have  $P(Y \leq y) = P(G(X) \leq y)$ .

According to the question,  $F(t)$  must be monotone increasing and differentiable. Applying  $F(\cdot)$  to both sides we have

$$P(G(X) \leq y) = P(F(G(X)) \leq F(y)) = P(X \leq F(y))$$

Because  $X \sim \text{Uniform}(0, 1)$ , its CDF takes the form

$$P(X \leq x) = x = F(y), \quad \forall u \in [0, 1)$$

Thus


$$P(Y \leq y) = F(y)$$


Applying the derivative to both side we have

$$p_Y(y) = p(y)$$

(b)

For  $N = 10^4$ , results are as follows:





For  $N = 10^6$ , we have:

