

of the anomaly observed by most theoretical approximations within TDDFT and Fermi liquid theories [7]. A strong failure of approximations based on TDDFT with a static exchange-correlation kernel is the lack of a damping mechanism which results in an infinite lifetime of plasmons for a region of wavevectors smaller than the critical wavevector that separates plasmonic and particle-hole excitations [7, 8]. The negative plasmon dispersion in low-density alkali metals, and in general the damping mechanism can be attributed to correlation effects or band structure [9–12]. There are pros and contras for both explanations in the literature [11, 12]. In heavier alkali metals such as Cs, electron transition to the near-Fermi-level $3d$ bands can occur [9, 10]. The transition energy of these electrons is comparable to the plasmon energy, potentially causing a negative dispersion. Additional corrections to the interacting response function or dielectric function can originate in a weak lattice potential and core polarization effects [7].

The random phase approximation (RPA) is a Greens function-based method that is often used to obtain the ground-state correlation energy of bulk and two-dimensional materials [13–17]. Although RPA relies on the linear response TDDFT framework, its excitation energies are inaccurate because of the overestimated short-range exchange-correlation effects [18]. The exact exchange-correlation kernel f_{xc} that would provide these effects is a functional derivative of the exchange-correlation potential. RPA is often interpreted within DFT to have roots in the adiabatic connection fluctuation dissipation theorem [19]. The bare RPA $f_{xc} = 0$ without a band structure does not yield a negative dispersion in heavy alkali metals [9]. Theoretical predictions from the late 80's in polycrystalline metals showed negative dispersion for Cs beyond RPA when the band structure was included [20]. Some investigations of the role of correlation in the plasmon dispersion discuss kernels, but most tests consider the exchange-correlation effects at the level of adiabatic local density functional approximation (ALDA) only. The work of Tatarczyk et al. [20] steps beyond this limitation to some extent by considering some more model kernels based either on the uniform electron gas paradigm or on other constraints.

In our current work we aim to fill the gap in analyzing the plasmon dispersion with recent exchange-correlation kernels, beyond the early ones developed in the 90's. Our work aims to go beyond a simple analysis of exchange-correlation effects with kernels. The major

$$f_{xc}^{ALDA}(q \rightarrow 0, \omega = 0) = -\frac{4\pi A}{k_F^2} \quad (2)$$

with $A = \frac{1}{4} - \frac{k_F^2}{4\pi} \frac{d^2(n\varepsilon_c)}{dn^2}$, where $k_F = (3\pi^2 n)^{1/3}$ is the Fermi wavevector and ε_c the correlation energy per particle of the uniform electron gas. $A = \frac{1}{4}$ belongs only to the exchange-only ALDA, while the density-dependent term in A gives the correlation beyond the high-density limit.

The real-space representation of ALDA is a delta function which indicates the spatial locality of this kernel. ALDA gives reasonable accuracy for low-frequency, long-wavelength excitations, but is not the right choice for a general correction to RPA [25]. The ALDA kernel was applied to the ground state correlation energy of the uniform electron gas but makes an error of ~ 0.5 eV [26]. This error is the same in magnitude but of opposite sign to the error that RPA makes for the same system.

The ALDA kernel can be made nonlocal, by applying a cutoff that makes the exchange-correlation kernel cancel the Hartree kernel for $q > 2k_{cut}$. The cutoff is introduced by the renormalized ALDA (rALDA) expression [27]

$$f_{xc}^{rALDA}(n, q) = -[\theta(k_{cut} - q) \frac{4\pi}{k_{cut}^2} + \theta(q - k_{cut}) \frac{4\pi}{q^2}] \quad (3)$$

with the cutoff wavevector $k_{cut} = \frac{k_F}{\sqrt{A}}$. By construction the rALDA_{xc} kernel keeps the correct $q \rightarrow 0$ limit of ALDA, but improves the wrong $q \rightarrow \infty$ behavior of ALDA_{xc}. For inhomogeneous systems, more ingredients like the density gradient or the kinetic energy density give more flexibility for kernels, as for ground state density functional approximations. The kinetic energy density is one of the ingredients of the nonlocal energy optimized (NEO) exchange-only kernel [28].

The NEO kernel improves the ground state correlation energy and structural properties beyond RPA. The NEO kernel is designed to satisfy further physical constraints beyond both ALDA or rALDA, and has the form

(28)

of real systems

$$f_x^{NEO} = -\frac{4\pi}{2q^2} [1 - e^{-\beta q^2/k_F^2}] \quad (4)$$

where $\beta = \frac{1}{4\tilde{c}(1-z^2)}$. The one-electron limit is satisfied by the ingredient $z = \frac{\tau^w}{\tau}$, where τ^w is the one-electron kinetic energy density [29] and τ the positive kinetic energy density constructed from the Kohn-Sham orbitals. In the uniform electron gas, z is zero, when $q \rightarrow 0$, the NEO kernel is properly independent of q for the uniform electron gas. The parameter \tilde{c} has a key relevance to the current work. In the construction of NEO \tilde{c} is designed to give a correction to the RPA correlation energy in the high-density limit. In other words, the standard $\tilde{c} = 0.264$ parameter in NEO provides a unique fit to the exact second-order correlation energy for the spin-unpolarized electron gas. The “second-order exchange” contribution to the second-order correlation energy of the uniform gas is the correction to direct RPA from wavefunction anti-symmetry that survives in the high-density limit. It can be evaluated from explicit expressions given by von Barth and Hedin for RPA [30] and by Langreth and Perdew [18] beyond RPA.

β can be chosen to satisfy another constraint relevant to the long-wavelength ($q \rightarrow 0$) limit. This choice is also made in the ALDA, rALDA, and in the CP07 dynamic exchange-correlation kernel constructed by Constantin and Pitarke [31]. The compressibility sum rule, [32] as

$$f_{xc}(n; q \rightarrow 0, \omega = 0) = \frac{d^2}{dn^2}[n\varepsilon_{xc}(n)], \quad (5)$$

is an important requirement for frequency-dependent exchange-correlation kernels. Satisfying the compressibility sum rule β becomes

~~$$\beta = -\frac{2k_F^2}{4\pi} \frac{d^2}{dn^2}[n\varepsilon_{xc}(n)] = A = \frac{-4\pi}{2k_F^2} \frac{1}{\tilde{c}} = \frac{-2k_F^2}{4\pi} \frac{d^2}{dn^2}[n\varepsilon_{xc}(n)]$$~~ (6)

or $\tilde{c} = 0.5$ in the high-density limit.

Thus the energy optimization of $\tilde{c} = 0.264$ in the high-density limit is different from the value $\tilde{c} = 0.5$ that yields the correct small- q kernel in the high-density limit. In the next section we will extensively discuss the impact of these physical constraints on the plasmon dispersion and provide a novel insight about the role of correlation effects. It will turn out that the difference between the ALDA and NEO kernels is important for the correlation

resolution electron energy loss (EELS) experiments indicate that the plasmon dispersion in heavy alkali metals such as Rb and Cs becomes negative, i.e., the plasmon frequency decreases with increasing wavevector q [7]. Since all the above experiments and calculations were performed for periodic crystals, it is difficult to decouple correlation and band structure effects [33] in the decay of the plasmon excitations.

Further calculations by Ku and Eguiluz in K crystal [10] indicate the relevance of band structure versus correlation and demonstrate that the exchange-correlation effects beyond RPA at the ALDA level have only a minor role in the decay of plasmons. A similar observation was made by Quong and Eguiluz [34] for Na crystal. Although crystal periodicity is important when modeling realistic conditions, the jellium model can offer an important way to separate the impact of many-body correlations and band structure. Within our work we want to provide an additional theoretical and numerical insight only about the many-body correlation effects and explain why in general no beyond-RPA approximation for jellium can predict the correct plasmon decay and lifetime for Cs. The justification of our results is based merely on exact physical constraints imposed on the construction of beyond-RPA approximations. By using the jellium model for alkali metals, we can build upon the conclusion that ALDA gives a minor improvement beyond RPA for the plasmon dispersion. With the possession of nonlocal exchange-correlation kernels developed since the later 90's, we can make further conclusions about how these recent approximations compare to ALDA and RPA in terms of correlation effects.

TDDFT has become a promising and affordable tool to assess the optical properties of solids. Most of the tests on optical properties still rely on the ALDA approximation. The shortcomings of ALDA become obvious for ground state correlation energies and for excitations of wide-frequency range in inhomogeneous systems. Relying on the observations of Refs [10, 34], we can keep the ALDA as our reference method when we compare the plasmon dispersion obtained from other exchange-correlation kernels to ALDA.

We can make two groups of assessed approximations. The first group includes exchange-only and exchange-correlation kernels based on the ALDA approximations. ALDAx and ALDAxc are both local kernels but differ in correlation contribution. rALDAx and rALDAxc

are nonlocal. The second group consists of NEO exchange-only kernels [28] with the \tilde{c} parameter constructed by satisfying different physical constraints. The default version of the NEO kernel yields the exact correction to the RPA correlation energy of jellium in the high-density limit:

$$e_c^{2X} = \frac{3}{8\pi^3} \int_0^\infty dQ Q^2 \tilde{G}_x(Q) \int_0^\infty dW Q^2 \{2\beta(Q, W)\}^2, \quad (7)$$

where $\tilde{G}_x(Q)$ refers to the kernel, according to the correspondence between kernels and local-field factors. $Q = \frac{q}{2k_F}$ is a dimensionless wavevector, and $W = \frac{\Omega}{2k_F^2}$ is a dimensionless frequency, to obtain the explicit expression for the second-order exchange energy e_c^{2X} by Langreth and Perdew [18] beyond RPA using the RPA correlation energy for the uniform electron gas given by von Barth and Hedin. The \tilde{c} parameter that corresponds to the second-order correlation energy was found to be 0.264.

delete

The exchange-only NEO kernel for the uniform electron gas at the long-wavelength limit is

$$f_x^{NEO}(n; q \rightarrow 0) = -\frac{4\pi}{2k_F^2} \frac{1}{4\tilde{c}} \quad (8)$$

Alternatively, in the long-wavelength limit we can use the compressibility sum rule formulated as $\frac{d^2}{dn^2} [n\varepsilon_{xc}(n)]$ to determine “ \tilde{c} ”. Then the c parameter of NEO can be estimated from the compressibility sum rule of the ALDAxc expression. This fitting delivers a different $\tilde{c} \cong 0.44$ at metallic densities. While formally the NEO approximation remains an exchange-only kernel, this kind of fitting to an exact physical condition with an exchange-correlation approach brings the correlation effects relevant for metallic densities into our NEO kernel [31].

$\tilde{c} = 0.43 - 0.47$ at metallic densities considered here.

Clearly “ \tilde{c} ” controls the correlation or screening within the NEO approximation starting from $\tilde{c} \rightarrow \infty$ in RPA. The impact of “ \tilde{c} ” as a screening parameter on ground state correlation energies was established by Bates et al in 2016. Changing “ \tilde{c} ” from its default 0.264 value was shown to yield different correlation energies in the uniform electron gas at $r_s=4$.

The exchange-only NEO kernel can be explicitly turned into an exchange-correlation ap-

proach by replacing “ \tilde{c} ” by an electron-density-dependent parameter. This approach was tested for jellium slab correlation energies at moderate densities, and resulted in improved integrated correlation energy [35]. For a given density, the density dependence can make “ \tilde{c} ” significantly smaller than its default value. In our analysis we investigate the effect of a low “ \tilde{c} ” parameter on the plasmon dispersion of various NFE metals. For testing purposes we choose $\tilde{c} = 0.0037$. Notice that this choice of \tilde{c} represents an unphysically low density according to Eq. (13) of Ref. 35.

At first we discuss the plasmon dispersion up to the wavevector region where plasmons decay into single particle excitations. We consider all the exchange-correlation kernels described above. Within the static approximation for the kernel $f_{xc}(q)$, the plasmon frequency $\omega_p(q)$ is found by solving the equation $\epsilon(q, \omega) = 1 - (v_c(q) + f_{xc}(q))\chi_0(q, \omega) = 0$, for $\lambda = 1$,¹⁷ and the solutions are undamped outside the particle-hole continuum, i.e., for $q < q_c$ where $\frac{\omega_p(q_c)}{\varepsilon_F} = 2 \left(\frac{q_c}{k_F} \right) + \left(\frac{q_c}{k_F} \right)^2$. For the r_s values considered here, $q_c/k_F \approx 1 \ll k_{cut}/k_F \approx 2$, where k_{cut} is the cutoff wavevector for a kernel. Thus rALDA and ALDA kernels will yield the same plasmon dispersion.

Al is a metal with rather high density, and RPA becomes relatively exact in the high-density limit [34]. Here, we model Al by jellium with $r_s = 2.07$. The small wavevector behavior is demonstrated by the correct plasmon energy known from an EELS experiment [36]. All ALDA and rALDA kernels return the correct long-wavelength limit of RPA. The $\tilde{c} \approx 0.47$ NEO exchange kernel keeps the correct long-wavelength of RPA. Since the compressibility sum rule delivers direct information about the long-wavelength limit, among all the approximations NEO $\tilde{c} \approx 0.47$ has a direct impact on the curvature of the dispersion. The fitting against this exact constraint designates that the plasmon dispersion must start out as horizontal at small wavevectors. NEO $\tilde{c} \approx 0.47$ exemplifies the best behavior in the long wavelength limit ($q \rightarrow 0$) that any static kernel for jellium at $r_s = 2.07$ can demonstrate.

Comparing the ALDA and rALDA kernels in the left panel of Fig. 2, it is apparent that the nonlocal feature of the rALDA is much relevant in the ground state correlation energy does not change the plasmon dispersion, and all ALDA and rALDA approximations yield the same dispersion curve. The NEO $\tilde{c} = 0.264$ and $\tilde{c} = 0.47$ methods basically agree, while

the NEO $\tilde{c}=0.0037$ is completely unphysical with low plasmon frequencies as an indication of the lack of exact constraints (See Fig. 2).




FIG. 2. The plasmon dispersion for Al (modeled as jellium with $r_s = 2.07$) up to the critical wavevector. The left panel shows the dispersion obtained with RPA and beyond-RPA with ALDAx, ALDAXc, rALDAx and rALDAXc approximations. The right panel shows the dispersion from RPA and the three NEO approximations with the \tilde{c} parameters corresponding to different choices.

For jellium at $r_s=3.93$ (as for Na), all ALDA and rALDA dispersion curves barely differ in the left panel of Fig. 3. As in Al, there is no significant change coming from the nonlocal kernels. According to the right subfigure, the NEO $\tilde{c} = 0.264$ and NEO $\tilde{c} = 0.44$ kernels differ more than they do in Al for a range of ~ 0.5 for dimensionless wavevector Q . NEO $\tilde{c} = 0.0037$ leads to unphysically low plasmon frequencies.

Cs is the alkali metal with the lowest density [37]. We modeled it here as jellium with $r_s = 5.62$. This characteristic manifests itself in the plasmon dispersion when comparing the approximations in the left and right panels of Fig. 4. Being correct at small q , the ALDAXc and rALDAXc are more suitable for lower densities in Cs, however, the nonlocality versus locality in rALDAXc and ALDAXc does not much affect the dispersion. Comparing the NEO approximations, NEO $\tilde{c} = 0.264$ results in more correction beyond-RPA than it does in the previous two metals. Furthermore NEO $\tilde{c} = 0.264$ yields more correction in the plasmon frequencies than any of the ALDA and rALDA kernels. The exact constraint-based NEO $\tilde{c} = 0.43$ behaves more like ALDAx or ALDAXc. NEO $\tilde{c} = 0.0037$ considerably lowers the plasmon dispersion against the other approximations. At the first glance this could be the

correlation kernels and RPA are properly horizontal at small Q wavevectors and become properly quadratic as Q increase. The horizontal line is a consequence of the exact physical constraints satisfied by these methods. NEO $\tilde{c} = 0.0037$ is not consistent with the exact constraint of the compressibility sum rule. While NEO $\tilde{c} = 0.264$ is based on the exact high-density limit of the correlation energy, the unphysical NEO deviates from this constraint and picks up a wrong negative quadratic dispersion.




FIG. 5. The long wavelength behavior of all approximations considered in this work.

We have also extracted the plasmon dispersion for the real metals Na and Cs by calculating the electron energy loss spectrum using the GPAW[38–41] code. The results are presented in Figure 6. We used the projector-augmented wave (PAW)[42] pseudopotential provided with the GPAW code, an energy cutoff of 600 eV, and a $16 \times 16 \times 16$ k-point mesh to sample the Brillouin zone. Our calculations including bandstructure effects confirm that the band structure significantly alters the plasmon dispersion for Cs, while its effect is negligible for Na. The plasmon frequency at $q \rightarrow 0$ shows a 0.5 eV renormalization compared to the RPA value within the jellium model. The effect of the band structure in Cs is significantly large enough to dominate over the changes from one kernel to another.




FIG. 6. The plasmon dispersion of Na (left) and Cs (right) with RPA and some exchange-correlation kernels within the jellium model. For Cs, the dispersion with RPA is also displayed with band-structure effects obtained from the GPAW code, showing results close to experiment [7] (also shown).

IV. THE DYNAMIC STRUCTURE FACTOR WITHIN AND BEYOND-RPA

The dynamic structure factor [19] or spectral function $S(\mathbf{q}, \omega)$ shows the distribution of frequencies ω for density fluctuations of wavevector \mathbf{q} in the ground state of the uniform electron gas. Although it arises from all density fluctuations of a given wavevector, the spectral function for small wavevector typically peaks around the frequency of a plasmon. The inverse frequency width of this peak, by the uncertainty principle, provides a lower bound on the decay time for such a fluctuation, while the dependence of the peak on \mathbf{q} reflects the plasmon dispersion. Here we will investigate the effects of various model exchange-correlation kernels on the spectral function. The dynamic structure factor $S(\mathbf{q}, \omega)$ is proportional to $\text{Im}\chi$ [43], the loss component of the dynamic density-density response function:

$$S(\mathbf{q}, \omega) = -\frac{1}{\pi} \text{Im}\chi(\mathbf{q}, \omega) \Theta(\omega) \quad (9)$$

This quantity has been investigated by Lewis and Berkelbach [44] using an equation-of-motion coupled cluster singles and doubles formalism, which unlike our TDDFT methods allows for plasmon decay via multi-pair electron-hole decay channels. With our real static exchange-correlation kernels, the plasmon at small finite \mathbf{q} does not decay.

In Figure 7, we analyze our approximations at three wavevectors: $\mathbf{q} = 0.1k_F$, $\mathbf{q} = 0.5k_F$,

electron gas. We will now show that the range of q/k_F (less than or about equal to 2) that distributes to the correlation energy is much greater than the range of q/k_F (less than or about equal to 1) that contributes to the plasmon dispersion. In the smaller range (but not in the larger one), the ALDA_{xc} kernel is nearly sufficient, while the default NEO kernel is nearly sufficient over the larger range. This completes our plasmon dispersion analysis with an explanation why a negative dispersion cannot exist in jellium at the density of Cs.

Exchange-correlation kernels can be applied to improve the ground state correlation energy of RPA through the adiabatic connection fluctuation dissipation theorem. This is the basis of the wavevector decomposition of the ground state exchange-correlation energy as known from Langreth and Perdew [19]:

$$E_{xc} = \int \frac{d^3 q}{(2\pi)^3} \frac{1}{2} \int_0^1 \frac{d\lambda}{\lambda} \left(\frac{4\pi\lambda}{q^2} \right) N[S_\lambda(\mathbf{q}) - 1], \quad (10)$$

where λ is the coupling constant along the adiabatic connection path and $S_\lambda(\mathbf{q})$ is the static structure factor found by integrating $S_\lambda(\mathbf{q}, \omega)$ over frequency and dividing by the electron number. According to the expression given by Eq. 10, the exchange-correlation energy depends upon the dynamic structure factor. The exchange energy E_x replaces S_λ by S_0 , and the correlation energy is $E_c = E_{xc} - E_x$. Figure 8 shows the wavevector decomposition of the correlation energy for all exchange and exchange-correlation kernels considered here. We plot this for $r_s = 4$ and $r_s = 5.62$.

The physical basis of our analysis is the exact exchange-correlation kernel $f_{xc}(q, \omega)$ of the uniform electron gas. For the correlation energy, the static version of the kernel $f_{xc}(q, 0)$ can be applied to a good approximation [26]. Note that the frequency dependence at least qualitatively can also be ignored for $\omega \approx \omega_p$ [45, 46]. The ALDA exchange-correlation kernel $f_{xc}^{ALDA}(q, 0)$ approaches the exact kernel for the uniform electron gas at $q \rightarrow 0$. In the long wavelength limit,

$$\lim_{q \rightarrow 0} f_{xc}(q, 0) = \lim_{q \rightarrow 0} f_{xc}^{ALDA}(q, 0). \quad (11)$$

As we will see below, f_{xc}^{ALDA} breaks down for $q/2k_F \geq 0.5$, where our constraint-based NEO kernels become less negative and more accurate. Therefore, as suggested by Fig. 1:

$$f_{xc}^{ALDA}(q, 0) < f_{xc}(q, 0) < 0$$

(12) (11)

The ALDA approximation becomes a lower bound to the static exact exchange-correlation kernel for the correlation energy of the uniform electron gas. Figure 8 visualizes the relation between ALDAxc, RPA and some other exchange-correlation kernels. The ALDAxc is shown in the left panel of Figure 8. The ALDAc is very accurate for small wavevectors but starts to deviate from RPA at $Q \approx 0.25$. At $q = 2k_F$ the ALDAxc yields a strong overestimation of the correction to RPA correlation energy. All static beyond-RPA kernels make the exchange-correlation energy of RPA less negative for any density including $r_s = 5.62$. The correlation energies from NEO $\tilde{c} = 0.264$ and from the NEO kernel fitted against the compressibility sum-rule are close to each other but the unphysical NEO $\tilde{c} = 0.0037$ adds a much larger correction to the RPA correlation energy.

The constraints of Equations 11 and 12 control the plasmon dispersion and the correlation energy. The dynamic structure factor becomes a link that couples the physics in the correlation energy and plasmon dispersion. From the correlation energy, the exact uniform-electron gas-based kernel must be less negative than the ALDAxc kernel. The NEO $\tilde{c} = 0.264$ kernel is uniform electron gas-based only through its energy optimization to the high-density limit, and performs reasonably for both plasmon dispersion and correlation energy in jellium.




FIG. 8. Wavevector analysis of the ground state correlation-only energy of jellium from the dynamic structure factor for reduced wavevector $Q = \frac{q}{2k_F}$. The area under the curve is the correlation energy.

The left figure shows the correlation-only energy for RPA, ALDA and NEO with the three choices for \tilde{c} , for $r_s=4$. The right figure shows the same for $r_s=5.62$ corresponding to Cs.

The area under each curve is the correlation energy.




FIG. 8. Wavevector analysis of the ground state correlation-only energy of jellium from the dynamic structure factor for reduced wavevector $Q = \frac{q}{2k_F}$. The left figure shows the correlation-only energy for RPA, ALDA and NEO with the three choices for \bar{c} , for $r_s=4$. The right figure shows the same for $r_s=5.62$ corresponding to Cs.

which arises from bandstructure

the heavy alkali metal Cs. The current exchange-correlation kernels do not have an explicit density dependence that could have a larger impact. For the exact exchange-correlation kernel, the ALDAxc is likely a lower bound (as suggested by Fig. 1). The ALDAxc is accurate for $\frac{q}{k_F} < 1$, the range of q that determines the plasmon dispersion, even though the ALDAxc kernel fails badly for $\frac{q}{k_F} \gg 1$, a range that contributes significantly to the correlation energy.

VI. ACKNOWLEDGMENT

The research of N. K. N., S. R. and A. R. was supported by the National Science Foundation under Grant No.DMR-1553022.

* niraj.nepal@temple.edu

† aruzsinszky@temple.edu

- [1] G. Onida, L. Reining, and A. Rubio, *Reviews of Modern Physics* **74**, 601 (2002).

- [2] H. Raether, *Excitation of plasmons and interband transitions by electrons*, Vol. 88 (Springer, 2006).