

Multi-variable calculus

Álvaro Fernández Barrero and Vicente Gabriel Piñar Ojeda

16-09-2026

1 Multi-variable function

We have been integrating and differentiating over maps with a single variable, but nothing in their definition prevents them from having more than one. Therefore, we can define a map $\phi : \mathbb{F}^n \rightarrow \mathbb{F}^m$, being $n, m \in \mathbb{N}$. In multi-variable calculus, just like single-variable calculus, we ignore the possibility of having a complex number as either an income or outcome to simplify the math and because there is a whole branch in mathematics that already studies how they behave as they have some interesting and unique rules. Thus, we just work with maps in the form $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$.

As you might have noticed already, this notation is revealing that when a function requires up to one variable or even give several results, what is actually using are n -dimensional vectors.

2 Limits in several variables

3 Continuity in several variables

4 Partial derivatives

When it comes to multi-variable functions, differentiating them is not as trivial as single-variable maps since we have now several directions to go through. Thus, something we can do is picking one variable and analyze the function in that direction.

Mathematically, we can adjust the derivative's definition to satisfy the conditions I just said, popping up the partial derivatives.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function:

$$\frac{\partial f}{\partial x_\mu} := \lim_{h \rightarrow 0} \frac{f(x_1, x_2, \dots, x_\mu + h, \dots, x_n) - f(x_1, x_2, \dots, x_\mu, \dots, x_n)}{h}$$

Another way to write this is by expressing the arguments as a vector:

$$\frac{\partial f}{\partial x_\mu} = \lim_{h \rightarrow 0} \frac{f(\vec{v} + h\hat{v}_\mu) - f(\vec{v})}{h}$$

Where $\hat{v}_\mu \in \mathbb{R}^n$ is an unit vector pointing to the variable's direction. We can define it like $\hat{v}_{\mu\nu} = \delta_{\mu\nu}$, where $\delta_{\mu\nu}$ is the Kronecker's delta.

At this point, maybe you are wondering how to differentiate it and the truth is that the method does not change whatsoever: as we are only moving in one variable's direction, that is the one taken as variable and the rest become constants at sight of the derivative.

For this kind of derivative, we can bump into a few notations such as:

$$f_{x_\mu} = \partial_{x_\mu} f = \frac{\partial}{\partial x_\mu} f = \frac{\partial f}{\partial x_\mu}$$

4.1 The n-th partial derivative

In the same way we can take the n -th derivative of a function if neither the derivability nor the continuity breaks, we can do the same with partial derivatives but as there are more variables, the notation must change.

Let us go for the simplest case: the n -th derivative with respect to the same variable all of the times.

$$f_{x_\mu x_\mu x_\mu \dots x_\mu} = \partial_{x_\mu x_\mu x_\mu \dots x_\mu} f = \frac{\partial^n}{\partial x_\mu^n} f = \frac{\partial^n f}{\partial x_\mu^n}$$

The notation is exactly the same as the n -th derivative's notation for single-variable functions but adding the new notation the partial derivatives introduces.

In case we are differentiating with respect to different variables, the notation might be a little more tedious since we need to specify every single one.

$$f_{x_{\mu_1} x_{\mu_2} x_{\mu_3} \dots x_{\mu_n}} = \partial_{x_{\mu_1} x_{\mu_2} x_{\mu_3} \dots x_{\mu_n}} f = \frac{\partial^n}{\partial x_{\mu_1} \partial x_{\mu_2} \partial x_{\mu_3} \dots \partial x_{\mu_n}} f = \frac{\partial^n f}{\partial x_{\mu_1} \partial x_{\mu_2} \partial x_{\mu_3} \dots \partial x_{\mu_n}}$$

4.2 Mean value theorem

In the same way we have modified the derivative's definition to fit better with the multi-variable analysis requisites, we can use the same philosophy for the Lagrange's mean value theorem: we will just use one variable and work with it. Thus, let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function in $[a, b]$ and differentiable in (a, b) through the μ -th variable, $\exists c \in (a, b)$:

$$\frac{f(x_1, x_2, \dots, x_{\mu-1}, a, x_{\mu+1}, \dots, x_n) - f(x_1, x_2, \dots, x_{\mu-1}, b, x_{\mu+1}, \dots, x_n)}{a - b} = \frac{\partial f}{\partial x_\mu}(c)$$

4.3 Schwarz's theorem: Symmetry of the second derivative

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous twice with respect to the variables x_μ and x_ν . We can arbitrarily write:

$$\begin{aligned} & f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n) \\ & - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n) + f(x_1, \dots, 0, \dots, 0, \dots, x_n) \end{aligned}$$

It looks messy, and it is, but we are just taking the function normally, subtracting it with the function with x_μ evaluated in 0, we repeat it but this time with the variable x_ν evaluated in 0, and finally add the function with both of these variables evaluated in 0.

Here, we will name a variable $u : \mathbb{R}^2 \rightarrow \mathbb{R}$ as :

$$u(x_\mu, x_\nu) = f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n)$$

Thus, replacing in the previous formula, we get:

$$\begin{aligned} & f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n) \\ & - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n) + f(x_1, \dots, 0, \dots, 0, \dots, x_n) \\ & = u(x_\mu, x_\nu) - u(x_\mu, 0) \end{aligned}$$

Here it is obvious we can apply the mean value theorem since we have a subtraction of two functions with a difference in a variable. Therefore, by applying it:

$$u(x_\mu, x_\nu) - u(x_\mu, 0) = (x_\nu - 0) \frac{\partial u}{\partial x_\nu}(\xi_1) = x_\nu \frac{\partial u}{\partial x_\nu}(x_\mu, \xi_1)$$

We know what u is, we just defined it. Now, we will undo that change and rewrite the result in terms of f :

$$x_\nu \frac{\partial u}{\partial x_\nu}(x_\mu, \xi_1) = x_\nu \left(\frac{\partial f}{\partial x_\nu}(x_1, \dots, x_\mu, \dots, \xi_1, \dots, x_n) - \frac{\partial f}{\partial x_\nu}(x_1, \dots, 0, \dots, \xi_1, \dots, x_n) \right)$$

We have obtained anew a difference of two functions with a change in one of its variables, therefore we can use again the mean value theorem.

$$\begin{aligned} & x_\nu \left(\frac{\partial f}{\partial x_\nu}(x_1, \dots, x_\mu, \dots, \xi_1, \dots, x_n) - \frac{\partial f}{\partial x_\nu}(x_1, \dots, 0, \dots, \xi_1, \dots, x_n) \right) \\ & = x_\nu x_\mu \frac{\partial^2 f}{\partial x_\mu \partial x_\nu}(x_1, \dots, \xi_2, \dots, \xi_1, \dots, x_n) \end{aligned}$$

After this whole work, we finally reached a pretty nice formula, and now, we need to do the same again. By applying commutativity to the very first formula, we can state that:

$$\begin{aligned} & f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n) \\ & - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n) + f(x_1, \dots, 0, \dots, 0, \dots, x_n) \\ & = f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n) \\ & - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n) + f(x_1, \dots, 0, \dots, 0, \dots, x_n) \end{aligned}$$

Therefore, we can name another function v such as:

$$v(x_\mu, x_\nu) = f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n)$$

And, just how we just have done, rewrite the previous formula in terms of v :

$$\begin{aligned}
& f(x_1, \dots, x_\mu, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, x_\mu, \dots, 0, \dots, x_n) \\
& - f(x_1, \dots, 0, \dots, x_\nu, \dots, x_n) + f(x_1, \dots, 0, \dots, 0, \dots, x_n) \\
& = v(x_\mu, x_\nu) - v(0, x_\nu)
\end{aligned}$$

If you check this result, it is exactly the same as the result the u function gave me but the variable that before we set to 0 is now its value and the other is now 0. Again, we repeat the process using by the same logic the mean value theorem.

$$v(x_\mu, x_\nu) - v(0, x_\nu) = x_\mu \frac{\partial v}{\partial x_\mu}(\zeta_1, x_\nu)$$

We rewrite it in terms for f :

$$x_\mu \frac{\partial v}{\partial x_\mu}(\zeta_1, x_\nu) = x_\mu \left(\frac{\partial f}{\partial x_\mu}(x_1, \dots, \zeta_1, \dots, x_\nu, \dots, x_n) - \frac{\partial f}{\partial x_\mu}(x_1, \dots, \zeta_1, \dots, 0, \dots, x_n) \right)$$

And finally, we apply again the mean value theorem.

$$\begin{aligned}
& x_\mu \left(\frac{\partial f}{\partial x_\mu}(x_1, \dots, \zeta_1, \dots, x_\nu, \dots, x_n) - \frac{\partial f}{\partial x_\mu}(x_1, \dots, \zeta_1, \dots, 0, \dots, x_n) \right) \\
& = x_\mu x_\nu \frac{\partial^2 f}{\partial x_\nu \partial x_\mu}(x_1, \dots, \zeta_1, \dots, \zeta_2, \dots, x_n)
\end{aligned}$$

Now, since we first stated that the two starting points to derive the last two formulas are the same, we can write that both results are the same:

$$x_\nu x_\mu \frac{\partial^2 f}{\partial x_\mu \partial x_\nu}(x_1, \dots, \xi_2, \dots, \xi_1, \dots, x_n) = x_\mu x_\nu \frac{\partial^2 f}{\partial x_\nu \partial x_\mu}(x_1, \dots, \zeta_1, \dots, \zeta_2, \dots, x_n)$$

By dividing both sides by $x_\mu x_\nu$:

$$\frac{\partial^2 f}{\partial x_\mu \partial x_\nu}(x_1, \dots, \xi_2, \dots, \xi_1, \dots, x_n) = \frac{\partial^2 f}{\partial x_\nu \partial x_\mu}(x_1, \dots, \zeta_1, \dots, \zeta_2, \dots, x_n)$$

Finally, we just derived that if the function behaves and it is continuous in the first and second derivative with respect to these two random variables, it actually does not matter the order of differentiation

$$\frac{\partial^2 f}{\partial x_\mu \partial x_\nu} = \frac{\partial^2 f}{\partial x_\nu \partial x_\mu}$$

5 Leibniz's rule: Differentiation under the integral sign

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a continuous function in the interval $[a, b]$, where a and b are two continuous functions $a : \mathbb{R}^m \rightarrow \mathbb{R}$, $b : \mathbb{R}^p \rightarrow \mathbb{R}$:

$$\frac{d}{dt} \int_{a(a_1, \dots, t, \dots, a_m)}^{b(b_1, \dots, t, \dots, b_p)} f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu$$

This is the most general way of writing the integral. We will solve this derivative by using the derivative's definition.

$$\begin{aligned} & \frac{d}{dt} \int_{a(a_1, \dots, t, \dots, a_m)}^{b(b_1, \dots, t, \dots, b_p)} f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu \\ &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_{a(a_1, \dots, t+\Delta t, \dots, a_m)}^{b(b_1, \dots, t+\Delta t, \dots, b_p)} f(x_1, \dots, t+\Delta t, \dots, x_\nu, \dots, x_n) dx_\nu \right. \\ & \quad \left. - \int_{a(a_1, \dots, t, \dots, a_m)}^{b(b_1, \dots, t, \dots, b_p)} f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu \right) \end{aligned}$$

First, we will rename a little the terms to simplify the notation and be able to comprehend what is afoot in every step without getting lost by it.

We will say that:

- $a = a(a_1, \dots, t, \dots, a_m)$
- $b = b(b_1, \dots, t, \dots, b_p)$
- $\Delta a = a(a_1, \dots, t + \Delta t, \dots, a_m) - a(a_1, \dots, t, \dots, a_m)$
- $\Delta b = b(b_1, \dots, t + \Delta t, \dots, b_p) - b(b_1, \dots, t, \dots, b_p)$
- $u(t, x_\nu) = f(x_1, \dots, t, \dots, x_\nu, \dots, x_n)$

Thus:

$$\begin{aligned} & \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_{a(a_1, \dots, t+\Delta t, \dots, a_m)}^{b(b_1, \dots, t+\Delta t, \dots, b_p)} f(x_1, \dots, t+\Delta t, \dots, x_\nu, \dots, x_n) dx_\nu \right. \\ & \quad \left. - \int_{a(a_1, \dots, t, \dots, a_m)}^{b(b_1, \dots, t, \dots, b_p)} f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu \right) \\ &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_{a+\Delta a}^{b+\Delta b} u(t+\Delta t, x_\nu) dx_\nu - \int_a^b u(t, x_\nu) dx_\nu \right) \end{aligned}$$

If we analyze the first integral, we can state that:

$$\begin{aligned} & \int_{a+\Delta a}^{b+\Delta b} u(t+\Delta t, x_\nu) dx_\nu \\ &= \int_a^b u(t+\Delta t, x_\nu) dx_\nu - \int_a^{a+\Delta a} u(t+\Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t+\Delta t, x_\nu) dx_\nu \end{aligned}$$

Therefore, we can write it on the formula to get the following result:

$$\begin{aligned}
& \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_{a+\Delta a}^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu - \int_a^b u(t, x_\nu) dx_\nu \right) \\
&= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_a^b u(t + \Delta t, x_\nu) dx_\nu - \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu \right. \\
&\quad \left. - \int_a^b u(t, x_\nu) dx_\nu \right)
\end{aligned}$$

As the integral is a linear operator, we can use its properties to gather together the two remaining integrals with the same limits as:

$$= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \left(\int_a^b [u(t + \Delta t, x_\nu) - u(t, x_\nu)] dx_\nu - \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu \right)$$

For the very first integral, we can change the function u by its definition, getting back the original function f and distribute the $1/\Delta t$ term:

$$\begin{aligned}
&= \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{f(x_1, \dots, t + \Delta t, \dots, x_\nu, \dots, x_n) - f(x_1, \dots, t, \dots, x_\nu, \dots, x_n)}{\Delta t} dx_\nu \right. \\
&\quad \left. + \frac{1}{\Delta t} \left[- \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu \right] \right)
\end{aligned}$$

As we see, we have the partial derivative's definition of f with respect to t .

$$= \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu + \frac{1}{\Delta t} \left[- \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu \right] \right)$$

For the rest of the terms, we need to use the mean value theorem. Let us take the third integral and say that U is a primitive of u .

$$\int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu = (b + \Delta b - b) \frac{\partial U}{\partial x_\nu}(t + \Delta t, \zeta) = \Delta b u(t + \Delta t, \zeta)$$

We can use the same logic for the second integral:

$$- \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu = -(a + \Delta a - a) \frac{\partial U}{\partial x_\nu}(t + \Delta t, \zeta) = -\Delta a u(t + \Delta t, \xi)$$

Using these results on the formula:

$$\begin{aligned}
& \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu + \frac{1}{\Delta t} \left[- \int_a^{a+\Delta a} u(t + \Delta t, x_\nu) dx_\nu + \int_b^{b+\Delta b} u(t + \Delta t, x_\nu) dx_\nu \right] \right) \\
&= \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu + \frac{1}{\Delta t} [-\Delta a u(t + \Delta t, \xi) + \Delta b u(t + \Delta t, \zeta)] \right)
\end{aligned}$$

Here, as Δt tends to 0, $t + \Delta t$ is clear that tends to t , but since ξ and ζ are defined by the mean value theorem, ξ tends to a and ζ to b .

$$= \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu + \frac{1}{\Delta t} [-\Delta a u(t, a) + \Delta b u(t, b)] \right)$$

After applying those changes, we can write the u in terms of f to get the original function back everywhere and distribute $1/\Delta t$.

$$= \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu - \frac{\Delta a}{\Delta t} f(x_1, \dots, t, \dots, a, \dots, x_n) + \frac{\Delta b}{\Delta t} f(x_1, \dots, t, \dots, b, \dots, x_n) \right)$$

As Δt goes to zero, we can check what happens with the fractions we have left.

$$\lim_{\Delta t \rightarrow 0} \frac{\Delta a}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{a(a_1, \dots, t + \Delta t, \dots, a_n) - a(a_1, \dots, t, \dots, a_n)}{\Delta t} = \frac{\partial a}{\partial t}$$

$$\lim_{\Delta t \rightarrow 0} \frac{\Delta b}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{b(b_1, \dots, t + \Delta t, \dots, b_n) - b(b_1, \dots, t, \dots, b_n)}{\Delta t} = \frac{\partial b}{\partial t}$$

Hence:

$$\begin{aligned} & \lim_{\Delta t \rightarrow 0} \left(\int_a^b \frac{\partial f}{\partial t} dx_\nu - \frac{\Delta a}{\Delta t} f(x_1, \dots, t, \dots, a, \dots, x_n) + \frac{\Delta b}{\Delta t} f(x_1, \dots, t, \dots, b, \dots, x_n) \right) \\ &= \int_a^b \frac{\partial f}{\partial t} dx_\nu - \frac{\partial a}{\partial t} f(x_1, \dots, t, \dots, a, \dots, x_n) + \frac{\partial b}{\partial t} f(x_1, \dots, t, \dots, b, \dots, x_n) \end{aligned}$$

Therefore, we just have proven that:

$$\begin{aligned} & \frac{d}{dt} \int_{a(a_1, \dots, t, \dots, a_n)}^{b(b_1, \dots, t, \dots, b_n)} f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu \\ &= \int_a^b \frac{\partial f}{\partial t} dx_\nu - \frac{\partial a}{\partial t} f(x_1, \dots, t, \dots, a, \dots, x_n) + \frac{\partial b}{\partial t} f(x_1, \dots, t, \dots, b, \dots, x_n) \end{aligned}$$

If it happens to the functions a and b to be a couple of single-variable maps with which variable is precisely the variable we are differentiating the integral, we can convert their partial derivatives to normal single-variable derivatives as:

$$\begin{aligned} &= \int_a^b \frac{\partial f}{\partial t} dx_\nu - \frac{da}{dt} f(x_1, \dots, t, \dots, a, \dots, x_n) + \frac{db}{dt} f(x_1, \dots, t, \dots, b, \dots, x_n) \\ &\iff a : \mathbb{R} \rightarrow \mathbb{R} \wedge b : \mathbb{R} \rightarrow \mathbb{R} \end{aligned}$$

Moreover, if it happens that a and b are just constants, since the derivative of a constants is always 0, some terms here vanishes leaving the result of:

$$\frac{d}{dt} \int_a^b f(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu = \int_a^b \frac{\partial f}{\partial t}(x_1, \dots, t, \dots, x_\nu, \dots, x_n) dx_\nu, \quad a, b \in \mathbb{R}$$

6 Directional derivative

When we are differentiating a multi-variable function, we can just use partial derivatives to differentiate with respect to a specific variable. Another way to look at it is by comprehending that we are taking a direction in an axis and differentiating the function thru that way. Taking this thinking, there are many more paths to differentiate thru than just the canonical axes and there is when the directional derivative pops up.

To commence, we need to know what path we want to take and for that we need to parameterize it.

Let $a, u \in \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuous. We define the directional derivative as:

$$D_u f := \lim_{t \rightarrow 0} \frac{f(a + tu) - f(a)}{t}$$

Here, the term $a + tu$ is a line parameterized in the u direction taking a as displacement. We will define a few more vectors as:

$$p_\mu = \begin{pmatrix} a_1 + tu_1 \\ a_2 + tu_2 \\ \vdots \\ a_\mu + tu_\mu \\ a_{\mu+1} \\ \vdots \\ a_n \end{pmatrix}, \quad \mu \in \{x \in \mathbb{N}, 1 \leq x \leq n\},$$

Here, we are defining a vector p_μ whose components $\leq \mu$ are their respective component in the vector $a + tu$ but the rest of components are their respective in a . By that, we can establish that $p_0 = a \wedge p_n = a + tu$. Therefore, we will take the nominator of the limit and rewrite it in terms of p_μ .

$$f(a + tu) - f(a) = f(p_n) - f(p_0)$$

This will allow us to telescope it and have a pretty neat serie:

$$f(a + tu) - f(a) = f(p_n) - f(p_0) = \sum_{\mu=1}^n (f(p_\mu) - f(p_{\mu-1}))$$

By dividing both sides by t :

$$\frac{f(a + tu) - f(a)}{t} = \sum_{\mu=1}^n \frac{f(p_\mu) - f(p_{\mu-1})}{t}$$

We have done it because we could make appear on the left-hand part the fraction from the definition we used for the directional derivative, obtaining a new way to express it.

Now, we will define another set of functions $g_\mu : \mathbb{R} \rightarrow \mathbb{R}$ such that:

$$g_\mu(s) = f(a_1 + tu_1, a_2 + tu_2, \dots, a_{\mu-1} + tu_{\mu-1}, s, a_{\mu+1}, \dots, a_n)$$

They are really similar to the vectors p_μ , for all the previous components to the $\mu - th$ one, we are just placing their respective component from $a + tu$, in the $\mu - th$ component we are placing a different number $s \in \mathbb{R}$ and the rest are going to be their respective component in a .

Hence, we can state that:

$$f(p_\mu) - f(p_{\mu-1}) = g_\mu(a_\mu + tu_\mu) - g_\mu(a_\mu)$$

Now, we can use the mean value theorem to claim that $\exists \theta_\mu(t) \in (0, 1)$ such that:

$$g_\mu(a_\mu + tu_\mu) - g_\mu(a_\mu) = (a_\mu + tu_\mu - a_\mu) \frac{dg_\mu}{ds}(a_\mu + \theta(t)tu_\mu) = tu_\mu \frac{dg_\mu}{ds}(a_\mu + \theta(t)tu_\mu)$$

By rewriting g'_μ in terms of f we see that:

$$\frac{dg_\mu}{ds} = \frac{\partial f}{\partial x_\mu}(a_1 + tu_1, \dots, a_{\mu-1} + tu_{\mu-1}, s, a_{\mu+1}, \dots, a_n)$$

Additionally, to simplify it, we can write another set of functions $q_\mu : \mathbb{R} \rightarrow \mathbb{R}^n$ such that:

$$q_\mu(t) = \begin{pmatrix} a_1 + tu_1 \\ \vdots \\ a_{\mu-1} + tu_{\mu-1} \\ a_\mu + \theta_\mu(t)tu_\mu \\ a_{\mu+1} \\ \vdots \\ a_n \end{pmatrix} \implies g_\mu(a_\mu + tu_\mu) - g_\mu(a_\mu) = tu_\mu \frac{\partial f}{\partial x_\mu}(q_\mu(t))$$

Therefore:

$$\begin{aligned} f(p_\mu) - f(p_{\mu-1}) &= g_\mu(a_\mu + tu_\mu) - g_\mu(a_\mu) = tu_\mu \frac{\partial f}{\partial x_\mu}(q_\mu(t)) \\ \implies \frac{f(a + tu) - f(a)}{t} &= \sum_{\mu=1}^n \frac{tu_\mu \frac{\partial f}{\partial x_\mu}(q_\mu(t))}{t} = \sum_{\mu=1}^n u_\mu \frac{\partial f}{\partial x_\mu}(q_\mu(t)) \end{aligned}$$

Applying this result to the full directional derivative's definition:

$$D_u f = \lim_{t \rightarrow 0} \frac{f(a + tu) - f(a)}{t} = \lim_{t \rightarrow 0} \sum_{\mu=1}^n u_\mu \frac{\partial f}{\partial x_\mu}(q_\mu(t))$$

Here, the only term we have relying on t is q_μ , but if we peek its definition we will see that when $t \rightarrow 0$, $q_\mu(t) \rightarrow a$.

$$= \sum_{\mu=1}^n u_\mu \frac{\partial f}{\partial x_\mu}(a)$$

We got the summation over components of a vector and partial derivatives, but we can use here the nabla operator to write it using the inner product.

$$= \langle \nabla f(a), u \rangle = \nabla f(a) \cdot u$$

That is why we can use also the notation $\nabla_u f$ to refer to the directional derivative.

$$D_u f(a) = \nabla_u f(a) = \langle \nabla f(a), u \rangle$$

7 The ∇ operator

There is a vector which we are going to be using quite often in multi-variable calculus which receives the name of "nabla" and it is represented with an upside-down triangle ∇ or with an arrow on top as a regular basic vector $\vec{\nabla}$ although in order to simplify the notation here we will be using exclusively the notation without the arrow.

This vector is super special as it is defined with partial derivatives like:

$$\nabla := \begin{pmatrix} \partial_{x_1} \\ \partial_{x_2} \\ \vdots \\ \partial_{x_n} \end{pmatrix}$$

Or, equivalently:

$$\nabla_\mu = \frac{\partial}{\partial x_\mu}$$

This is just notation, but we can do some really nice tricks as multiplying this vector with a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$:

$$\nabla f = \begin{pmatrix} \partial_{x_1} f \\ \partial_{x_2} f \\ \vdots \\ \partial_{x_n} f \end{pmatrix}$$

8 Gradient

Let us take a scalar field $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and a constant $\lambda \in \mathbb{R}$. We parameterize the level curve of f under the function $\vec{r} : \mathbb{R} \rightarrow \mathbb{R}^n$. Thus, we have the equation:

$$f(\vec{r}(t)) = \lambda$$

By differentiating both sides:

$$\frac{d}{dt} f(\vec{r}(t)) = \frac{d}{dt} \lambda$$

We see that in the left-hand part we have to use the chain rule and on the right-hand part we just have 0:

$$\sum_{\mu=1}^n \frac{\partial f}{\partial x_{\mu}} \frac{dx_{\mu}}{dt} = 0$$

Here we have exactly the inner product:

$$\langle \nabla f, \vec{r}' \rangle = 0$$

Therefore, we just found out that the operation $\text{grad}(f) = \nabla f$ is a vector perpendicular to the parameterized path of an arbitrary level curve of f , which means that the gradient is pointing at any point to either a relative maximum or a relative minimum. By the nature of derivatives, we know that they are positive when the next value is greater than the current one, which means we can claim that they are actually pointing to a relative maximum.

9 Chain rule

Let us attempt to derive the following function:

$$\frac{d}{dt_{\mu}} f(x_1(t_1, \dots, t_{\mu}, \dots, t_n), x_2(t_1, \dots, t_{\mu}, \dots, t_n), \dots, x_m(t_1, \dots, t_{\mu}, \dots, t_n))$$

We will use vector notation defining $\vec{v} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ like:

$$\frac{d}{dt_{\mu}} f(\vec{v}(t_1, \dots, t_{\mu}, \dots, t_n))$$

Now, we will just use the standard derivative's definition.

$$\frac{d}{dt_{\mu}} f(\vec{v}(t_1, \dots, t_{\mu}, \dots, t_n)) = \lim_{h \rightarrow 0} \frac{f(\vec{v}(t_1, \dots, t_{\mu} + h, \dots, t_n)) - f(\vec{v}(t_1, \dots, t_{\mu}, \dots, t_n))}{h}$$

Before going on with this, we will work with the derivative of $\vec{v}$ with respect to the variable t_{μ} :

$$\frac{\partial \vec{v}}{\partial t_{\mu}} = \lim_{h \rightarrow 0} \frac{\vec{v}(t_1, \dots, t_{\mu} + h, \dots, t_n) - \vec{v}(t_1, \dots, t_{\mu}, \dots, t_n)}{h}$$

Next, we will add an arbitrary function $E(h)$ whose definition is unknown but we must establish that when $h \rightarrow 0$, $E(h) \rightarrow 0$ too.

$$\frac{\partial \vec{v}}{\partial t_{\mu}} \approx \frac{\vec{v}(t_1, \dots, t_{\mu} + h, \dots, t_n) - \vec{v}(t_1, \dots, t_{\mu}, \dots, t_n)}{h} + E(h)$$

We can solve this equation for the first term in the nominator by subtracting both sides by $E(h)$, multiplying by h and adding the second term of the nominator.

$$\vec{v}(t_1, \dots, t_{\mu} + h, \dots, t_n) \approx \vec{v}(t_1, \dots, t_{\mu}, \dots, t_n) + h \frac{\partial \vec{v}}{\partial t_{\mu}} - h E(h)$$

We can simplify a little the expression by using another function $o(h)$, which is a function such that:

$$h \rightarrow 0, \quad \frac{|o(h)|}{h} \rightarrow 0$$

$hE(h)$ satisfies that condition, thus we can replace it.

$$\vec{v}(t_1, \dots, t_\mu + h, \dots, t_n) \approx \vec{v}(t_1, \dots, t_\mu, \dots, t_n) + h \frac{\partial \vec{v}}{\partial t_\mu} - o(h)$$

We just found another way to write the variables of the function f in the first term of the derivative. Thus:

$$\frac{d}{dt_\mu} f(\vec{v}(t_1, \dots, t_\mu, \dots, t_n)) = \lim_{h \rightarrow 0} \frac{f(\vec{v}(t_1, \dots, t_\mu, \dots, t_n) + h \frac{\partial \vec{v}}{\partial t_\mu} - o(h)) - f(\vec{v}(t_1, \dots, t_\mu, \dots, t_n))}{h}$$

By reading the result, it is quite noticeable that the result is in fact a directional derivative.

$$\frac{d}{dt_\mu} f(\vec{v}(t_1, \dots, t_\mu, \dots, t_n)) = \langle \nabla f, \partial_{t_\mu} \vec{v} \rangle = \sum_{\nu=1}^n \frac{\partial f}{\partial x_\nu} \frac{\partial x_\nu}{\partial t_\mu}$$

Therefore, a more general way to compute a derivative for a function with m variable whose variables could also be functions of n variables which rely on more values and taking one of them as the variable for the differentiation is:

$$\frac{df}{dt_\mu} = \sum_{\nu=1}^n \frac{\partial f}{\partial x_\nu} \frac{\partial x_\nu}{\partial t_\mu}$$

But, in case the variables of the function only relies on a single variable, we can use the typical Leibniz's notation for single-variable functions's derivatives.

$$\frac{df}{dt} = \sum_{\nu=1}^n \frac{\partial f}{\partial x_\nu} \frac{dx_\nu}{dt}$$

And here, if we multiply both side by dt , we have a more general definition for the differential of a function:

$$df = \sum_{\nu=1}^n \frac{\partial f}{\partial x_\nu} dx_\nu$$

10 Multiple integrals

10.1 Fubini's theorem