

# New critical and collective phenomena in the random networks.

A. Gorsky (IITP RAS)

April 20 , Yandex

- In collaboration with V. Avetisov(Moscow), M. Hovanessian(Erevan), S.Maslov(Chicago) ,S. Nechaev(Moscow+Paris), M. Tamm(Moscow), O. Valba(Moscow)
- Phys.Rev E93,012302(2016), Phys. Rev E94,062313 (2016)  
Phys.Rev E96,062303(2017) + arXiv 1801.03912 + to appear

# Outline of the talk

- Motivation and key questions
- Numerical findings for constrained Erdos-Renyi network(CERN) and regular random graph (RRG)
- Spectral analysis. Eigenvalue tunneling and the ground state of constrained networks.
- Propagation of the excitation and spectral properties
- Application. Critical phenomena in generalized Schelling model of segregation
- Application to connectome

# Properties of networks

- Degree distribution
- Clusterization
- Typical path on the network
- Number of layers and interlayer interaction
- Synchronization
- Propagation of the signal on the network
- Phase transition in the exponential networks

# Network Models: Summary

## Erdős-Renyi model

- short path lengths
- Poisson distribution (no hubs)
- no clustering

## Watts-Strogatz Small World model


- short path lengths
- high clustering  
( $N$  independent)
- almost constant degrees

## Barabási-Albert scale-free model

- short path lengths
- power-law distribution for degrees
- robustness
- no clustering (may be fixed)



## Real-world networks

- short path lengths
- high clustering
- broad degree distributions,  
often power laws




Typical examples of network organization

# Multilayer networks




# Exponential networks — network ensembles with weights for some motifs


3-node motifs



4-node motifs



5-node motifs



# Mixed network ensembles

- Investigation of the mixed network ensemble. Set of local constraints(degree conservation) + additional chemical potential for some motif(substructure)
- The degree of the vertex fixed. Standard condition in chemistry and biology. May be fixed by some topological arguments
- Properties of the mixed ensembles differ from the properties of canonical or microcanonical ensembles



# Multicluster networks

- How multicluster networks emerge?
- How interaction between the clusters depends on the parameters of networks? When the clusters get synchronized?
- How the external probe and excitations behave on the clustered networks?





# Experimental data for one color

- The CERN network or RRG
- The degree of all nodes in the network is conserved-constraint
- Chemical potential for the triangles. **Mixed ensemble**
- Phase transitions. Cliques formation above critical chemical potential
- For CERN network  $[1/p]$  cliques emerge (average number of links at vertex equals  $pN$ )


# Model.Mixed ensemble

$$H = -\mu N_{\Delta}, \quad (\mu > 0) \quad (= \text{Tr } A^3, A \text{ -adjacency matrix})$$

- The possible moves in the network(= kinetic term)

undirected subgraphs- triads				
	[0]	[1]	[2]	[3]
concentration	$c_0$	$c_1$	$c_2$	$c_3$

(a)



(b)

Maslov,  
Sneppen  
2004

Figure 1: a) Possible triads in a non-directed network; b) Single link permutation: links (12) and (34) are removed, and links (13) and (24) are created. Triad {135} goes from type [0] to type [1], triads {125, 345} – from type [2] to type [1], and triad {245} – from type [2] to type [3]: three new triads of type [1] and one triad of type [3] are created instead of three triads of type [2] and one of type [0], compare to Eq.(1).




FIG. 1: The number of clusters  $N_{cl}$  as a function of the probability  $p$  in ER graph. The numerical data are obtained by averaging over 100 randomly generated graphs up to 512 vertices. Numerical values are fitted by the curve  $p^{-0.95}$ ; the behavior in doubly logarithmic scale is shown in the insert.

The network is completely defragmented into the finite number of weakly coupled dense droplets above the critical point. Both for CERN and RRG

Model without degree conservation — Strauss model (1986), solved  
In mean field approximation (Newman, Park -2004)





FIG. 5: Few typical samples of intermediate stages of the network evolution: upper panel – evolution with fixed vertex degree; lower panel – evolution with non-fixed vertex degree.

Intermediate «spin glass- like» pattern?

# Typical evolution of the random initial network to the ground state




Number of the droplets in the ground state can be predicted!  
All clusters are almost complete graphs.

# Spectral anatomy of transitions



- Second zone formation from the separated eigenvalues moving from the central zone.
- Number of isolated eigenvalues equals the number of clusters (Newman et al, 13)
- Semicircle distribution before the phase transition. «Triangle»-shape density in the central zone after the phase transition + second zone.
- Strong effect of intercluster interaction.

# Spectral density of adjacency matrix before and after phase transition





Second zone formation from clusters. Number of the Isolated eigenvalues of the adjacency matrix equals to the number of clusters





The second zone in the spectrum of adjacency matrix corresponds to the soft modes in spectrum of graph Laplacian

# Spectral density for unicolor case

- Spectral density for each cluster


Account of intercluster entanglement amounts to the very different total spectral density



# Application of the random network theory

- Transport networks
- Linguistic networks
- Evolutionary network
- Social networks and segregation models.
- Connectome- brain network
- Network of space-time and «Holographic Universe».

# Schelling model of segregation. Single parameter-level of tolerance




# Model for two colors-communities


- Chemical potentials for trimers are the same for all colors. Degrees of nodes conserved
- Two possibilities. Colored CERN network and colored RRG. The results for two cases are the same.
- Communities — countries, genders etc

Generalization of Schelling model with two parameters.  
One parameter- the weight for the intercommunity links  
Second parameter — the weight for the formation of triad inside community.


$$Z = \sum e^{-(\mu_G N_G + \mu_R N_R + \nu D N_{RG})}$$



Phase1.      Bipartite phase  
(intercommunity links win)



The line of phase transition



Phase 2.

Network = 2 unicolor clusters connected with  
A few links

(intra-connections win)







FIG. 3: (a)  $N_{RG}$  at  $\nu = 0.1$  and the behavior of the second eigenvalue of Laplacian matrix,  $\lambda_2$  as functions of  $\mu$ ; (b)  $N_{RG}$  as a function of  $\nu$  at  $\mu = 0.1$  and the behavior of the second eigenvalue of Laplacian matrix.




Near the transition between the in-dominated and Inter-dominated phases the interaction occurs via hubs




# What it means for social dynamics?

- If the weights for the preferences of intra- or inter- connections are high the transitions between phases only via sharp phase transition. The playground for manipulations
- Any evolutionary transition occurs in the crossover region of low weights.
- When the number of links between the communities is small the interaction occurs via the hubs- ambassadors.

# A bit of spectral analysis



CERN



RRG

The dependence of the first nonvanishing eigenvalue of Laplacian matrix (red) and dependence of the number of interlayer links (blue) on the chemical potential for trimers



Trimer

$$\lambda_2(\gamma) = cN_{bw}(\gamma)$$

c- some constant

The problem of evolution of  $\lambda_2$ .

The key phenomena — intersection and rearrangements of the spectrum of first modes  
In the block matrices

$$\det \begin{pmatrix} (A - \lambda) & C \\ C^T & (D - \lambda) \end{pmatrix} = \det(A - \lambda) \det(D - C^T(A - \lambda)^{-1}C) = 0$$

Chemical potential  
for trimers

Condition for the plateau  
entrance


$$\lambda_2^A(\gamma) = \lambda_3(L)$$

Eigenvalue of the  
one layer Laplacian

Eigenvalue of the  
Laplacian of the whole  
network


Two layers are absolutely correlated at the plateau!

The **semiinfinite** plateau in more simple two-layer network only with interlayer interaction- phenomena of superdiffusion (Arenas et al 2013)



$$\mathcal{L} = \left( \begin{array}{c|c} D_1 L_1 + D_x I & -D_x I \\ \hline -D_x I & D_2 L_2 + D_x I \end{array} \right)$$

Laplacian of the whole network



Analogue of our plateau in the model with One-one interaction between layers

The layers are absolutely synchronized at plateau. Phenomena of Superdiffusion due to synchronization

# Matrix model description


Adjacency matrix is symmetric random matrix involving 1 and 0 only

$$Z = \int dM \exp(a \text{Tr} M^2 - \mu \text{Tr} M^3)$$

Additional constraint: sum of elements in each row and each column is fixed


a- parameter of the network . Chemical potential for the number of triangles yields «interaction cubic term».

The matrix model counterpart of the cluster formation.  
Eigenvalue tunneling — instanton, nonperturbative  
phenomena in many physical situations. Formation  
of stable D-branes in the string theory from unstable  
branes.  
Baby-universes in 2d gravity




# Spontaneous breaking of the symmetry in the two-color network

( no degree conservation)




The parameters of the black and white subsystems are the same!





# Phase transition in two-color network above the critical weight for triads (no degree conservation)

liquid



White cluster

Black cluster

# Transport on the network

- How excitations behave on the network?
- Diffusion on the network is defined by the largest eigenvalue of the adjacency matrix
- Is there localization of the excitation on the network?

## Standard criteria for localization


- Level spacing distribution (between the neighbor levels)

$$\begin{cases} P_{deloc}(s) = s e^{-as^2} \\ P_{loc}(s) = e^{-s} \end{cases}$$

- Participation ratio or inverse participation ratio
- Area or volume law for the entanglement


Let us treat our RRG and CERN as the «Fock space» for some interacting many-body system. Could we say smth about this many-body systems using our new critical phenomena?

## First check- level spacing distribution



Level spacing distribution in the «nonperturbative» zone.  
Poisson distribution — insulator zone.

## The spectral density of adjacency matrix above the phase transition



The delocalization in «perturbative» and localiation in  
«nonperturbative» bands

## Non-ergodicity of delocalized modes


- We have considered the artificial network of clusters with the same parameters as our initial random network. Its spectral Density is different from the network prepared from the initial random network. Hence in the delocalized regime there is dependence on the initial condition! Mark of non-ergodicity.

# Connectome

- Only the connectome of C.Elegance is completely known -312 neurons(nodes)
- Connectome Program(USA-3 млрд \$, 2015-2025 гг)
- The number of neurons increases during first 2 years, is approximately stable till 60 and then starts to decrease
- There are structural and functional networks in connectome

Eduardo Izquierdo

<http://www.basis.ncl.ac.uk>




<http://neuroinformatics2012.org/>


<http://www.genome.gov/10000570>






# MRI Acquisition





# Connectom. What is known?

- Small-world network. Hierarchical network
- Average clusterization
- Many hubs
- Anomalously large number of «open triads»
- Relation between the diseases and structural changes in connectome
- Complicated synchronization of functional networks





## Коннектом человека сильнее отличается от своей случайной копии




Рис.3 Спектр коннектома человека и его рандомизированного аналога

# New findings

- The human connectom is much more non-ergodic compared with other animals
- The local clusterization seems to be very important in connectom evolution(at least for human)
- The localization properties of the excitation in human connectome(level spacing distribution) are unusual(seems that mobility edge exists)
- K.Anokhin, V.Avetisov, S.Nechaev,N.Pospelov  
O.Valba, A.G. To appear

# И раньше догадывались, что случайные сети приносят неожиданные результаты

«Тятя,тятя,наши сети притащили мертвеца» (А.С. Пушкин)

И в советское время.

