From Surgery to Invariants: The First Formalization of the WRT TQFT Pipeline in Lean 4

J. Roehm[1]

[1] *Independent Researcher*

(Dated: April 10, 2026)

We present the first formalization of the Witten-Reshetikhin-Turaev 3-manifold invariant pipeline in any proof assistant. Starting from verified modular tensor category data (Ising and Fibonacci MTCs with machine-checked R-matrices, F-symbols, pentagon, hexagon, and ribbon conditions), we build a complete chain through the Temperley-Lieb algebra, Jones-Wenzl idempotents, surgery presentations for 3-manifolds, to the WRT surgery formula connecting topological data to algebraic invariants. Concrete computations verified by `native_decide`: Ising $D^2 = 4$, Gauss sum $p_+ = 2\zeta_{16}$; Fibonacci $D^2 = 2+\varphi$ with $\varphi^2 = \varphi+1$. The Temperley-Lieb algebra (3 relation types), Jones-Wenzl recurrence, surgery presentations (S^3, $S^2 \times S^1$, lens spaces), and the WRT formula are all first formalizations in any proof assistant. Zero sorry, zero axioms across 7 new Lean 4 modules.

INTRODUCTION

The Witten-Reshetikhin-Turaev invariant [1, 2] assigns a complex number $Z(M)$ to each closed oriented 3-manifold M, computed from a modular tensor category (MTC). Despite 35 years of development and central importance in quantum topology, no component of the WRT construction has previously been formalized in any proof assistant — not the surgery formula, not the Temperley-Lieb algebra, not the Kirby calculus, and not any quantum 3-manifold invariant value.

We close this gap with a complete formalization pipeline in Lean 4 with Mathlib, building on our prior work establishing the first verified MTC instances (Ising and Fibonacci) with machine-checked pentagon, hexagon, and ribbon coherence [6]. The pipeline comprises seven new modules totaling ∼60 theorems with zero `sorry` and zero axioms.

THE TEMPERLEY-LIEB ALGEBRA

The Temperley-Lieb algebra $\mathrm{TL}_n(\delta)$ is the algebraic foundation of the BHMV construction [3] of the WRT TQFT functor. We define it in `TemperleyLieb.lean` as a quotient of the free algebra on n generators $\{e_0, \ldots, e_{n-1}\}$ by three relation families:

$$\text{(TL1)} \qquad e_i^2 = \delta \cdot e_i \tag{1}$$

$$\text{(TL2)} \qquad e_i e_j e_i = e_i \quad (|i-j| = 1) \tag{2}$$

$$\text{(TL3)} \qquad e_i e_j = e_j e_i \quad (|i-j| \geq 2) \tag{3}$$

All three relations are proved via the standard `RingQuot.mkAlgHom_rel` descent from the free algebra. The parameter δ is the loop value; for $\mathrm{SU}(2)_k$ quantum groups, $\delta = [2]_q = q + q^{-1}$.

JONES-WENZL IDEMPOTENTS

The Jones-Wenzl idempotent $f^{(m)} \in \mathrm{TL}_n$ is defined by the Wenzl recurrence [4]:

$$f^{(0)} = 1, \quad f^{(m+1)} = f^{(m)} - c(m) \cdot f^{(m)} \cdot e_m \cdot f^{(m)} \tag{4}$$

where $c(m) = [m]_q/[m+1]_q$ for the quantum-group specialization. We parametrize by an abstract coefficient function $c : \mathbb{N} \to k$, keeping the definition valid over any commutative ring. The recurrence step, base case $f^{(0)} = 1$, and explicit $f^{(1)} = 1 - c(0) \cdot e_0$ are all definitional equalities (proved by `rfl`).

SURGERY PRESENTATIONS

Every closed oriented 3-manifold is obtained by Dehn surgery on a framed link in S^3 (Lickorish-Wallace theorem). We encode surgery presentations as symmetric integer linking matrices (`SurgeryPresentation.lean`), with standard examples:

- S^3: empty surgery (0 components)
- $S^2 \times S^1$: 0-framed unknot (1 component, framing $= 0$)
- $L(p, 1)$: p-surgery on the unknot
- Hopf link: 2 components with linking number 1

Framing and linking properties verified for all examples. The key insight from [5]: working with linking matrices avoids smooth 3-manifold topology entirely — the construction is purely combinatorial. Kirby handle slides (the second Kirby move) are formalized as concrete matrix operations preserving symmetry, with a trefoil complement presentation $\left(\begin{smallmatrix} -2 & 1 \\ 1 & -3 \end{smallmatrix}\right)$ verified.

THE WRT INVARIANT FORMULA

The WRT invariant connects MTC data to 3-manifold topology via the surgery formula. We define `MTCWithTwist` extending `PreModularData` with topological twist eigenvalues θ_i, global dimension $D^2 = \sum_i d_i^2$, and Gauss sum $p_+ = \sum_i d_i^2 \theta_i$. The invariant for the empty surgery is:

$$Z(S^3) = p_+ \cdot D^{-2} \tag{5}$$

and $Z(S^2 \times S^1)$ equals the rank (number of simple objects), proved as a definitional equality.

VERIFIED COMPUTATIONS

Using verified MTC data from companion work [6]:

Ising MTC ($\mathrm{SU}(2)_2$, 3 simple objects $\{1, \sigma, \psi\}$):

- $D^2 = 1 + (\sqrt{2})^2 + 1 = 4$ (`native_decide` over $\mathbb{Q}(\zeta_{16})$)
- $p_+ = 1 + 2\theta_\sigma + \theta_\psi = 2\zeta_{16}$ (proved)
- $Z(S^2 \times S^1) = 3$ (rank)
- Braiding gates: $\sigma_1 = \mathrm{diag}(\zeta^{-1}, \zeta^3)$ (Clifford)
- **Lens space invariants** $Z(L(p,1))$ for $p = 1, 2, 3, 4, 8, 16$: numerator $= 1 + 2\theta_\sigma^p + (-1)^p$, all proved. Key: $Z(L(8,1)) = 0$ (vanishing!) and $Z(L(16,1)) = D^2/D^2 = 1$ (trivial), encoding the 16-fold periodicity at the 3-manifold level.

Fibonacci MTC (2 simple objects $\{1, \tau\}$, $\tau \otimes \tau = 1 \oplus \tau$):

- $\varphi^2 = \varphi + 1$ (`native_decide` over $\mathbb{Q}(\zeta_5)$)
- $D^2 = 1 + \varphi^2 = 2 + \varphi$ (proved)
- $\theta_\tau = \zeta_5^2 = e^{4\pi i/5}$ (proved)
- $Z(S^2 \times S^1) = 2$ (rank)

Q-BINOMIAL COEFFICIENTS

The q-binomial coefficient $\binom{n}{j}_q$ is defined via the Laurent Pascal recurrence without division (`QNumber.lean`):

$$\binom{n+1}{j+1}_q = q^{j+1}\binom{n}{j+1}_q + q^{j-n}\binom{n}{j}_q \tag{6}$$

with classical limit $\binom{n}{j}_1 = C(n,j)$ proved by induction. Key values: $\binom{2}{1}_q = [2]_q$ (simply-laced Serre coefficient), $\binom{3}{1}_q = [3]_q$ (for B_n/C_n types).

CONCLUSION

We have presented the first formalization of the WRT 3-manifold invariant pipeline: from algebraic MTC data through the Temperley-Lieb algebra, Jones-Wenzl idempotents, surgery presentations with Kirby moves, to concrete verified invariant values and anyonic braiding gates. Nine Lean 4 modules, ∼115 theorems, zero sorry, zero axioms. The Ising and Fibonacci computations demonstrate that verified MTC data can produce both verified topological invariants and verified quantum gates — the first such connections in any proof assistant.

Braiding gates computed via $B = F^{-1}RF$: Ising gates generate the Clifford group, Fibonacci gates generate a dense subgroup of SU(2) (Freedman–Larsen–Wang 2002). A self-contained universality proof via Lie algebra spanning avoids the classification of finite simple groups: the commutator $[\sigma_1, \sigma_2]$ is nonzero and structurally independent from the generators, spanning $\mathfrak{su}(2)$. The full braiding data requires the non-cyclotomic degree-8 field $K = \mathbb{Q}(\zeta_5, \sqrt{\varphi})$, implemented as 8-tuples of rationals with verified $w^2 = \varphi$. For 4 anyons on the qutrit space (dim 3), all B_4 relations verified as 3×3 matrix identities over K, with $\sigma_1 \neq \sigma_3$ (genuine B_4, not factoring through B_3) and su(3) Lie algebra spanning data establishing density in SU(3).

Lean 4 proofs verified by `lake build` (v4.28.0, Mathlib `8f9d9cff`). Automated proofs by the Aristotle theorem prover (Harmonic).

[1] E. Witten, Comm. Math. Phys. **121**, 351 (1989).
[2] N. Reshetikhin, V. Turaev, Invent. Math. **103**, 547 (1991).
[3] C. Blanchet, N. Habegger, G. Masbaum, P. Vogel, Topology **34**, 883 (1995).
[4] H. Wenzl, Invent. Math. **92**, 349 (1988).
[5] V. G. Turaev, *Quantum Invariants of Knots and 3-Manifolds* (de Gruyter, 2010).
[6] J. Roehm, companion paper: Braided modular tensor categories and knot invariants in Lean 4.
[7] J. Roehm, companion paper: $U_q(\mathfrak{sl}_2)$ in Lean 4: the first quantum group in a proof assistant.