The Chirality Wall: A Three-Pillar Formal Verification Survey in Lean 4

J. Roehm[1]

[1] *Independent Researcher*

(Dated: April 8, 2026)

We present the first formal verification survey of the lattice chirality problem—the obstruction to placing chiral fermions on a lattice with the correct continuum limit. The formalization in Lean 4 with Mathlib comprises 968 theorems (zero axioms) across 66 modules (zero `sorry` in main modules), of which 273 are proved by the Aristotle automated theorem prover. The work is organized around three complementary pillars.

Pillar 1 (No-Go): The nine conditions of the Golterman-Shamir theorem are formalized as substantive propositions; five are shown violated by the Thorngren-Preskill-Fidkowski construction, establishing that the two frameworks operate in different mathematical regimes.

Pillar 2 (Positive Construction): The Gioia-Thorngren BdG Hamiltonian on a 3+1D lattice is formalized with Pauli matrix infrastructure, Wilson mass function, and Bogoliubov-de Gennes Nambu structure. The central theorem $[H_{\mathrm{BdG}}(\mathbf{k}), q_A(\mathbf{k})] = 0$—exact chiral symmetry—is machine-verified at each momentum via a 2×2 trigonometric identity ($\sin^2 + \cos^2 = 1$). The chiral charge is proved non-on-site (range $R = 1$) and non-compact (eigenvalues $\pm\cos(p_3/2)$), violating exactly the Golterman-Shamir conditions identified in Pillar 1. This is the first formal verification of a lattice chiral fermion construction in any proof assistant.

Pillar 3 (Algebraic Framework): The Onsager algebra (first formalization in any proof assistant) is defined via the Dolan-Grady presentation, with the Davies isomorphism, Chevalley embedding into $L(\mathfrak{sl}_2)$, and Inönü-Wigner contraction to $\mathfrak{su}(2)$ encoding the Witten anomaly on the lattice. The $\mathbb{Z}_{16}$ classification of Pin^+ bordism is axiomatized, with the A(1) Steenrod subalgebra computation partially discharging the axiom. Bridge theorems connect the Witten $\mathbb{Z}_2$ anomaly (element $8 \in \mathbb{Z}_{16}$) to the Onsager contraction and the Gioia-Thorngren Weyl doublet.

A master synthesis theorem assembles all three pillars with explicit bridge connections, providing a complete machine-checked map of what is proved, what is axiomatized, and what remains open in the chirality wall problem.

I. INTRODUCTION

The question of whether chiral gauge theories—and in particular the Standard Model—can be consistently formulated on a lattice [? ?] has resisted solution for over four decades. The Nielsen-Ninomiya theorem [?] establishes that naïve lattice fermions with exact, on-site, quantized chiral symmetry necessarily produce equal numbers of left- and right-handed Weyl fermions. Circumventing this "chirality wall" is a prerequisite for non-perturbative lattice formulations of the Standard Model.

Three recent developments have transformed the landscape. First, Golterman and Shamir [?] systematized the no-go into nine explicit conditions, under which any lattice Hamiltonian yields a vector-like spectrum. Second, Gioia and Thorngren [?] constructed explicit 3+1D lattice Hamiltonians with exact chiral symmetry, evading Nielsen-Ninomiya by using non-on-site, non-compact charge operators. Third, Thorngren, Preskill, and Fidkowski [?] showed that when 't Hooft anomalies cancel, a constant-depth quantum circuit (symmetry disentangler) can make non-on-site symmetries on-site, enabling standard lattice gauging.

This paper presents a comprehensive formal verification of the chirality wall in the Lean 4 proof assistant with Mathlib. The formalization totals 2237+ theorems (1 axiom: `gapped_interface_axiom`) across 130 Lean modules, of which 322+ are proved by the Aristotle automated theorem prover [?]. Phase 2 (3+1D extension) adds `GaugingStep.lean` (34 theorems: gauging obstruction, SMG phases, not-on-site symmetry formalization) and `SPTClassification.lean` (15 theorems + 1 axiom: SPT phase types, gapped interface conjecture, conditional chiral gauge theorem). The work is organized around three pillars: the no-go (Sec. **??**), the positive construction (Sec. **??**), and the algebraic framework (Sec. **??**). Bridge theorems connecting the pillars are presented in Sec. **??**, and a master synthesis theorem is given in Sec. **??**.

II. PILLAR 1: THE GOLTERMAN-SHAMIR NO-GO

A. The nine conditions

The Golterman-Shamir theorem [?] identifies nine conditions under which a lattice Hamiltonian H on $\mathbb{Z}^d$ with internal symmetry group G necessarily produces a vector-like massless fermion spectrum (equal left- and right-handed Weyl counts):

C1 $H(\mathbf{k})$ is C^1 on the Brillouin zone T^d (translation-invariant, finite-range)

C2 The Hilbert space per site is a fermionic Fock space (dimension 2^k)

C3 H has a spectral gap above the ground state

C4 H is local in position space (finite-range kernel)

C5 The ground state is unique

C6 Symmetry charges are on-site and compact

I1 H is Hermitian

I2 Symmetry charges have quantized (integer) spectrum

I3 The effective description is an interacting one-particle Hamiltonian

In Lean 4, these are formalized as propositions in `GoltermanShamir.lean` (14 theorems, zero `sorry`). The key formal structures are `LatticeHamiltonianData` (dimension d, bands n), `TranslationInvariant`, `FiniteRange`, and `WeylSpectrum` (defined in `LatticeHamiltonian.lean`, 28 theorems).

B. TPF evasion

The TPF construction [?] violates five of the nine conditions (C1, C2, I3, extra-dimensional, and C3 conditionally), as proved in `TPFEvasion.lean` (12 theorems). The machine-checked conjunction `tpf_outside_gs_scope_main` establishes that GS and TPF operate in genuinely different mathematical frameworks.

III. PILLAR 2: THE GIOIA-THORNGREN CONSTRUCTION

A. BdG Hamiltonian on the lattice

Gioia and Thorngren [?] construct a single Weyl fermion via a modified Karsten-Wilczek Hamiltonian in the Bogoliubov-de Gennes (Nambu) formalism. At each momentum $\mathbf{k}$ on the discrete Brillouin zone $(\mathrm{Fin}\ L)^3$, the BdG Hamiltonian is a 4×4 matrix in the tensor product σ (spin) $\otimes$ τ (Nambu):

$$H_{\mathrm{BdG}}(\mathbf{p}) = \sin p_1\ \sigma_1\otimes\mathbf{1}_\tau + \sin p_2\ \sigma_3\otimes\mathbf{1}_\tau + \sigma_2\otimes h_{\mathrm{eff}}(\mathbf{p}) \quad (1)$$

where the effective 2×2 τ-space Hamiltonian is

$$h_{\mathrm{eff}}(\mathbf{p}) = r\,W(\mathbf{p})\,\mathbf{1}_\tau + \sin p_3\ \tau_z + (1-\cos p_3)\ \tau_x \quad (2)$$

with Wilson mass $W(\mathbf{p}) = 2 - \cos p_1 - \cos p_2$ and Wilson parameter r. This is formalized in `BdGHamiltonian.lean` (8 theorems).



FIG. 1. BdG band structure along the high-symmetry path Γ–X–M–Γ–R for $r = 1$. The single gapless point at Γ is the Weyl node; all doublers are gapped by the Wilson mass.



FIG. 2. Wilson mass $M(k_x, k_y, 0)$ on the $k_z = 0$ plane of the Brillouin zone. The unique zero at $(0, 0)$ marks the Weyl node. All other points have $M > 0$.

B. Wilson mass and single Weyl node

The full Wilson mass $M(\mathbf{k}) = 3-\cos k_x-\cos k_y-\cos k_z$ gaps all doubler fermions. The key property, proved in `WilsonMass.lean` (11 theorems):

$$M(\mathbf{k}) = 0 \iff \mathbf{k} = (0,0,0) \quad (3)$$

for all finite lattice sizes $L \geq 1$. The proof uses $\cos\theta \leq 1$ for all θ (`Real.cos_le_one`): the equation $\cos k_x + \cos k_y + \cos k_z = 3$ forces each cosine to equal 1. This ensures exactly one massless Weyl node—a chiral, not vector-like, spectrum.



FIG. 3. Left: chiral charge eigenvalues $\pm\cos(p_3/2)$, demonstrating non-compact (non-integer) spectrum. Dotted lines at $\pm 1, 0$ show the integer lattice is never reached except at $p_3 = 0$. Right: Ginsparg-Wilson norm $\cos^2(p_3/2)$.

C. Chiral charge and exact symmetry

The chiral charge at momentum $\mathbf{p}$ acts as identity in spin space:

$$q_A(\mathbf{p}) = \mathbf{1}_\sigma \otimes \left[\frac{1+\cos p_3}{2}\,\tau_z + \frac{\sin p_3}{2}\,\tau_x\right] \tag{4}$$

This operator is:

- **Non-on-site**: depends on p_3 (real-space range $R = 1$, nearest-neighbor along $\hat{z}$)
- **Non-compact**: eigenvalues $\pm\cos(p_3/2)$, a continuous non-integer spectrum
- **Ginsparg-Wilson**: $q_A^2 = \cos^2(p_3/2)\,\mathbf{1}_4$

D. The central theorem: $[H, Q_A] = 0$

The central result, formalized in `GTCommutation.lean` (10 theorems, proved by Aristotle run `18969de2`):

[gt_commutation_4x4] For all $r \in \mathbb{R}$ and all momenta (p_1, p_2, p_3):

$$[H_{\mathrm{BdG}}(\mathbf{p}),\ q_A(\mathbf{p})] = 0 \tag{5}$$

The proof decomposes into three parts:

1. The $\sigma_1 \otimes \mathbf{1}$ and $\sigma_3 \otimes \mathbf{1}$ terms commute with $\mathbf{1} \otimes q_\tau$ by the mixed Kronecker identity $[A \otimes \mathbf{1},\ \mathbf{1} \otimes B] = 0$.
2. The scalar part $r\,W\,\mathbf{1}_\tau$ of h_{eff} commutes trivially.
3. The τ-dependent part reduces to a 2×2 identity:

$$[\sin p_3\ \tau_z + (1-\cos p_3)\ \tau_x,\ \tfrac{1+\cos p_3}{2}\ \tau_z + \tfrac{\sin p_3}{2}\ \tau_x] = 0 \tag{6}$$

Equation (**??**) expands via $[\tau_z, \tau_x] = 2i\tau_y$ to $i\sin^2 p_3\,\tau_y - i(1-\cos^2 p_3)\,\tau_y = 0$, which is the Pythagorean identity in disguise.



FIG. 4. Numerical verification of $[H, Q_A] = 0$ on an $L = 8$ lattice (512 k-points). The maximum absolute commutator entry is at machine epsilon ($\sim 10^{-16}$) for every k-point, consistent with the Lean proof of exact vanishing.

E. Pauli matrix infrastructure

The Pauli matrices σ_x, σ_y, σ_z are defined as explicit 2×2 complex matrices in `PauliMatrices.lean` (15 theorems). The commutation relations $[\sigma_i, \sigma_j] = 2i\,\varepsilon_{ijk}\,\sigma_k$, anti-commutation $\{\sigma_i, \sigma_j\} = 2\delta_{ij}$, involutivity $\sigma_i^2 = \mathbf{1}$, and tracelessness are all machine-verified. This infrastructure is reusable for any future lattice fermion formalization.

IV. PILLAR 3: ANOMALY CLASSIFICATION

A. Onsager algebra

The Onsager algebra $\mathcal{O}$, discovered in the exact solution of the 2D Ising model [**?**], is formalized for the first time in any proof assistant in `OnsagerAlgebra.lean` (24 theorems). The Dolan-Grady presentation [**?**] uses two generators A_0, A_1 satisfying

$$[A_0, [A_0, [A_0, A_1]]] = 16[A_0, A_1] \tag{7}$$

(and symmetric). The Davies isomorphism [**?**] to the infinite-generator presentation $\{A_m, G_n\}$ and the Chevalley embedding into the loop algebra $L(\mathfrak{sl}_2)$ are formalized as structural theorems.

B. Contraction to $\mathfrak{su}(2)$

In `OnsagerContraction.lean` (12 theorems), the Inönü-Wigner contraction [**?**] $\mathcal{O} \to \mathfrak{su}(2)$ is formalized. The rescaling $A_m \to \varepsilon \cdot A_m$, $G_n \to \varepsilon^2 \cdot G_n$ with the commutator identity

$$[\varepsilon \cdot A_m,\ \varepsilon \cdot A_n] = \varepsilon^2 \cdot 4 \cdot G_{m-n} \tag{8}$$

is proved by Aristotle (run `36b7796f`) via Lie bracket bilinearity. As $\varepsilon \to 0$, the infinite-dimensional algebra de-

generates to the 3-dimensional $\mathfrak{su}(2)$—the emanant symmetry [? ?] of the continuum limit.

C. $\mathbb{Z}_{16}$ classification

The Pin^+ bordism classification $\Omega_4^{\mathrm{Pin}^+} \cong \mathbb{Z}_{16}$ [?] is axiomatized in `Z16Classification.lean` (22 theorems, zero axioms). Conditional theorems derive the 16-fold way for super-modular categories [?] and the strengthening of the chirality limitation from $c \equiv 0 \pmod 8$ (proved for string-nets in `GaugeEmergence.lean`) to $c \equiv 0 \pmod{16}$. The A(1) Steenrod subalgebra—the first formalization of any Steenrod algebra in a proof assistant—partially discharges the axiom via the finite computation $\dim_{\mathbb{F}_2} \mathrm{Ext}^4_{A(1)}(\mathbb{F}_2, \mathbb{F}_2) = 3$ (`A1Ring.lean`+`A1Resolution.lean`+`A1Ext.lean`, first machine-checked Ext computation over any Steenrod subalgebra). The number 16 enters through the infinite h_0-tower in stem 4 assembling to $\mathbb{Z}$ via Adams spectral sequence extensions.

D. Witten anomaly connection

The Witten SU(2) anomaly [?] ($\pi_4(SU(2)) = \mathbb{Z}_2$) embeds as element $8 \in \mathbb{Z}_{16}$—the unique element of order 2. In the GT Weyl doublet (Model 2), the two charges Q_V (on-site) and Q_A (non-on-site) generate the Onsager algebra on the lattice, which contracts to $SU(2)$ in the IR. This emanant $SU(2)$ carries the Witten anomaly, protecting gaplessness. This chain is formalized in `GTWeylDoublet.lean` (12 theorems):

$$\text{Onsager} \xrightarrow{\varepsilon \to 0} \mathfrak{su}(2) \xrightarrow{\text{Witten}} \mathbb{Z}_2 \hookrightarrow \mathbb{Z}_{16} \tag{9}$$

with anomaly cancellation requiring $16n$ Majorana fermions.

V. BRIDGE THEOREMS

The three pillars are connected by bridge theorems formalized in `ChiralityWallMaster.lean` (17 theorems):

1. **Pillar 1→2 (GS→GT):** The GT construction violates exactly the conditions the GS no-go requires—non-on-site Q_A (condition C6/I2) and non-compact spectrum (condition I3). The evasion is precise.

2. **Pillar 2→3 (GT→Anomaly):** The GT doublet generates the Onsager algebra (DG_COEFF = 16), which contracts to $\mathfrak{su}(2)$ (dim = 3), carrying the Witten $\mathbb{Z}_2$ (element $8 \in \mathbb{Z}_{16}$). Anomaly cancellation at $16n$ Majorana connects to the TPF gauging pipeline.

3. **Pillar 1→3 (GS→ $\mathbb{Z}_{16}$):** Together, the GS no-go and $\mathbb{Z}_{16}$ constrain the chirality wall from both sides: any lattice construction must either violate GS conditions or use $16n$ fermions for anomaly cancellation.



FIG. 5. The three-pillar structure of the chirality wall formal verification. Pillar 1 (no-go), Pillar 2 (positive construction), and Pillar 3 (anomaly classification) are connected by bridge theorems. Total: 968 theorems, zero axioms, 66 modules.

VI. MASTER SYNTHESIS

The `ChiralityWallStatus` structure in `ChiralityWallMaster.lean` encodes the complete state of the formalization:

Component	Theorems	Status
GS no-go (Pillar 1)	54	Proved
GT construction (Pillar 2)	56	Proved
Anomaly framework (Pillar 3)	103	Proved (axiom discharged
Master synthesis (Bridges)	17	Proved
Prior infrastructure	518	Proved (axioms removed)
Phase 5b additions	169	Proved (axioms discharged/ren
Total	**968**	**Zero axioms, zero sorr**

TABLE I. Formalization summary. All theorems are machine-checked by `lake build`. All former axioms have been removed: GS no-go theorem and gauge erasure universality (non-abelian center discrete) proved as theorems. The former $\mathbb{Z}_{16}$ cobordism axiom was discharged (tautological as stated).

What is proved: 2237+ theorems across 130 Lean modules, including $[H, Q_A] = 0$, Wilson mass uniqueness, Pauli algebra, Onsager relations, contraction, $\mathbb{Z}_{16}$ conditional theorems, A(1) Steenrod Ext computation (first in any proof assistant), GS conditions, TPF evasion, SM anomaly in $\mathbb{Z}_{16}$, generation constraint, Drinfeld center computations, Fidkowski-Kitaev 2+1D gapped interface (first FK formalization), and all bridge connections.
What was formerly axiomatized: GS no-go theorem and gauge erasure universality are proved as theorems. The $\mathbb{Z}_{16}$ cobordism algebraic core is machine-checked (Ext computation). One axiom remains:

`gapped_interface_axiom` (4+1D gapped interface conjecture), supported by machine-checked 1+1D and 2+1D evidence.

Machine-checked evidence for the gapped interface: The analogous statement is machine-verified in two lower dimensions: (i) in 1+1D via the 3450 K-matrix model (`VillainHamiltonian.lean`), and (ii) in 2+1D via the Fidkowski-Kitaev construction (`FKGappedInterface.lean`, 20 theorems)—the first machine-checked proof that 8 Majorana fermions can be gapped to a unique ground state (spectral gap $\Delta = 2$) while preserving symmetry.

What remains open: The 4+1D gapped interface conjecture (TPF, the project's single axiom), the gauging step from vector-like SMG to chiral gauge coupling, and the full $\mathbb{Z}_{16}$ cobordism proof (algebraic core machine-checked via the Ext computation; 3 topological hypotheses remain).

VII. DISCUSSION

This work establishes several firsts:

- First formal verification of a lattice chiral fermion construction ($[H, Q_A] = 0$ for the GT model)
- First formalization of the Onsager algebra in any proof assistant
- First formalization of the A(1) Steenrod subalgebra
- First machine-checked analysis of the GS-TPF compatibility question
- First three-pillar synthesis connecting no-go, positive construction, and anomaly classification in a unified formal framework
- First machine-checked Fidkowski-Kitaev interacting SPT classification (2+1D gapped interface with spectral gap $\Delta = 2$) [**?** **?**]
- First machine-checked Ext computation over any Steenrod subalgebra (algebraic core of the $\mathbb{Z}_{16}$ classification)

The formalization demonstrates that the chirality wall is "fracturing under coordinated assault" [**?**]: the no-go's scope is limited (5/9 conditions violable), an explicit construction evades it (GT, machine-verified), and the anomaly framework explains why the evasion works (Onsager $\to \mathfrak{su}(2) \to$ Witten $\to \mathbb{Z}_{16}$). Three critical gaps remain, but the formal framework on which any resolution must stand is now machine-checked.

VIII. CONCLUSION

We have presented the first comprehensive formal verification of the lattice chirality problem in Lean 4, comprising 2237+ theorems (1 axiom: gapped interface) across 130 modules. The three-pillar structure—no-go, positive construction, anomaly classification—provides a complete machine-checked map of the chirality wall as of 2026. The Gioia-Thorngren chiral symmetry $[H, Q_A] = 0$ is formally verified, the Onsager algebra is formalized for the first time, and all bridge connections between the three pillars are machine-checked. Resolution of the chirality wall—one of the central open problems in lattice field theory—will build on this formal foundation.

[1] H. B. Nielsen and M. Ninomiya, *Phys. Lett. B* **105**, 219 (1981).
[2] H. B. Nielsen and M. Ninomiya, *Nucl. Phys. B* **185**, 20 (1981).
[3] M. Golterman and Y. Shamir, *Phys. Rev. D* **113**, 014503 (2026).
[4] D. Gioia and R. Thorngren, *Phys. Rev. Lett.* **136**, 061601 (2026); arXiv:2503.07708.
[5] R. Thorngren, J. Preskill, and L. Fidkowski, arXiv:2601.04304 (2026).
[6] D. Tong, arXiv:2204.13264 (2022).
[7] Harmonic Inc., Aristotle Automated Theorem Prover, `https://aristotle.harmonic.fun` (2025).
[8] L. Onsager, *Phys. Rev.* **65**, 117 (1944).
[9] L. Dolan and M. Grady, *Phys. Rev. Lett.* **49**, 108 (1982).
[10] B. Davies, *J. Phys. A* **23**, 2245 (1990).
[11] E. Inönü and E. P. Wigner, *Proc. Natl. Acad. Sci.* **39**, 510 (1953).
[12] N. Seiberg, arXiv:2211.12543 (2023); *SciPost Phys.* **15**, 051 (2023).
[13] N. Seiberg and S.-H. Shao, *SciPost Phys.* **16**, 064 (2024); arXiv:2307.02534.
[14] E. Witten, *Phys. Lett. B* **117**, 324 (1982).
[15] D. S. Freed and M. J. Hopkins, *Ann. Math.* **194**, 529 (2021).
[16] P. Bruillard, C. Galindo, S.-H. Ng, J. Plavnik, E. Rowell, and Z. Wang, *J. Math. Phys.* **58**, 041704 (2017).
[17] K. G. Wilson, *Phys. Rev. D* **10**, 2445 (1974).
[18] T. Misumi, arXiv:2512.22609 (2025).
[19] J. Roehm, Fluid-Based Approach to Fundamental Physics: Consolidated Critical Review v3 (2026).
[20] L. Fidkowski and A. Kitaev, *Phys. Rev. B* **81**, 134509 (2010).
[21] L. Fidkowski and A. Kitaev, *Phys. Rev. B* **83**, 075103 (2011).
[22] D. Tong and C. Turner, *SciPost Phys. Lect. Notes* **14** (2020); arXiv:1906.07199.