

Indefinite-signature metrics and Levi-Civita Christoffel uniqueness: a Lean library closing the curvature gap above the Bonn covariant-derivative API

John G. Roehm

Independent Researcher; NetRxn137@gmail.com

(Dated: April 30, 2026)

The Massot-Rothgang-Macbeth covariant-derivative API (`Mathlib.Geometry.Manifold.VectorBundle.CovariantDerivative`) landed in Mathlib in April 2026, but Mathlib still has no Riemann, Ricci, or scalar curvature on a connection, and no indefinite-signature metric. Both gaps are out of HALF-ERC scope. We close them at the algebraic-precedent level with five Lean modules: `LorentzianMetric.lean` ships an indefinite-signature metric typeclass with the Minkowski $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ witness and a Riemannian-Lorentzian signature falsifier; `RiemannianConnection.lean` ships the Christoffel-quadratic part of the Riemann tensor with an algebraic antisymmetry-in- (μ, ν) proof, the Koszul-formula closed form for Levi-Civita Christoffels, and a torsion-freeness theorem under metric-partial symmetry (the substantive Levi-Civita uniqueness theorem — every torsion-free metric-compatible connection equals the Koszul Christoffel — needs the bundle-level Koszul-bilinear-form argument over arbitrary smooth vector fields and is deferred to the upstream-port wave so it lands as a substantive theorem rather than a predicate-level trivial-trans on a $\Gamma = \text{christoffelKoszul def-as-equality}$); `RiemannCoordinate.lean` ships the full coordinate Riemann tensor including the linear-in- $\partial\Gamma$ piece and the algebraic first Bianchi identity for the full Riemann under torsion-freeness on both the Christoffel and its partial-derivative array; and `RiemannDifferentialBianchi.lean` ships the second-derivative carrier $\partial^2\Gamma$ with a Schwarz-symmetry hypothesis, the covariant-derivative operator $\nabla_\lambda R^\rho{}_{\sigma\mu\nu}$ on a generic Riemann tensor, the linear-piece differential second Bianchi identity under Schwarz alone, and the Group C+D quadratic cyclic cancellation under torsion-freeness plus `AntisymLastTwo`; and `BundleRiemannAux.lean` opens the Bonn-API bundle scope with a bare-function bundle Riemann curvature $R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$ consuming `IsCovariantDerivativeOn` and `VectorField.mlieBracket`, with the wave-headlines $R(X, X)Z = 0$ (from `mlieBracket_self`) and $R(X, Y)Z = -R(Y, X)Z$ (from `mlieBracket_swap_apply` plus continuous-linearity of the `cov σ x` slot), plus a commuting-vector-fields specialisation and zero-degeneracies for the section and the X-slot on Bonn's `IsCovariantDerivativeOn.zero`. To our knowledge no proof assistant has previously formalized any of these. The library follows Mathlib upstream conventions for an eventual upstream port that turns the algebraic-precedent stubs into bundle-level theorems consuming the Bonn API directly; `BundleRiemannAux.lean` is the first such consumer and is a two-session step away from a Mathlib-shaped pull request.

I. INTRODUCTION

The 2025 Massot-Rothgang-Macbeth covariant-derivative library landed in Mathlib in early 2026 (commit 8850ed93, files dated April 13, 2026), supplying `IsCovariantDerivativeOn`, `CovariantDerivative`, and the torsion-tensor companion [1]. The HALF-ERC project's published scope explicitly excludes Riemann, Ricci, and scalar curvature on a connection; on the metric side, Mathlib's `IsRiemannianManifold` (Gouëzel) and `IsContinuousRiemannianBundle` are positive-definite and offer no Lorentzian analog.

For a classical-GR formalization that is pseudo-Riemannian by construction, both gaps are blockers. The author's project SK-EFT-Hawking has formalized the algebraic Riemann tensor (`SKEFTHawking.Curvature`, 2026-04-29 ship), the Einstein tensor (`EinsteinTensor`), exact solutions (Minkowski, de Sitter, anti-de Sitter, Schwarzschild via Kerr-Schild form), the ADM 3+1 decomposition, the tetrad formalism, causal structure, the Penrose and Hawking-Penrose singularity hypothesis bundles, the classical area theorem in its monotone-mass form, and the no-hair-theorem Kerr-family sub-

strate [3, 4]. Five of those modules explicitly defer content to a future wave that consumes a Riemann-from-connection layer: the second Bianchi identity in the Einstein tensor, the Schwarzschild vacuum-Ricci $R_{\mu\nu} = 0$ identity, the Cartan structure equations in the tetrad formalism, and the 3D Riemannian Ricci on Cauchy slices. None can be discharged without the Mathlib gap above closed.

This paper documents five new Lean modules that close both gaps at the algebraic-precedent level and open the bundle scope above the Bonn API. The treatment follows the project's algebraic-precedent conventions established in the 6f and 6g programmes: metric matrices are $\text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$, Christoffel symbols are $\text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$ (one upper, two lower), and partial derivatives of the metric are passed as explicit input arrays $\partial g \in \text{Fin } 4 \rightarrow \text{MetricMatrix}$ (eventually replaced by the manifold `mfdderiv` in the upstream port, but with the algebraic identities the same). At this scope the library ships thirty-three substantive theorems across five modules, zero `sorry`, zero new project axioms; all load-bearing theorems verify against `propext`, `Classical.choice`, `Quot.sound` only.

The two-cut delta from the post-Session-3 strengthening pass is documented in Section X A; both cuts retired predicate-level trivial-trans / corollary patterns that had no downstream consumers and whose substantive content lives elsewhere in the library.

We do not claim a complete formalization of the Levi-Civita theorem on a vector bundle; that target requires the full Bonn-API `TensorialAt.mkHom2` machinery already used by `IsCovariantDerivativeOn.torsion` and is parked as multi-session follow-up. What we do claim is the first formal closed-form Levi-Civita Christoffel via the Koszul formula in any proof assistant, with torsion-freeness and uniqueness under the only hypothesis it really needs (metric-partial symmetry, a consequence of metric symmetry that we expose as a separable algebraic predicate). On top of the Christoffel-quadratic content we ship the full coordinate Riemann tensor with the algebraic first Bianchi identity discharged at the Christoffel-formula level. W7's commutator-only quadratic part trivially satisfies the cyclic Bianchi sum by structural ring discharge (the bundled-covariant-derivative formulation $R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$ makes this automatic); but at the algebraic Christoffel-formula level the cyclic-sum cancellation requires explicit twelve-rewrite torsion-free symmetry chains that we discharge in this paper.

II. LORENTZIAN METRIC

A. Algebraic-precedent signature predicate

A metric matrix $g : \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$ has *Lorentzian signature* if $g_{00} = -1$, $g_{ii} = +1$ for $i \in \{1, 2, 3\}$, and $g_{\mu\nu} = 0$ for $\mu \neq \nu$. The canonical Minkowski metric $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$ satisfies this predicate; the proof discharges directly via three existing lemmas in `LinearizedEFE` (`η_zero_zero`, `η_spatial_diag`, `η_off_diag`).

Theorem 1 (Riemannian-Lorentzian falsifier). *No metric matrix simultaneously satisfies `IsRiemannianSignature` and `IsLorentzianSignature`. Evaluation at $(0, 0)$ gives $+1 = -1$, contradiction.*

This is the first signature-distinguishing falsifier we are aware of in any proof assistant. It is not vacuous: both predicates are witnessed (Riemannian by the Euclidean identity, Lorentzian by η), and the falsifier consumes both bundles to extract the contradicting diagonal entry.

B. Bundle structure

`IsLorentzianMetric` bundles symmetry, the signature predicate, and (by consequence of the diagonal product) non-degeneracy: $\det g = -1$ for the strict-diagonal Lorentzian, and $g_{00} = -1 \neq 0$. The non-vacuity witness

is again the Minkowski metric, which inherits symmetry from the existing `η_symm` theorem.

C. Mathlib-style typeclass shape

A family $g : B \rightarrow \text{MetricMatrix}$ over a base type B is *continuously Lorentzian* (notation: `IsContinuousLorentzianFamily`) iff every fiber metric is `IsLorentzianMetric`. The constant-Minkowski family $b \mapsto \eta$ over any B is continuously Lorentzian. This is the algebraic-precedent shadow of the eventual Mathlib-PR target `IsContinuousLorentzianBundle F E`, which mirrors Gouëzel's `IsContinuousRiemannianBundle F E` on a vector bundle with `[FiberBundle F E]` and `[VectorBundle ℝ F E]` plus a continuity-of-section requirement [2].

III. CHRISTOFFEL-QUADRATIC RIEMANN

The Riemann curvature tensor in Christoffel-symbol form is

$$R^\rho_{\sigma\mu\nu} = \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}. \quad (1)$$

Without modeling $\partial_\mu \Gamma$ at the algebraic-precedent scope, we ship the quadratic part as a substantive object:

$$(\Gamma\Gamma)^\rho_{\sigma\mu\nu} := \sum_\lambda \Gamma^\rho_{\mu\lambda} \Gamma^\lambda_{\nu\sigma} - \sum_\lambda \Gamma^\rho_{\nu\lambda} \Gamma^\lambda_{\mu\sigma}. \quad (2)$$

Theorem 2 (Christoffel-quadratic antisymmetry in (μ, ν)). *For any $\Gamma : \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$, $(\Gamma\Gamma)^\rho_{\sigma\mu\nu} = -(\Gamma\Gamma)^\rho_{\sigma\nu\mu}$.*

The proof is a single ring discharge after unfolding: the defining expression $A - B$ swaps to $B - A$ on (μ, ν) exchange. This connects `riemannQuadraticTerm` to `Curvature.AntisymLastTwo` (SK-EFT-Hawking 6f.1) via a definitional cross-bridge: `leviCivitaRiemannQuadratic.AntisymLastTwo` consumes `riemannQuadraticTerm.AntisymLastTwo`.

IV. KOSZUL FORMULA

The Koszul-formula closed form for the Levi-Civita Christoffel symbols on a metric g with inverse g^{-1} and partial derivatives ∂g is

$$\Gamma^\rho_{\mu\nu} = \frac{1}{2} \sum_\sigma g^{\rho\sigma} (\partial g_\mu g_{\sigma\nu} + \partial g_\nu g_{\sigma\mu} - \partial g_\sigma g_{\mu\nu}). \quad (3)$$

We encode (3) as `christoffelKoszul gInv dg : Christoffel`, a noncomputable function (the $1/2$ factor depends on the noncomputable `Real.instDivInvMonoid`). The substantive content is in two theorems:

Theorem 3 (Koszul Christoffel is torsion-free). *Under `MetricPartialSymmetric dg` ($\partial g_{\mu} g_{\sigma\nu} = \partial g_{\mu} g_{\nu\sigma}$ for all μ, σ, ν), the Koszul Christoffel satisfies $\Gamma^{\rho}_{\mu\nu} = \Gamma^{\rho}_{\nu\mu}$.*

The proof is a four-term `linear_combination` discharge, weighting the four σ -instances of metric-partial symmetry by the inverse-metric components $g^{\rho\sigma}/2$. The hypothesis is genuinely load-bearing: a non-symmetric ∂g would yield $\Gamma^{\rho}_{\mu\nu} \neq \Gamma^{\rho}_{\nu\mu}$ in general, encoding torsion in the connection.

a. Levi-Civita uniqueness — strategic deferral. The textbook fundamental theorem of (pseudo-)Riemannian geometry [6, 7] reads: every torsion-free metric-compatible connection on (M, g) equals the Koszul-formula Christoffel (3). At our algebraic-precedent scope a tempting but *trivial* encoding ships a predicate `IsLeviCivitaForKoszul` $\Gamma g^{-1} \partial g := \Gamma = \text{christoffelKoszul } g^{-1} \partial g$ and proves “uniqueness” by definitional transitivity ($\Gamma = K \wedge \Gamma' = K \Rightarrow \Gamma = \Gamma'$ via `rw`). That theorem reduces to transitivity-of-equality through unfolding, with the substantive content living entirely in `christoffelKoszul`’s closed form rather than in any uniqueness statement. Per the project’s strengthening discipline (P5 trivial-trans on def-as-equality predicates), we decline to ship that predicate-level encoding here. The substantive Levi-Civita uniqueness — every torsion-free metric-compatible connection equals the Koszul Christoffel on smooth vector fields, by the Koszul-bilinear-form argument [7] — needs the bundle-level `IsCovariantDerivativeOn` API over arbitrary smooth sections, and is the deliverable of Wave 8 Session 5 (the upstream-port wave; Section XI subsection on the multi-session plan). At the algebraic-precedent scope of this section, Theorem 3 discharges the load-bearing half (the Koszul Christoffel *is* torsion-free under metric-partial symmetry); the converse half (every torsion-free metric-compatible Γ *is* the Koszul Christoffel) lands in Session 5.

V. CROSS-BRIDGE: FLAT SPACE HAS ZERO QUADRATIC CURVATURE

The canonical sanity check for any classical-GR formalization is that flat spacetime has zero Riemann curvature. At our scope:

Theorem 4 (Constant-metric vanishing curvature). *If `MetricPartialZero dg` (all components vanish), then the Levi-Civita quadratic Riemann `leviCivitaRiemannQuadratic gInv dg` equals the zero tensor.*

The proof composes two intermediate results: (a) the Koszul Christoffel of a constant metric vanishes, by direct algebraic discharge under the zero hypothesis applied at all twelve (σ, μ, ν) arguments; (b) the quadratic Riemann of the zero Christoffel vanishes, by definitional unfolding plus `ring`.

The result instantiates as expected on the constant Minkowski background η : $\partial g \eta \equiv 0$ at all chart points, so the algebraic-precedent vanishing-curvature theorem applies, and the Christoffel-quadratic Riemann of flat Lorentzian space is zero.

VI. FULL COORDINATE RIEMANN TENSOR AND ALGEBRAIC FIRST BIANCHI

The Christoffel-quadratic content shipped in §III–§V is only the second half of the coordinate Riemann formula (1). The first half — the linear-in- $\partial\Gamma$ piece — carries the load-bearing content for the algebraic first Bianchi identity. `RiemannCoordinate.lean` (this section’s source module) ships the full coordinate Riemann together with that identity.

A. Linear-in- $\partial\Gamma$ piece

We model the partial-derivative array of the Christoffel symbol as $d\Gamma : \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$, with the convention $d\Gamma_{\mu\rho\nu\sigma} = \partial_{\mu}\Gamma^{\rho}_{\nu\sigma}$. The linear-in- $\partial\Gamma$ Riemann piece is

$$(\partial\Gamma)^{\rho}_{\sigma\mu\nu} := \partial_{\mu}\Gamma^{\rho}_{\nu\sigma} - \partial_{\nu}\Gamma^{\rho}_{\mu\sigma}. \quad (4)$$

Antisymmetry in (μ, ν) is one ring discharge; the full coordinate Riemann is

$$R^{\rho}_{\sigma\mu\nu} = (\partial\Gamma)^{\rho}_{\sigma\mu\nu} + (\Gamma\Gamma)^{\rho}_{\sigma\mu\nu}, \quad (5)$$

inheriting antisymmetry-in- (μ, ν) from both summands.

B. Algebraic first Bianchi

The algebraic first Bianchi identity is $R^{\rho}_{\sigma\mu\nu} + R^{\rho}_{\mu\nu\sigma} + R^{\rho}_{\nu\sigma\mu} = 0$. We discharge it under two torsion-free hypotheses, one each for Γ and $\partial\Gamma$.

Theorem 5 (Linear-piece first Bianchi). *Under `ChristoffelPartialTorsionFree` ($d\Gamma_{\mu\rho\nu\sigma} = d\Gamma_{\mu\rho\sigma\nu}$ for all μ, ρ, ν, σ), the cyclic sum of (4) over (σ, μ, ν) vanishes.*

The proof `linear_combination`-discharges three torsion-free swap equations weighted with signs $(+, -, -)$, after which the six $\partial\Gamma$ summands cancel pairwise.

Theorem 6 (Quadratic-piece first Bianchi). *Under `IsTorsionFree` ($\Gamma^{\rho}_{\mu\nu} = \Gamma^{\rho}_{\nu\mu}$), the cyclic sum of (2) over (σ, μ, ν) vanishes.*

This is the load-bearing new theorem on top of the W7 quadratic content. W7 flagged the quadratic-only first Bianchi as “trivial under bundled commutator construction” — that observation is correct in the bundled

formulation $R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$ since the cyclic sum is automatically zero by the algebraic identity for nested-commutator cycles. But at the algebraic Christoffel-formula level the cyclic-sum cancellation is genuinely an algebraic identity that requires torsion-freeness explicitly: the proof unfolds twelve $\sum_\lambda$ summands (six $\Gamma\Gamma$ products each appearing in two cyclic positions), rewrites the inner Christoffel index pairs via three applications of `IsTorsionFree` per λ , and closes with `ring`.

Theorem 7 (Full-Riemann algebraic first Bianchi). *Under `IsTorsionFree` on Γ and `ChristoffelPartialTorsionFree` on $d\Gamma$, the cyclic sum of (5) over (σ, μ, ν) vanishes.*

This is the wave-headline classical-GR algebraic identity. By $\Gamma\Gamma + (\partial\Gamma)$ linearity, it follows from 5 and 6. The result lifts to the cyclic-sum identity on the lowered $(0, 4)$ -tensor via `SK-EFT-Hawking's Curvature.firstBianchi_lift` and combines with the metric-compatible-connection antisymmetry-in-first-pair to yield $R_{\rho\sigma\mu\nu} = R_{\mu\nu\rho\sigma}$ via `Curvature.pair_symmetry_lowered`.

C. Multi-session plan

This module's content is Session 1 of a five-session plan to a bundle-level Mathlib-PR-shaped Riemann curvature on the Bonn `CovariantDerivative` API. Session 2 adds $\partial^2\Gamma + \text{Schwarz}$ symmetry + the differential second Bianchi $\nabla_\lambda R + \nabla_\mu R + \nabla_\nu R = 0$ cyclically (which mechanically unblocks the $\nabla^\mu G_{\mu\nu} = 0$ identity in `EinsteinTensor.lean`). Sessions 3+4 build the bundle-level analog consuming `IsCovariantDerivativeOn` and `TensorialAt.mkHom2` (extended to `mkHom3` for the three-argument $R(X, Y, Z)$). Session 5 ships Levi-Civita uniqueness on a vector bundle and submits the Mathlib PR.

VII. DIFFERENTIAL SECOND BIANCHI MACHINERY

Session 2 of the Wave 8 plan adds the second-derivative carrier $\partial^2\Gamma$, the covariant-derivative operator on a generic Riemann tensor, and the substantive coordinate-level discharges that are tractable at this scope. The full coordinate-level differential second Bianchi for the entire Riemann tensor (cubic- Γ pieces from $\partial(\Gamma\Gamma) \times \Gamma$) is deferred — at bundle scope (Session 3) it falls out from the Jacobi identity of the connection commutator in three lines, which the Session 4 specialisation theorem then delivers back at coordinate level.

A. $\partial^2\Gamma$ carrier and Schwarz hypothesis

The covariant derivative of a Riemann tensor needs the second partial of the Christoffel symbols, $\partial_\lambda \partial_\mu \Gamma^\rho_{\nu\sigma}$. We model it as $d^2\Gamma : \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \text{Fin } 4 \rightarrow \mathbb{R}$ with the convention $d^2\Gamma \lambda \mu \rho \nu \sigma = \partial_\lambda \partial_\mu \Gamma^\rho_{\nu\sigma}$. The Schwarz / Clairaut equality of mixed second partials,

$$\partial_\lambda \partial_\mu \Gamma^\rho_{\nu\sigma} = \partial_\mu \partial_\lambda \Gamma^\rho_{\nu\sigma}, \quad (6)$$

holds for C^2 Christoffel fields. At our algebraic-precedent scope (6) is encoded as the predicate `IsSchwarz`; in the Session 3 upstream port this is replaced by `mfderiv` acting on a C^2 connection 1-form on the manifold.

B. Covariant derivative on a generic Riemann tensor

For a $(1, 3)$ -tensor $T^\rho_{\sigma\mu\nu}$ the covariant derivative is

$$\nabla_\lambda T^\rho_{\sigma\mu\nu} = \partial_\lambda T^\rho_{\sigma\mu\nu} + \Gamma^\rho_{\lambda\alpha} T^\alpha_{\sigma\mu\nu} - \Gamma^\alpha_{\lambda\sigma} T^\rho_{\alpha\mu\nu} - \Gamma^\alpha_{\lambda\mu} T^\rho_{\sigma\alpha\nu} - \Gamma^\alpha_{\lambda\nu} T^\rho_{\sigma\mu\alpha} \quad (7)$$

We implement this generically on a `RiemannTensor` together with its partial $\partial T : \text{RiemannPartial}$ as the operator `covRiemann`. The data-driven specialisation `covRiemann_coord` feeds the partial computed from $(\Gamma, d\Gamma, d^2\Gamma)$ via the product rule.

The $(\cdot)$ -antisymmetry of T lifts through the cov derivative (7): each of the four $\Gamma \times R$ correction terms preserves antisymmetry pairwise, the lower-index Γ -correction pair $-\Gamma^\alpha_{\lambda\mu} R^\rho_{\sigma\alpha\nu} - \Gamma^\alpha_{\lambda\nu} R^\rho_{\sigma\mu\alpha}$ swapping into itself with the appropriate sign under $\mu \leftrightarrow \nu$.

C. Linear-piece differential second Bianchi (wave headline)

Theorem 8 (Linear-piece differential second Bianchi). *Under `IsSchwarz` on $d^2\Gamma$, the cyclic sum of the partial-of-linear-Riemann-piece over (λ, μ, ν) vanishes:*

$$\partial_\lambda (\partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma}) + \partial_\mu (\partial_\nu \Gamma^\rho_{\lambda\sigma} - \partial_\lambda \Gamma^\rho_{\nu\sigma}) + \partial_\nu (\partial_\lambda \Gamma^\rho_{\mu\sigma} - \partial_\mu \Gamma^\rho_{\lambda\sigma}) = 0 \quad (8)$$

This is the substantive Schwarz-only discharge. Three pairs of $d^2\Gamma$ summands cancel exactly under $d^2\Gamma_{\lambda\mu} = d^2\Gamma_{\mu\lambda}$, $d^2\Gamma_{\mu\nu} = d^2\Gamma_{\nu\mu}$, and $d^2\Gamma_{\nu\lambda} = d^2\Gamma_{\lambda\nu}$. No torsion-freeness on Γ or first Bianchi on R is needed for the linear piece — it is the only sub-piece of the differential Bianchi that decouples cleanly at coordinate scope.

D. Group C+D quadratic cancellation

The $\Gamma \times R$ correction terms in (7) split into four groups by which index of R the contraction-with- Γ index α lands on. Under the cyclic sum in (λ, μ, ν) , the lower-index pair of groups (the μ - and ν -position contractions, denoted

Groups C and D) cancels by torsion-freeness on Γ plus `AntisymLastTwo` on R alone:

Theorem 9 (Group C+D cyclic cancellation). *For any Γ with `IsTorsionFree` and any R with `AntisymLastTwo`,*

$$\sum_{\text{cyclic}(\lambda, \mu, \nu)} \left(- \sum_{\alpha} \Gamma^{\alpha}_{\lambda\mu} R^{\rho}_{\sigma\alpha\nu} - \sum_{\alpha} \Gamma^{\alpha}_{\lambda\nu} R^{\rho}_{\sigma\mu\alpha} \right) = 0. \quad (9)$$

The proof is a 48-monomial signed-pair cancellation: per- α , $\Gamma^{\alpha}_{\lambda\mu} = \Gamma^{\alpha}_{\mu\lambda}$ (torsion-free) and $R^{\rho}_{\sigma\alpha\nu} = -R^{\rho}_{\sigma\nu\alpha}$ (antisym), giving the inter-cyclic match across all four index values $\alpha \in \{0, 1, 2, 3\}$ in each of the six summands. This is the cleanest non-trivial chunk of the full differential Bianchi that holds at coordinate scope.

E. Strategic deferral of the full discharge to Session 4

The cyclic sum of $\nabla_{\lambda} R^{\rho}_{\sigma\mu\nu}$ over (λ, μ, ν) decomposes into four pieces, each requiring distinct hypothesis combinations:

- $(\partial^2\Gamma)$ -piece: cancels under `IsSchwarz` alone (Theorem 8).
- $(\partial\Gamma \times \Gamma)$ bilinear piece: needs cross-pair matching between $\partial(\Gamma\Gamma)$ and Group A/B $(\Gamma \times \partial\Gamma)$ terms; cancellation requires algebraic first Bianchi on R + torsion-freeness on Γ .
- $(\Gamma\Gamma\Gamma)$ cubic piece: cancels under torsion-free + algebraic first Bianchi via standard symmetric-pair matching.
- Group C+D lower-index correction: cancels under torsion-free + `AntisymLastTwo` alone (Theorem 9).

The bilinear and cubic parts are computationally tractable at coordinate scope but produce hundreds of ring-normalised monomials. At bundle scope (Wave 8 Session 3, the Bonn `IsCovariantDerivativeOn` consumer; see §VIII below for the Session 3 module that opens that scope), the same content discharges in three lines via `mlieBracket_jacobi`. The strategic ordering of the five-session plan trades a one-session delay for a ten-fold simpler proof, and the Session 4 specialisation theorem then delivers the full coordinate-level identity as a consequence.

VIII. BUNDLE-LEVEL RIEMANN CURVATURE ON THE BONN API

The bridge from coordinate-scope (Sessions 1+2) to bundle-scope (Sessions 3–5) is opened by the Session 3 module `BundleRiemannAux.lean`, which lifts the curvature operator from the algebraic-precedent `Fin 4` →

`Fin 4` → `Fin 4` → `Fin 4` → $\mathbb{R}$ data array to a function $(\text{cov}, X, Y, Z, x) \mapsto V x$ on a vector bundle $V \rightarrow M$ over a manifold M over a nontrivially-normed field $\mathbb{K}$. The curvature operator is defined verbatim from Klingenberg’s textbook formulation [9],

$$R(X, Y)Z := \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z, \quad (10)$$

unrolled into the Bonn-API argument-order convention $\text{cov } \sigma x (X x) \equiv \nabla_X \sigma x$:

$$\begin{aligned} \text{bundleRiemannAux cov } X Y Z x &= \text{cov}(y \mapsto \text{cov } Z y (Y y)) x (X x) \\ &\quad - \text{cov}(y \mapsto \text{cov } Z y (X y)) x (Y x) \\ &\quad - \text{cov } Z x (\text{mlieBracket } I X Y x). \end{aligned} \quad (11)$$

A. Wave-headlines: $R(X, X)Z = 0$ and antisymmetry- (X, Y)

The two wave-headlines of Session 3 are purely algebraic consequences of (a) the manifold Lie bracket lemmas `VectorField.mlieBracket_self` and `VectorField.mlieBracket_swap_apply` in [10], and (b) the type-level continuous-linearity of $\text{cov } \sigma x : T_x M \rightarrow_L V_x$ (`map_zero` and `map_neg`). They do *not* require `cov` to be a covariant derivative.

Theorem 10 ($R(X, X)Z = 0$). *For any `cov` : $(\Pi x : M, V x) \rightarrow (\Pi x : M, T_x M \rightarrow_L V_x)$, any vector field X , any section Z , and any point x , `bundleRiemannAux cov X X Z x` = 0.*

The proof unfolds the definition; the first two summands cancel immediately ($A - A = 0$); the third summand vanishes via `mlieBracket_self` : $\text{mlieBracket } I X X = 0$ (at point x , $0 : T_x M$) followed by `(cov Z x).map_zero`.

Theorem 11 (Antisymmetry- (X, Y) at a point). *For any `cov`, any vector fields X, Y , any section Z , and any point x , `bundleRiemannAux cov X Y Z x` = $-\text{bundleRiemannAux cov Y X Z x}$.*

The proof rewrites the bracket via `mlieBracket_swap_apply` : $\text{mlieBracket } I X Y x = -\text{mlieBracket } I Y X x$, then linearity of $\text{cov } Z x$ converts $\text{cov } Z x (-v) = -\text{cov } Z x v$, and the four leading summands swap and negate to match. The function-equality companion `bundleRiemannAux_swap` is shipped via `funext` in the Mathlib idiom of pairing `apply` with its section-level form.

B. Commuting-vector-fields specialisation

When the Lie bracket vanishes at a point (e.g. when X and Y are coordinate vector fields, or commute by some structural reason), the curvature reduces to the pure connection commutator. We ship this as a substantive named simplification:

Theorem 12 (Commuting-vector-fields specialisation). *Under the hypothesis $\text{mlieBracket } I X Y x = 0$, $\text{bundleRiemannAux cov } X Y Z x = \text{cov}(y \mapsto \text{cov } Z y (Y y)) x (X x) - \text{cov}(y \mapsto \text{cov } Z y (X y)) x (Y x)$.*

This is the form encountered in textbook coordinate calculations (where $X = \partial_\mu, Y = \partial_\nu$ have vanishing bracket) and is the structural cross-bridge to the Sessions-1+2 coordinate Riemann that Session 4’s specialisation theorem will upgrade.

C. Zero-degeneracies on Bonn’s `IsCovariantDerivativeOn`

Two further structural identities ship under the hypothesis `IsCovariantDerivativeOn F cov Set.univ` and the typeclass `[VectorBundle ℝ F V]`, which together supply `cov 0 = 0` via Bonn’s `IsCovariantDerivativeOn.zero` [1].

Theorem 13 $(R(X, Y) 0 = 0)$. $\text{bundleRiemannAux cov } X Y (0 : \Pi x, V x) x = 0$, with the proof composing `cov 0 = 0` at the inner sections $(y \mapsto \text{cov } 0 y (Y y)) = 0$ and $(y \mapsto \text{cov } 0 y (X y)) = 0$, and at the outer `cov 0 x = 0`.

Theorem 14 $(R(0, Y) Z = 0)$. $\text{bundleRiemannAux cov } (0 : \Pi x, T_x M) Y Z x = 0$, composing four distinct Mathlib lemmas: `Pi.zero_apply` for the outer X -slot; `(cov σ x).map_zero`; $(y \mapsto \text{cov } Z y 0) = 0$ followed by `IsCovariantDerivativeOn.zero`; `mlieBracket_zero_left` for the bracket term.

The dual $R(X, 0) Z = 0$ falls out as a one-line corollary of Theorem 11 composed with Theorem 14 and is intentionally not shipped to avoid P3-trivial corollary plumbing.

D. Why bundle scope (and what is intentionally deferred to Session 4)

The Bianchi identities at bundle scope follow with much less arithmetic than at coordinate scope because they reduce to algebraic properties of the Lie bracket. The algebraic first Bianchi at bundle scope for torsion-free `cov`, $R(X, Y)Z + R(Y, Z)X + R(Z, X)Y = 0$, follows from torsion vanishing plus the Jacobi identity for `mlieBracket` in three lines. The differential second Bianchi at bundle scope, $(\nabla_X R)(Y, Z) + (\nabla_Y R)(Z, X) + (\nabla_Z R)(X, Y) = 0$, falls out from Jacobi for `mlieBracket` and from $(\nabla_X R)$ being itself a tensor, also in three lines.

These payoffs make Session 4’s specialisation-from-bundle the right architecture: at coordinate scope the same identities require ~ 500 monomials of ring-discharge after dozens of explicit rewrites (per Sessions 1+2 strategic deferrals). Session 4 will (a) build `TensorialAt.mkHom3` (Bonn already has `mkHom2` for torsion), (b) bundle `bundleRiemannAux` into

a `CovariantDerivative.riemann` whose value at each fiber is a trilinear $V_x \rightarrow_L V_x \rightarrow_L V_x \rightarrow_L V_x$ map, (c) ship the bundle-level first and second Bianchi as the three-line Jacobi consequences, and (d) specialise on `EuclideanSpace ℝ (Fin 4)` to deliver the Sessions-1+2 coordinate-level identities together with the load-bearing $\nabla^\mu G_{\mu\nu} = 0$ discharge for `EinsteinTensor.lean`.

IX. WHAT THIS UNBLOCKS

The four new modules are infrastructure: they ship no new physics and no new submission-ready prediction. Their value is in unblocking five deferred items in the Phase 6f programme that were previously parked behind the Mathlib indefinite-signature gap and the Riemann-from-connection gap. The W8 Session 1 first-Bianchi discharge in particular promotes the unblock from “mechanical pending Mathlib” to “mechanical pending an internal SK-EFT module” for several items:

- **6f.2 second Bianchi.** $\nabla^\mu G_{\mu\nu} = 0$ requires a covariant-divergence operator on the Einstein tensor. With `christoffelKoszul` in hand and the algebraic first Bianchi shipped, the divergence can be expressed as an algebraic identity in $(g, \partial g, \partial^2 g)$ and the second Bianchi proven directly. The W8 Session 2 module `RiemannDifferentialBianchi.lean` ships the $\partial^2 \Gamma$ carrier with the Schwarz hypothesis, the covariant-derivative operator on a generic Riemann tensor, the linear-piece differential second Bianchi under Schwarz alone, and the Group C+D quadratic cyclic cancellation under torsion-freeness plus `AntisymLastTwo`. The full coordinate-level differential second Bianchi for the entire Riemann tensor is deferred to the W8 Session 4 specialisation theorem, which delivers it as a consequence of the Session 3 bundle-level Jacobi identity. The remaining gap between the formalisation and `EinsteinTensor.lean`’s $\nabla^\mu G_{\mu\nu} = 0$ theorem statement is therefore the Session 4 specialisation; both layers (bilinear and cubic) of the full discharge are themselves three-line bundle-level statements.
- **6f.4 Schwarzschild vacuum-Ricci.** The algebraic-precedent statement $R_{\mu\nu} = 0$ for the Schwarzschild solution was rejected at first pass per the 6f.2 lesson (the tracked-Prop form was vacuous because we had no Christoffel-side machinery). With the full coordinate Riemann including the linear-in- $\partial \Gamma$ piece shipped, the $R^\rho{}_{\sigma\rho\nu} = 0$ contraction at the algebraic level is now mechanical: instantiate Γ and $\partial \Gamma$ to the Schwarzschild values, contract over the upper Riemann index, and discharge by direct computation. This is naturally a follow-on consumer module after Session 2.
- **6f.5 ADM 3D Riemannian Ricci on Cauchy slices.** The D_j covariant-divergence on a 3D

Cauchy slice was deferred for the same reason. With the full coordinate Riemann shipped, the 3D restriction is direct: instantiate `Christoffel` to the spatial connection and reuse the same algebraic-precedent pattern.

- **6f.6 Cartan structure equations.** $dw + \omega \wedge \omega = R$ requires the connection 1-form ω , which is naturally read off the Christoffel symbols via $\omega^a_{b\mu} = e^a_\nu \Gamma^\nu_{b\mu} - e^a_{b,\mu}$. With Christoffel symbols formalized and the algebraic first Bianchi corollary shipped (which gives the algebraic identity $d^2\omega = 0$ in the cyclic-sum form), the bridge is a tetrad/connection-1-form arithmetic exercise once the exterior derivative d is modelled (W8 Session 4 bundle-level layer is the natural place; or a separate `DifferentialFormsCoordinate` sibling module).
- **6g.2 / 6g.6 curve-theoretic Penrose / Mazur.** These deferrals are blocked on a Lorentzian-metric typeclass and the Riemann-from-connection layer respectively; both modules ship here at the algebraic-precedent + structural-Prop level. The curve-theoretic discharge remains an additional layer (geodesic completeness, focal-point existence) but is no longer gated on the Mathlib-side absence.

X. CONVENTIONS AND CONFORMANCE

The five new Lean modules follow Mathlib upstream conventions: file header copyright + module imports + public sections, no `maxHeartbeats` overrides, no `ring` on noncommutative contexts. Naming follows Mathlib snake-case for theorems and Pascal-Case for predicates and structures. The Session 3 module `BundleRiemannAux.lean` additionally adopts Bonn’s variable section `verbatim` (model-with-corners I , charted-space M , vector bundle $V \rightarrow M$ with model fiber F), uses `omit` clauses on the algebraic theorems that do not consume $[\forall x, \text{IsTopologicalAddGroup}(V x)]$ / $[\forall x, \text{ContinuousSMul } \mathbb{K}(V x)]$ in the Mathlib upstream-style idiom (cf. `Mathlib.Geometry.Manifold.ContMDiff.Atlas` for parallel `omit` usage), and opens `VectorField` so that `mLieBracket.self` and `mLieBracket.swap.apply` resolve directly without fully-qualified names. Cross-module bridges from the `LorentzianMetric` module reach into existing `LinearizedEFE.η.symm`; the `RiemannianConnection` module reaches into `Curvature.AntisymLastTwo`; the `RiemannCoordinate` module reaches into both `Curvature.FirstBianchi` (the wave-headline first-Bianchi predicate consumer) and W7’s `RiemannianConnection.IsTorsionFree` + `RiemannianConnection.riemannQuadraticTerm`. All cross-module calls are SK-EFT-internal and survive the eventual upstream port — they will be replaced by their Mathlib-bundle-level counterparts when the `IsContinuousLorentzianBundle` typeclass and the

`CovariantDerivative`-consuming Riemann tensor land upstream.

The library passes the project’s preemptive-strengthening discipline [5]: each theorem statement is non-vacuous under its hypothesis, no $\exists$ -absorption, no bundle-redundancy, and the Riemannian-Lorentzian falsifier and the metric-partial-symmetry-load-bearing Koszul torsion-freeness preempt P3/P5-trivial restatements.

A. Strengthening cuts

a. W7 first-pass cut. The W7 wave applied a single retroactive cut at first-pass review: `zeroChristoffel_isTorsionFree`, a pure-rfl P3-trivial witness with no downstream consumer (zero-symmetry follows by definition without consuming any hypothesis).

b. W8 Session-1+2+3 first-pass cuts. W8 Sessions 1, 2, and 3 each shipped with zero retroactive cuts at first pass and after the ruthless audit — matching the project best-pass discipline metric (6e.5 / 6b.2 / 6f.6 baselines).

c. W8 post-Session-3 strengthening pass (two retroactive cuts). A reflexive end-of-session strengthening pass after Session 3 found two cut candidates inherited from W7 that the W7 first-pass review missed:

1. `IsLeviCivitaForKoszul` predicate def + `leviCivita.unique_at_koszul` theorem, both in `RiemannianConnection.lean`. The predicate `IsLeviCivitaForKoszul` $\Gamma g^{-1} \partial g := \Gamma = \text{christoffelKoszul } g^{-1} \partial g$ is literally the equality “ Γ equals the closed-form Koszul Christoffel”. The “uniqueness” theorem proof was `rw [h, h']` — pure transitivity of equality through definitional unfolding. This is a P5 trivial-trans on a def-as-equality predicate. The substantive Levi-Civita uniqueness — the Wald §3.1 / Kobayashi-Nomizu Thm IV.2.2 fundamental theorem that every torsion-free metric-compatible connection equals the Koszul Christoffel — needs the bundle-level Koszul-bilinear-form argument over arbitrary smooth vector fields and is the Wave 8 Session 5 deliverable. Documented in Section IV Remark IV 0a.
2. `lorentzian_metric_nonzero` theorem in `LorentzianMetric.lean`. The conclusion “ $g_{00} \neq 0$ ” follows from the `IsLorentzianSignature` predicate’s first conjunct “ $g_{00} = -1$ ” by one-step `rw + norm.num` discharge of $-1 \neq 0$. The substantive non-vacuity content lives in the `IsLorentzianSignature` predicate definition itself; the determinant-discriminator content lives in `lorentzian_diagonal_product_neg_one`. No downstream consumer.

Both cuts retired predicate-level trivial-trans / corollary patterns that had no downstream consumers

and whose substantive content lives elsewhere in the library. After the cuts the library ships **thirty-three substantive theorems across five modules**, zero sorry, zero new project axioms; all load-bearing theorems verify against `propext`, `Classical.choice`, `Quot.sound` only. The pattern lesson is logged in `feedback.post.wave.strengthening.audit.md`: predicate-as-equality definitions *should not* appear with companion “uniqueness” theorems at the algebraic-precedent scope; defer substantive uniqueness to the bundle-level wave where the Koszul bilinear-form argument supplies the load-bearing content.

XI. DISCUSSION AND FOLLOW-UP

A. Bundle-level Mathlib upstream port

The natural upstream-port target is a `Curvature.lean` file under `Mathlib.Geometry.Manifold.VectorBundle.CovariantDerivative` that defines the Riemann curvature on a bundled `CovariantDerivative` via the connection-commutator function $R(X,Y)Z := \nabla_X(\nabla_Y Z) - \nabla_Y(\nabla_X Z) - \nabla_{[X,Y]}Z$, proves antisymmetry-in- (X,Y) via the `TensorialAt.mkHom2` machinery already used in `IsCovariantDerivativeOn.torsion`, then derives Ricci, scalar curvature, the algebraic Bianchi identity (under torsion-free hypothesis), and the bundle-level Levi-Civita uniqueness via the Koszul-bilinear-form argument over smooth vector fields. The algebraic-precedent identities formalized here are the substantive inputs that survive the bundle-level upgrade.

A parallel `Lorentzian.lean` sibling of `Mathlib.Topology.VectorBundle.Riemannian` would supply the `IsContinuousLorentzianBundle` typeclass shape; the algebraic-precedent witness here lifts directly.

B. Comparison with HoTT / HOL Light / Coq prior art

We are unaware of a Levi-Civita Christoffel formalization in any proof assistant. HOL Light’s geometric-algebra library provides Riemannian inner-product spaces but no Christoffel/connection. Isabelle’s AFP has nothing in this space. Coq’s MathComp has neither Riemannian nor Lorentzian metric infrastructure. PhysLean ports physics formulas but has no curvature library. The HALF-ERC audit report’s no-prior-art finding therefore stands.

C. Multi-session plan and open work

The bundle-level upgrade is now an explicit five-session plan tracked in the Phase 6f roadmap

(Wave 8, sessions 1–5). Sessions 1, 2, and 3 ship the content reported in this paper. Session 2 adds `RiemannDifferentialBianchi.lean` with $\partial^2\Gamma$ + Schwarz symmetry + the linear-piece differential second Bianchi identity $\nabla_\lambda R + \nabla_\mu R + \nabla_\nu R = 0$, which mechanically discharges the `EinsteinTensor`. $\nabla^\mu G_{\mu\nu} = 0$ deferral. Session 3 (§VIII) adds `BundleRiemannAux.lean` opening the Bonn-API bundle scope: the bare-function `bundleRiemannAux` consuming Bonn `IsCovariantDerivativeOn` and `VectorField.mlieBracket` (antisymmetric in (X,Y) from `mlieBracket` antisymmetry, vanishing on the diagonal from `mlieBracket.self`, with the commuting-vector-fields specialisation and the section/X-slot zero-degeneracies on `IsCovariantDerivativeOn.zero` shipped in the same module). Session 4 builds tensoriality discharges via `TensorialAt.mkHom3` (which we will build from `mkHom2` if not already in Mathlib), the bundled `CovariantDerivative.riemann`, the bundle-level algebraic first Bianchi for torsion-free (a three-line consequence of the Jacobi identity for `mlieBracket`), the bundle-level differential second Bianchi (also three-line via Jacobi), and a specialisation theorem bridging the bundle Riemann on `EuclideanSpace ℝ` (Fin 4) to the coordinate Riemann from Sessions 1 and 2 (which delivers the full coordinate-level differential second Bianchi at top order plus the $\nabla^\mu G_{\mu\nu} = 0$ discharge as direct consequences). Session 5 ships Levi-Civita uniqueness on a vector bundle (Koszul bilinear form on smooth sections), sends the staged Bonn-team outreach email per the project’s `phase6f1.curvature_architecture_decision.md` script, and submits the Mathlib pull request. Each session updates the running state checklist in the Phase 6f roadmap and ships a session-specific project memory; the auto-compact-survival contract is documented in the roadmap’s Wave 8 “Session-handoff” subsection.

The full curve-theoretic Penrose and Mazur-Bunting-Israel proofs, also blocked, have additional gates beyond the Lorentzian-metric availability (geodesic existence, focal-point existence, σ -model rigidity); those are phase-level commitments. Within the SK-EFT-Hawking programme they sit alongside the Phase 6g.7 cosmic-censorship backlog.

ACKNOWLEDGMENTS

We thank the Mathlib community, particularly Patrick Massot, Michael Rothgang, and Heather Macbeth for the covariant-derivative library; Sébastien Gouëzel for the Riemannian-bundle infrastructure that the Lorentzian companion mirrors; and the SK-EFT-Hawking project’s preemptive-strengthening discipline for catching the `zeroChristoffel.isTorsionFree` P3-trivial cut at the post-wave review.

-
- [1] P. Massot, M. Rothgang, H. Macbeth. Mathlib4 covariant derivatives library. *Mathlib4 commit 8850ed93*, April 2026. `Mathlib.Geometry.Manifold.VectorBundle.CovariantDerivatives.Basic.Torsion1`.
- [2] S. Gouëzel. Riemannian vector bundles in Mathlib4. *Mathlib4 commit 8850ed93*, March 2026. `Mathlib.Topology.VectorBundle.Riemannian`; `Mathlib.Geometry.Manifold.VectorBundle.Riemannian`.
- [3] J. G. Roehm. Phase 6f waves 1–6 module roster: classical-GR algebraic backbone in Lean. *SK-EFT-Hawking project, in preparation*, inventory snapshot 2026-04-29.
- [4] J. G. Roehm. Phase 6g waves 1–6: causal structure, singularity hypothesis bundles, area theorem, no-hair theorem in Lean. *SK-EFT-Hawking project, in preparation*, inventory snapshot 2026-04-29.
- [5] J. G. Roehm. Wave-execution pipeline and preemptive-strengthening discipline for theorem-driven theoretical-physics workflows. *SK-EFT-Hawking project, in preparation*, `docs/WAVE_EXECUTION_PIPELINE.md`.
- [6] R. M. Wald. *General Relativity*. University of Chicago Press, 1984, §3.1.1.
- [7] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry, Vol. I*. Wiley-Interscience, 1963. Theorem IV.2.2 (Levi-Civita uniqueness).
- [8] S. Carroll. *Spacetime and Geometry*. Addison-Wesley, 2004. §3.2 (Koszul formula).
- [9] W. Klingenberg. *Riemannian Geometry*, 2nd ed. de Gruyter Studies in Mathematics, vol. 1, 1995. §1.3 (curvature operator on a vector bundle).
- [10] S. Gouëzel. Lie brackets of vector fields on manifolds in Mathlib4. *Mathlib4 commit 8850ed93*, 2024–2026. `Mathlib.Geometry.Manifold.VectorField.LieBracket (mlieBracket, mlieBracket_self, mlieBracket_swap_apply, mlieBracket_zero_left, leibniz_identity_mlieBracket)`.