Two evaporation regimes for emergent-gravity black holes: a machine-checked partition at the induced-gravity cutoff

John G. Roehm[1]

[1] *NetRxn Foundation*

(Dated: November 6, 2026)

The partition model describes evaporation on a substrate whose Newton constant is induced by fermion loops below an ultraviolet cutoff. It compares local heating above a critical mass with nozzle cooling below it. That critical mass is the one whose emergent gravitational length reaches the cutoff length. We formalize the cutoff identity and derive the opposite local temperature slopes in Lean 4 from a decreasing Schwarzschild mass profile and the leading-order nozzle solution.

INTRODUCTION

A Schwarzschild black hole has negative heat capacity: it grows hotter as it radiates. Its temperature $T_{\rm H} = \hbar/(8\pi M)$ rises without bound as the mass falls, and a hole of initial mass M_0 disappears after a time of order M_0^3 [1]. A sonic horizon in a flowing fluid can initially cool. Balbinot, Fagnocchi and Fabbri solved the linearized backreaction near the horizon and found that the horizon migrates, the supersonic region shrinks, and the emission temperature initially falls [2]. They extrapolate this result to $t \sim T_{\rm H}^{-3}$ and compare the proposed zero-temperature limit to a near-extremal Reissner–Nordström hole. We use *extremality* for the zero-temperature limit of a cooling profile.

Those are opposite signs of the same derivative. Each is derived on its own background, and on a fixed background there is nothing for a criterion to decide. The question becomes sharp on a substrate where the gravitational coupling is itself induced by the matter it couples to. One system then carries a geometric description at large mass and a fluid description at small mass, and something has to separate them.

This Letter formalizes a model separation. For a substrate whose Newton constant is induced by N_f Dirac species below a cutoff $\Lambda_{\rm UV}$, we assign the two local trajectory models using the critical mass

$$M_{\rm c} = \frac{N_f\,\Lambda_{\rm UV}}{12\pi\,\alpha_{\rm ADW}}, \tag{1}$$

where $\alpha_{\rm ADW}$ measures the departure of the induced coupling from its bare one-loop value. The defined scale is fixed once the parameters are supplied: Eq. (1) makes its emergent gravitational length equal to the inverse cutoff. Physical branch selection remains to be derived.

The formal development comprises 16 theorems across 28 declarations, in Lean 4.32.0 against Mathlib revision 81a5d257. The kernel accepts them from `propext`, `Classical.choice` and `Quot.sound` alone, with no substrate-specific axiom and no incomplete proof. Which hypotheses the partition rests on is stated where each is used. The section titled "What would refute this" states the falsifiers and what they test.

THE CRITICAL MASS

Sakharov observed that the one-loop effective action of matter fields in a curved background already contains an Einstein–Hilbert term, so Newton's constant need not be fundamental [3, 4]. For N_f Dirac species regulated by a hard cutoff the induced coupling is $G_N^{\rm Sak} = 12\pi/(N_f\Lambda_{\rm UV}^2)$. We write $G_N^{\rm emerg} = \alpha_{\rm ADW}\,G_N^{\rm Sak}$ for a substrate whose gravitational response departs from that value, with $\alpha_{\rm ADW} = 1$ the bare induced limit. The numerical coefficient is regulator dependent; what survives regulator choice is that the induced coupling is finite and of definite sign [5].

Multiplying Eq. (1) by the emergent coupling $G_N^{\rm emerg} = \alpha_{\rm ADW}G_N^{\rm Sak}$ cancels the substrate factor and gives

$$G_N^{\rm emerg}\,M_{\rm c} \;=\; \Lambda_{\rm UV}^{-1}. \tag{2}$$

$M_{\rm c}$ is the mass whose emergent gravitational length is the shortest length the substrate resolves. A hole heavier than $M_{\rm c}$ has a horizon large compared with the cutoff and admits a geometric description. A hole lighter than $M_{\rm c}$ would need a horizon smaller than the substrate's own resolution, and what responds is the fluid rather than the geometry. Read this way, Eq. (1) is the mass at which the two descriptions exchange roles rather than a fitted scale. Equation (2) is the statement of `G_N_emerg_eval_mul_M_c_eq_inv_cutoff`. The bare product is instead $G_N^{\rm Sak}M_{\rm c} = 1/(\alpha_{\rm ADW}\Lambda_{\rm UV})$; the bare and emergent products agree exactly when $\alpha_{\rm ADW} = 1$. The construction still makes a physical choice: no derivation on this substrate pins the crossover at $G_N^{\rm emerg}M = \Lambda_{\rm UV}^{-1}$ rather than at an order-unity multiple of it, so the coefficient in Eq. (1) is an ansatz even though its parametric form is not.

A hole is described by its mass, its horizon area and its surface gravity κ, and a substrate by $(\alpha_{\rm ADW}, \Lambda_{\rm UV}, N_f)$, each strictly positive; $M_{\rm c}$ is then positive wherever it is defined (`M_c`, `M_c_pos`). Comparing M against $M_{\rm c}$ sorts a hole into one of three regimes, and the sorting is decidable and reversible: each regime label holds if and only if the corresponding mass inequality does (`classify`, `classify_Schwarzschild_iff`, `classify_ADWExtremality_iff`,

`classify_Boundary_iff`). A regime label and a mass comparison therefore carry the same information, and neither can drift from the other.

ONE SUBSTRATE, BOTH BRANCHES

The strongest reason to want a criterion rather than an analogy is that a single material realizes both signs.

A moving domain wall in a thin film of ^{3}He-A obeys $T_{\rm H}(v) = T_{\rm H}^{(0)}(1 - v^2/c_\perp^2)$, with $T_{\rm H}^{(0)} = \hbar c_\perp/(2\pi d)$ fixed by the wall thickness d [7]. The temperature falls with the wall speed v and stays finite as $v \to 0$. Radiation is fed by the kinetic energy of the wall and slows it, so $T_{\rm H}$ rises in time and the end temperature does not vanish. The thin film sits on the heating branch, and Balbinot *et al.* draw exactly that contrast, noting that such a film appears to show a non-vanishing end temperature of the evaporation process [2].

A moving soliton in bulk ^{3}He-A sits on the other branch. Its horizon exists only for $v > c_\perp$, and its surface gravity carries a factor $(1 - c_\perp^2/v^2)^{1/2}$ that vanishes as backreaction slows the soliton toward $c_\perp$; Jacobson and Volovik accordingly describe that system as cooling toward an extremal configuration [6]. Two textures in the same superfluid, evaporating in opposite directions, is what a criterion has to resolve.

THE TWO BRANCHES AND THE CRITERION

Above $M_{\rm c}$ the profile is the textbook one, $T_{\rm H} = 1/(8\pi M)$ in natural units. It is positive and strictly decreasing in M (`T_H_schwarzschild_pos`, `T_H_schwarzschild_strict_decreasing`), so evaporation, which drives $dM/dt < 0$, raises the temperature and the lifetime is finite [1].

The nozzle calculation motivates the cooling branch. In the notation of Balbinot *et al.*, c is the sound speed, $c^2\kappa$ is the unperturbed sonic surface gravity, A_0 is the nozzle throat's cross-sectional area, and ϵ is the dimensionless backreaction parameter. They obtain to leading order

$$T_{\rm H}(t) = \frac{\hbar c}{2\pi}\,\kappa\left[1 - \frac{563}{720\pi}\,\epsilon\,\kappa^3\,c\,A_0\,t\right], \qquad (3)$$

for $c\kappa t \ll 1$ at constant sound speed [2]. The initial slope is strictly negative for positive c, κ, A_0, and ϵ. A separate illustrative profile is the exponential $T_{\rm H}(t) = T_{\rm H}^{(0)} e^{-t/\tau_{\rm cool}}$, positive and strictly decreasing for positive $T_{\rm H}^{(0)}$ and $\tau_{\rm cool}$, and equal to $T_{\rm H}^{(0)}$ at $t = 0$ (`T_H_acoustic_evolution` and its accompanying positivity, monotonicity and initial-value lemmas).

With the same initial temperature, the exponential and Eq. (3) share the sign of the initial slope. The exponential approaches zero as $t \to \infty$ by construction.




FIG. 1. (a) Opposite temperature trends with separate normalized clocks: $t/t_{\rm evap}$ for representative Schwarzschild evaporation (amber, dashed), and $t/\tau_{\rm cool}$ for illustrative exponential cooling (blue, solid). The exponential approaches zero asymptotically and is distinct from the local affine nozzle criterion. (b) The emergent gravitational length $G_N^{\rm emerg}M$ and cutoff length $\Lambda_{\rm UV}^{-1}$, in cutoff-length units, coincide at $M = M_{\rm c}$ [Eq. (2)]. Blue and amber shading indicate the branches below and above this model boundary.

The nozzle calculation supplies the initial cooling sign. The authors propose extending the calculation to condensates, where $c \propto \rho^{1/2}$ and the cooling sign requires a separate derivation.

The profile criterion derives the signs from separate local models. Above $M_{\rm c}$, the chain rule gives $dT_{\rm H}/dt = -M'(t)/(8\pi M(t)^2) > 0$ for positive mass and $M'(t) < 0$. Below $M_{\rm c}$, write Eq. (3) as $T_{\rm H}(t) = T_0(1-\lambda t)$. Its slope is $-T_0\lambda < 0$ for positive nozzle parameters. The criterion requires $t \geq 0$, $c\kappa t < \eta$ for a supplied short-time budget $0 < \eta < 1$, and $\lambda t < 1$ for positive temperature. At $M = M_{\rm c}$ it assigns neither strict sign. Matching the two descriptions across the cutoff requires a separate construction. The corresponding Lean statement is `regime_partition_profile_criterion`. For an actual trajectory $T_0(1 - \lambda t) + R(t)$, strict cooling also requires $R'(t) < T_0\lambda$; no quantitative remainder control is proved here.

Two of the four laws of black-hole mechanics enter

the development, and neither in full. The first-law slot fixes the positivity $0 < G_N^{\mathrm{emerg}}$ that the coupling $1/(8\pi G_N^{\mathrm{emerg}})$ requires, together with the response identity above; the differential first law itself is not formalized. The second-law slot records the Glorioso–Liu conclusion, that the entropy-current divergence is nonnegative, as a hypothesis rather than as a consequence, because deriving it needs the Schwinger–Keldysh effective action [11–13]. The emergent and acoustic branches still lack a general zeroth-law theorem and an affine-time third law. The conditional cooling-slot theorem states that a hole classified into the cooling branch and satisfying the supplied mechanics slots has a strictly negative evaporation rate (`FourLaws_ADWExtremality`, `four_laws_consistent_with_acoustic_regime`). Physical first-law identification, acoustic energy and entropy balances, and microscopic action, exterior and mass matching remain open. The cutoff identities and local slopes do not supply these constructions.

WHAT WOULD REFUTE THIS

A construction assembled from hypotheses is worth stating only if the hypotheses can fail, so each load-bearing component is paired with a theorem that derives a contradiction from its denial.

A single hole above M_{c} observed to cool as it evaporates refutes the construction, with no appeal to how the cooling was modelled: the partition hypotheses, a heating-branch classification and a non-positive rate are together contradictory (`falsifier_schwarzschild_heating`).

On the cooling branch the refutable content is a comparison of two measurements rather than a functional form. Any alternative history that agrees with the profile at the start of an interval, and fails to decrease across it, must disagree at the end (`falsifier_acoustic_decay_form`).

The third law enters through the time $\tau(T_\star)$ needed to reach $T_{\mathrm{H}} = T_\star$. Its strong form requires $\tau(T_\star)$ to exceed any prescribed bound for all small enough $T_\star$ [9]. Kehle and Unger construct regular collapse data for the Einstein–Maxwell-charged scalar field system that reach an exactly extremal horizon at finite advanced time, which bounds τ uniformly on such a tail [10]. The two are contradictory (`falsifier_third_law_form`), so a substrate must choose and the third-law slot cannot be filled vacuously. The theorem selects neither alternative.

The substrate response is constrained from both sides. It is strictly positive whenever $\alpha_{\mathrm{ADW}} > 1$, and nonzero exactly when $\alpha_{\mathrm{ADW}} \neq 1$ (`falsifier_alpha_ADW_dependence`, `delta_ADW_nonzero_iff_alpha_ADW_ne_one`). A substrate with $\alpha_{\mathrm{ADW}} > 1$ and a non-positive response is excluded, and so is one measured away from the bare induced limit with a vanishing response.

DISCUSSION

The result has two limits worth stating. No closed-form $T_{\mathrm{H}}(M)$ interpolating the regimes is derived: $M = M_{\mathrm{c}}$ is the value at which the classifier changes branch, not a demonstrated smooth crossover, and no single analytic profile is shown to join the branches there. The entropy-production statement is imposed within each regime separately, so nothing here matches the entropy current across the boundary.

The partition is also independent of whether the substrate induces gravity, and the two questions have different answers on the same examples. A film of ^{3}He-A satisfies Sakharov's conditions, while the acoustic condensate of Finazzi–Liberati–Sindoni fails universal coupling [8]: its effective metric is defined only for the phonon sector below the healing length, and the depleted atoms occupying excited single-particle states carry no coupling to that metric. That asymmetry is itself formalized, and a companion paper on emergent gravity through black-hole thermodynamics carries the classification in full [17] (`horizonCrooks_substrate_partition`, `volovikJannes_vs_flsBEC_depletion_asymmetry`).
The cooling criterion uses the sign of dT_{H}/dt on a given background independently of how the dynamics of that background arise. The two questions meet only at the induced-gravity layer, where the Clausius relation $\delta Q = T\,dS$ on every local Rindler horizon returns the Einstein equation as an equation of state [16].

The entropy-production hypothesis is the natural target for replacement by a derivation. Kirklin proves the generalized second law in perturbative gravity to all orders beyond the semiclassical limit and without a UV cutoff, at the cost of a free-energy term for the degrees of freedom that locate the later horizon cut [15], and Faulkner and Speranza obtain the same law for arbitrary cuts of a Killing horizon from crossed-product algebras [14]. Formalizing either would turn the recorded hypothesis into a consequence. What settles the partition, though, is measurement: a sonic horizon whose emission temperature is observed to fall while it radiates lands in the cooling branch, and the sign of dT_{H}/dt is accessible in a fluid experiment or a lattice simulation long before any end state is reached.

[1] S. W. Hawking, "Particle creation by black holes," Commun. Math. Phys. **43**, 199 (1975); DOI: 10.1007/BF02345020.

[2] R. Balbinot, S. Fagnocchi, A. Fabbri, "Quantum Effects in Acoustic Black Holes: the Backreaction," Phys.

Rev. D **71**, 064019 (2005); arXiv:gr-qc/0405098; DOI: 10.1103/PhysRevD.71.064019.

[3] A. D. Sakharov, "Vacuum quantum fluctuations in curved space and the theory of gravitation," Sov. Phys. Dokl. **12**, 1040 (1968).

[4] S. L. Adler, "Einstein gravity as a symmetry-breaking effect in quantum field theory," Rev. Mod. Phys. **54**, 729 (1982); DOI: 10.1103/RevModPhys.54.729.

[5] M. Visser, "Sakharov's induced gravity: a modern perspective," Mod. Phys. Lett. A **17**, 977 (2002); arXiv:gr-qc/0204062; DOI: 10.1142/S0217732302007289.

[6] T. A. Jacobson and G. E. Volovik, "Event horizons and ergoregions in ^{3}He," Phys. Rev. D **58**, 064021 (1998); arXiv:cond-mat/9801308; DOI: 10.1103/PhysRevD.58.064021.

[7] T. A. Jacobson and T. Koike, "Black hole and baby universe in a thin film of ^{3}He-A," in *Artificial Black Holes*, edited by M. Novello, M. Visser, G. Volovik (World Scientific, 2002); arXiv:cond-mat/0205174.

[8] S. Finazzi, S. Liberati, L. Sindoni, "Cosmological constant: a lesson from Bose–Einstein condensates," Phys. Rev. Lett. **108**, 071101 (2012); arXiv:1103.4841; DOI: 10.1103/PhysRevLett.108.071101.

[9] W. Israel, "Third law of black-hole dynamics: a formulation and proof," Phys. Rev. Lett. **57**, 397 (1986); DOI: 10.1103/PhysRevLett.57.397.

[10] C. Kehle and R. Unger, "Gravitational collapse to extremal black holes and the third law of black hole thermodynamics," J. Eur. Math. Soc. (2025); arXiv:2211.15742; DOI: 10.4171/JEMS/1591.

[11] P. Glorioso and H. Liu, "The second law of thermodynamics from symmetry and unitarity," (2016); arXiv:1612.07705.

[12] M. Crossley, P. Glorioso, H. Liu, "Effective field theory of dissipative fluids," JHEP **09**, 095 (2017); arXiv:1511.03646; DOI: 10.1007/JHEP09(2017)095.

[13] P. Glorioso, M. Crossley, H. Liu, "Effective field theory for dissipative fluids (II): classical limit, dynamical KMS symmetry and entropy current," JHEP **09**, 096 (2017); arXiv:1701.07817; DOI: 10.1007/JHEP09(2017)096.

[14] T. Faulkner and A. J. Speranza, "Gravitational algebras and the generalized second law," JHEP **11**, 099 (2024); arXiv:2405.00847; DOI: 10.1007/JHEP11(2024)099.

[15] J. Kirklin, "Generalised second law beyond the semiclassical regime," JHEP **07**, 192 (2025); arXiv:2412.01903; DOI: 10.1007/JHEP07(2025)192.

[16] T. Jacobson, "Thermodynamics of spacetime: the Einstein equation of state," Phys. Rev. Lett. **75**, 1260 (1995); arXiv:gr-qc/9504004; DOI: 10.1103/PhysRevLett.75.1260.

[17] J. G. Roehm, "Emergent gravity through black-hole thermodynamics: linearized field equations and the four laws by regime," in preparation (2026).