

# Higher-curvature structure from the Dirac heat kernel: microscopic predictions and observational ceilings

John G. Roehm

Independent Researcher; *NetRxn137@gmail.com*

(Dated: August 9, 2026)

We formalize, in the Lean 4 proof assistant, the basis change from the 3-scalar Wave 1 Dirac  $a_4$  coefficients  $\{R^2, R_{\mu\nu}R^{\mu\nu}, R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}\}$  [1, 2] into Stelle’s canonical  $\{R^2, C^2, \mathcal{G}\}$  basis [3], where  $C^2$  is the Weyl-tensor-squared and  $\mathcal{G}$  the Gauss–Bonnet Euler density (topological in 4D [4]). The closed-form coefficients  $(\alpha, \beta, \gamma)$  at fermion count  $N_f$  are fixed by a  $3 \times 3$  linear-system solve, with sign-definite behaviour:  $\alpha < 0$ ,  $\beta < 0$ ,  $\gamma > 0$  for any  $N_f > 0$ . The substantive cross-bridge to Wave 1 is encoded as a Lean **ring**-closing identity. We then formalize a numerical correctness-push that compares the predicted dimensionless higher-curvature couplings (at SM-relevant counts  $N_f \in [24, 27]$ ) to the four canonical observational ceilings [5, 6]: LIGO/Virgo speed-of-graviton, Hulse–Taylor binary-pulsar period decay, Eöt-Wash short-range gravity, and the Cassini post-Newtonian solar-system test. The Hulse–Taylor pulsar bound is the tightest at  $|\beta| \lesssim 10^{59}$ ; the predicted Wave 1 coefficients sit at  $\mathcal{O}(10^{-3})$ , some 62 orders of magnitude below. The **HigherCurvatureStructure.lean** module ships 11 substantive theorems, zero **sorry**, and zero new axioms; the Python mirror in **src/higher\_curvature/** backs them with 41 cross-checks.

## I. INTRODUCTION

The Sakharov–Adler induced-gravity programme yields, at one loop in the heat-kernel expansion, an Einstein–Hilbert action plus a calculable tower of higher-curvature corrections. In Phase 6e Wave 1 we formalized the leading  $a_2$  coefficient and its exact match to the linearized 6a.1 emergent Newton constant at the Sakharov baseline  $\alpha_{\text{ADW}} = 1$  [7]. Wave 2 takes the next coefficient,  $a_4$ , and unpacks its three-scalar expansion

$$a_4(x) = \frac{N_f}{(4\pi)^2} \left[ c_R R^2 + c_{\text{Ric}} R_{\mu\nu} R^{\mu\nu} + c_{\text{Riem}} R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} \right], \quad (1)$$

with the Christensen–Duff–Vassilevich rational coefficients  $(c_R, c_{\text{Ric}}, c_{\text{Riem}}) = \left( -\frac{5}{12 \cdot 180}, \frac{7}{12 \cdot 180}, -\frac{12}{12 \cdot 180} \right)$ . At the local effective-action level only *two* combinations of these scalars survive in 4D: the trace-free Weyl-squared  $C^2 = R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 2R_{\mu\nu} R^{\mu\nu} + \frac{1}{3}R^2$  and an arbitrary Ricci-trace piece. The Gauss–Bonnet density  $\mathcal{G} = R^2 - 4R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$  integrates to  $32\pi^2 \chi(M)$  on closed 4-manifolds and contributes nothing to the local equations of motion.

The Wave 2 deliverable is twofold: (i) the closed-form basis change to Stelle’s  $\{R^2, C^2, \mathcal{G}\}$  representation with sign-definite coefficients, formalized as a **ring**-closing identity at the density level; (ii) the correctness-push comparison of the resulting dimensionless couplings against current observational ceilings.

## II. CURVATURE BASIS: GAUSS–BONNET AND WEYL-SQUARED

We formalize the two algebraically-independent curvature scalars beyond  $R^2$  as Lean definitions

**gaussBonnet4D** and **weylSquared4D**:

$$\mathcal{G} \equiv R^2 - 4R_{\mu\nu} R^{\mu\nu} + R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}, \quad (2)$$

$$C^2 \equiv R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 2R_{\mu\nu} R^{\mu\nu} + \frac{1}{3}R^2. \quad (3)$$

The Lean theorem **weylSquared4D.eq\_zero\_iff\_conformally\_flat** encodes the conformal-flatness biconditional  $C^2 = 0 \iff R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} = 2R_{\mu\nu} R^{\mu\nu} - \frac{1}{3}R^2$ . Sanity check: de Sitter at Hubble parameter  $H$  has  $(R^2, R_{\mu\nu} R^{\mu\nu}, R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}) = (144H^4, 36H^4, 24H^4)$ , giving  $C^2 = 24 - 72 + 48 = 0$  identically (de Sitter is conformally flat).

The substantive algebraic engine for the Wave 2 basis change is the identity

$$\mathcal{G} - C^2 = \frac{2}{3}R^2 - 2R_{\mu\nu} R^{\mu\nu}, \quad (4)$$

formalized as **gaussBonnet\_minus\_weyl.eq\_R\_minus\_Ricci\_combina** and closed by **ring** after unfolding both Eqs. (2)–(3).

## III. STELLE-BASIS COEFFICIENTS $(\alpha, \beta, \gamma)$

In the canonical  $\{R^2, C^2, \mathcal{G}\}$  basis, the Wave 1  $a_4$  density takes the form

$$a_4(x) = \alpha(N_f) R^2 + \beta(N_f) C^2 + \gamma(N_f) \mathcal{G}, \quad (5)$$

where the coefficients  $(\alpha, \beta, \gamma)$  are determined by the linear system

$$\begin{aligned} \alpha + \frac{\beta}{3} + \gamma &= c_R, \\ -2\beta - 4\gamma &= c_{\text{Ric}}, \\ \beta + \gamma &= c_{\text{Riem}}. \end{aligned} \quad (6)$$

Solving Eq. (6) yields the closed forms

$$\alpha = -\frac{N_f}{324(4\pi)^2}, \quad \beta = -\frac{41N_f}{4320(4\pi)^2}, \quad \gamma = +\frac{17N_f}{4320(4\pi)^2}. \quad (7)$$

Three sign-definite Lean theorems (`a4_alpha_neg`, `a4_beta_neg`, `a4_gamma_pos`) encode the immediate consequence

$$N_f > 0 \implies \alpha(N_f) < 0, \beta(N_f) < 0, \gamma(N_f) > 0. \quad (8)$$

The positive sign of the topological coefficient  $\gamma$  is the fingerprint of the chiral-anomaly contribution to the Euler density:  $\gamma > 0$  couples the Christensen–Duff Dirac sector to the topological sector with a definite orientation.

### Main basis-change identity

The substantive cross-bridge between the original 3-scalar basis and Stelle’s  $\{R^2, C^2, \mathcal{G}\}$  basis is the Lean theorem

```
theorem
a4_density_eq_a4_density_in_RC2GB_basis
(N_f R_sq Ricci_sq Riemann_sq : ) :
a4_density N_f R_sq Ricci_sq Riemann_sq
=
a4_density_in_RC2GB_basis
N_f R_sq Ricci_sq Riemann_sq
```

proved by `ring` after unfolding the Wave 1 coefficients `a4_R_sq_coef`, `a4_Ricci_sq_coef`, `a4_Riemann_sq_coef`, the Wave 2 coefficients `a4_alpha`, `a4_beta`, `a4_gamma`, and the definitions `weylSquared4D`, `gaussBonnet4D`. This is drift-protection per the project’s `feedback.python.lean.refs.drift.md` memo: the proof body literally invokes the Wave 1 names, so any future renaming would break the build immediately.

### IV. OBSERVATIONAL CEILINGS ON DIMENSIONLESS HIGHER-CURVATURE COUPLINGS

In the Stelle truncation  $L = (1/16\pi G_N)(R + \alpha R^2 + \beta C^2)$  the dimensionless coefficients  $(\alpha, \beta)$  are bounded by four canonical observations, translated into action-coefficient ceilings via the EFT framework of [5, 6]:

- LIGO/Virgo speed-of-graviton (GW170817 [8]):  $|\beta| \lesssim 10^{62}$
- Eöt-Wash short-range gravity (50  $\mu\text{m}$ , [9]):  $|\alpha| \lesssim 10^{61}$
- Hulse–Taylor binary-pulsar period decay (PSR B1913+16, [10]):  $|\beta| \lesssim 10^{59}$
- Cassini post-Newtonian solar-system ([11]):  $|\beta| \lesssim 10^{62}$

The Hulse–Taylor pulsar bound is tightest by  $\sim 3$  orders of magnitude. We formalize the four bounds as Lean definitions `hc_bound_LIGO`, `hc_bound_SRG`, `hc_bound_pulsar`, `hc_bound_cassini`.




FIG. 1. Higher-curvature predictions vs observational ceilings. **Left:** Magnitudes of the three Wave 1 Dirac  $a_4$  Christensen–Duff coefficients (per  $(4\pi)^{-2}$ ) at SM-relevant fermion counts  $N_f \in \{24, 27, 100\}$ ; the  $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$  coefficient dominates at every  $N_f$ . **Right:** log-scale comparison to the four canonical observational ceilings; the Hulse–Taylor pulsar bound is tightest at  $|\beta| \lesssim 10^{59}$ , and the predicted  $\mathcal{O}(10^{-3})$  coefficient sits  $\sim 62$  orders of magnitude below. Lean: `higher_curvature_below_pulsar_bound`; `higher_curvature_predictions_strictly_positive`; `H.HigherCurvatureWithinObservationalBounds.pulsar_witness`.

### V. CORRECTNESS-PUSH: PREDICTIONS VS CEILINGS

The structural Wave 2 correctness-push is encoded as the Lean theorem

```
theorem higher_curvature_below_pulsar_bound
{N_f : } (hN_pos : 0 < N_f) (hN_max : N_f 100) :
|a4_R_sq_coef N_f| < hc_bound_pulsar
|a4_Ricci_sq_coef N_f| < hc_bound_pulsar
|a4_Riemann_sq_coef N_f| < hc_bound_pulsar
```

proved by combining (i) the bound  $|1| (4\pi)^2$  from `Real.pi_gt_three` and `nlinarith` (helper lemmas `fourPiSq_gt_one` and `fourPiSqInv_lt_one`); (ii) the trivial numerical estimate  $100 \cdot 12 / (12 \cdot 180) \cdot 1 < 1$ ; and (iii) the chain  $1 < 10^{59}$ . The proof body invokes the Wave 1 `a4_R_sq_coef`, `a4_Ricci_sq_coef`, `a4_Riemann_sq_coef` by name (P6 cross-module bridge integrity).

The 3-conjunct bundle is *not* a P2-redundancy: each conjunct invokes a distinct Wave 1 coefficient and a distinct positivity proof (the  $R^2$  and  $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$  coefficients are negative and need an explicit `abs_neg` step; the  $R_{\mu\nu}R^{\mu\nu}$  coefficient is positive and uses `abs_of_pos` directly).

*Anchoring the “62 orders below” claim.* At the SM-with- $\nu_R$  count  $N_f = 27$  the dominant Riemann<sup>2</sup> coefficient evaluates to

$$|c_{\text{Riem}}(N_f = 27)| = \frac{27}{180(4\pi)^2} \approx 9.499 \times 10^{-4}, \quad (9)$$

and the same form bounds the smaller  $|c_R|, |c_{\text{Ric}}|$  in the same dimensionless natural-units convention. Comparing against the binary-pulsar ceiling  $|\beta| \lesssim 10^{59}$  gives  $10^{59}/(9.499 \times 10^{-4}) \approx 10^{62}$ , i.e. the predicted coupling sits  $\sim 62$  orders of magnitude below the tightest observational bound. The numerical anchors are autogenerated from `papers/paper40_higher_curvature/tables.py` SCALARS spec (regenerate via `scripts/render_paper_tables.py`); the boolean-shape and golden-shape tests live in `tests/test_higher_curvature.py` (`TestObservationalBandFormula`).

A complementary falsifier `higher_curvature_predictions_strictly_positive` ensures that the bounds are *not* passed by trivial vanishing: every Wave 1  $a_4$  coefficient is non-zero for  $N_f > 0$ . This rules out the unhelpful reading “all bounds are passed because all predictions are zero”.

The Lean Prop predicate `H_HigherCurvatureWithinObservationalBounds B` is parameterised by an upper bound  $B$ :

$$H_{\text{HC}}(B) : 0 < B \wedge \forall N_f, 0 < N_f \leq 100,$$

$$|a_4(R^2)| \leq B, |a_4(R_{\mu\nu}R^{\mu\nu})| \leq B, |a_4(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma})| \leq B$$

Unlike a typical tracked-hypothesis Prop, this predicate is *fully discharged* for  $B = 10^{59}$  inside the module: the Lean theorem `H_HigherCurvatureWithinObservationalBounds_pulsar_witness` proves  $H_{\text{HC}}(10^{59})$  by invoking the correctness-push, so consumers receive a closed proof rather than an undischarged hypothesis. The parameterisation over  $B$  is a forward-compatibility hook: future tighter ceilings (e.g. from next-generation pulsar timing) can be discharged at the new  $B$  without changing downstream consumers.

## VI. LEAN MODULE STRUCTURE

The module `lean/SKEFTHawking/HigherCurvatureStructure` ships 11 substantive theorems organised in six sections:

**§1:** `gaussBonnet4D`, `weylSquared4D` definitions plus the conformal-flatness biconditional and the `gaussBonnet_minus_weyl` algebraic identity (2 theorems).

**§2:** Stelle-basis coefficients `a4.alpha`, `a4.beta`, `a4.gamma` (definitions) plus sign-definiteness for  $N_f > 0$  (3 theorems).

**§3:** Density-level definitions `a4.density`, `a4.density_in_RC2GB_basis`, plus the substantive basis-change identity `a4.density_eq_a4.density_in_RC2GB_basis` (1 theorem).

**§4:** Helper bounds `fourPiSq_gt_one`, `fourPiSqInv_lt_one` (2 theorems).

**§5:** Observational-ceiling defs.

**§6:** Correctness-push `higher_curvature_below_pulsar_bound`, falsifier `higher_curvature_predictions_strictly_positive`, plus the tracked-Prop witness `H_HigherCurvatureWithinObservationalBounds_pulsar_witness` (3 theorems).

The module ships zero `sorry` and zero new axioms beyond the Lean 4 kernel’s standard `propext`, `Classical.choice`, `Quot.sound`. The Python mirror in `src/higher_curvature/` provides numerical evaluation of the  $\{R^2, C^2, \mathcal{G}\}$  basis, the Stelle coefficients, and the observational-bound check; a 41-case `pytest` suite cross-checks each Lean theorem against the Wave 1 numerics.

## VII. CONCLUSION

The Wave 1 Christensen–Duff Dirac  $a_4$  coefficient triple is re-expressed in Stelle’s canonical  $\{R^2, C^2, \mathcal{G}\}$  basis with closed-form sign-definite Stelle coefficients  $(\alpha, \beta, \gamma)$ . The basis change is recorded as a Lean ring identity at the density level (cross-bridge to Wave 1). A correctness-push theorem certifies that, in the natural fermion-count window  $0 < N_f \leq 100$ , every predicted dimensionless higher-curvature coefficient sits  $\sim 62$  orders of magnitude below the tightest observational ceiling (the  $\sim 10^{62}$  bound for binary pulsar) — no tension with current observation. The complementary falsifier rules out the trivial reading.

Wave 3 (`NonlinearDiffInvariance.lean`) consumes the  $\{R^2, C^2, \mathcal{G}\}$  basis to formalize order-by-order diff-invariance of the heat-kernel effective action; the structural correctness-push there is whether  $a_4$  contributions to the nonlinear effective Einstein equations are diff-invariant or not — a clean falsifiable test of the ADW emergent-gravity claim.

[1] S. M. Christensen and M. J. Duff, *New gravitational index theorems and supertheorems*, Nucl. Phys. B **154**, 301

(1979).

[2] D. V. Vassilevich, *Heat kernel expansion: user’s manual*,

- Phys. Rep. **388**, 279 (2003) [arXiv:hep-th/0306138].
- [3] K. S. Stelle, *Renormalization of higher-derivative quantum gravity*, Phys. Rev. D **16**, 953 (1977).
  - [4] D. Lovelock, *The Einstein tensor and its generalizations*, J. Math. Phys. **12**, 498 (1971).
  - [5] X. Calmet, S. Capozziello and D. Pryer, *Gravitational effective action at second order in curvature and gravitational waves*, Eur. Phys. J. C **77**, 589 (2017) [arXiv:1708.08253].
  - [6] E. Berti *et al.*, *Testing general relativity with present and future astrophysical observations*, Class. Quantum Grav. **32**, 243001 (2015) [arXiv:1501.07274].
  - [7] J. G. Roehm, *Formal heat-kernel calibration of induced Newton's constant from the ADW Dirac substrate* (in preparation, Phase 6e Wave 1).
  - [8] B. P. Abbott *et al.* (LIGO/Virgo + EM partners), *Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A*, Astrophys. J. Lett. **848**, L13 (2017) [arXiv:1710.05834].
  - [9] D. J. Kapner, E. G. Adelberger, B. R. Heckel, C. D. Hoyle, U. Schmidt and H. E. Swanson, *Tests of the gravitational inverse-square law below the dark-energy length scale*, Phys. Rev. Lett. **98**, 021101 (2007) [arXiv:hep-ph/0611184].
  - [10] J. M. Weisberg and Y. Huang, *Relativistic measurements from timing the binary pulsar PSR B1913+16*, Astrophys. J. **829**, 55 (2016) [arXiv:1606.02744].
  - [11] B. Bertotti, L. Iess and P. Tortora, *A test of general relativity using radio links with the Cassini spacecraft*, Nature **425**, 374 (2003).