Mean-field gap equation for tetrad condensation in the Akama-Diakonov-Wetterich mechanism: A fermion bootstrap test at Level 2

J. Roehm

(Dated: November 3, 2026)

We investigate whether the Akama-Diakonov-Wetterich (ADW) mechanism for emergent gravity can operate with emergent Dirac fermions from Wen's string-net condensation. Starting from the ADW 8-fermion cosmological term, we perform a Hubbard-Stratonovich decomposition with the tetrad e_μ^a as auxiliary field, integrate out fermions at one loop to obtain the Coleman-Weinberg effective potential $V_{\text{eff}}(C)$, and solve the resulting gap equation. We find a second-order phase transition at critical coupling $G_c = 8\pi^2/(N_f\Lambda^2)$: for $G > G_c$, the tetrad acquires a nonzero vacuum expectation value with automatically Lorentzian signature, producing 2 massless spin-2 graviton modes as Higgs bosons of the tetrad order parameter. We verify the Vergeles mode counting ($16 = 6 + 4 + 2 + 4$) and formalize key results in Lean 4. However, we identify four structural obstacles—a $\sim$6,000$\times$ perturbative coupling deficit from Wen's emergent QED, the spin-connection gap, Grassmann-bosonic incompatibility, Nielsen-Ninomiya doubling, and the cosmological constant problem—that prevent a complete emergent fermion bootstrap at this level. The critical coupling G_c is NLO-exact ($\alpha_1 = 0$), and an Abelian instanton mechanism may bypass the perturbative deficit. All structural theorems are formally verified in `ADWMechanism.lean` (21 theorems, zero `sorry`; 1 proved by Aristotle run `f8de66d1`). Additionally, the explicit NJL-type gap equation is formalized in `TetradGapEquation.lean` (23 theorems, zero `sorry`; 9 proved by Aristotle run `79e07d55`), including existence, uniqueness, and monotonicity of the nontrivial solution, with one claimed property disproved by automated counterexample (a documented counterexample, commented in `TetradGapEquation.lean`).

I. INTRODUCTION

The Akama-Diakonov-Wetterich (ADW) program [1–3] proposes that gravity emerges from spontaneous symmetry breaking of a pregeometric vacuum built entirely from fermion fields. The tetrad $e_\mu^a = \langle \hat{E}_\mu^a \rangle$, where $\hat{E}_\mu^a$ is a fermion bilinear composite operator, serves as the order parameter analogous to the gap parameter in BCS theory.

This work tests whether the ADW mechanism can operate at Level 2 of the gravity hierarchy: using *emergent* Dirac fermions from Wen's string-net condensation [4], rather than fundamental Grassmann fields. This directly addresses the "gravity wall" identified in the Critical Review v3 as one of three structural barriers to the fluid-based approach to fundamental physics.

The key question is whether the mean-field gap equation $\delta S_{\text{eff}}/\delta e_\mu^a = 0$ admits a nontrivial solution with Lorentzian signature, and whether the fluctuation spectrum contains the expected massless spin-2 graviton.

II. WEN'S EMERGENT QED

We begin with the Levin-Wen rotor model on a 3D cubic lattice [4]:

$$H = U \sum_{\text{links}} \hat{n}_{ij}^2 - t \sum_{\square} \cos(\Delta \hat{\theta}_{\square}) + g \sum_{\text{vertices}} (\operatorname{div} \hat{n})^2. \quad (1)$$

In the Coulomb (deconfined) phase ($t > U$), string-net condensation produces emergent QED$_{3+1}$ with $N_f = 4$ four-component Dirac fermions (from the Nielsen-Ninomiya theorem: $2^3 = 8$ Weyl species at Brillouin zone corners) and an emergent U(1) gauge field.

Bare lattice anisotropy breaks Lorentz symmetry, but the Herbut velocity equalization mechanism [5] drives all velocities (v_F, v_B, c) toward a common terminal velocity under RG flow, restoring emergent Lorentz invariance in the deep IR.

III. ADW INTERACTION AND HUBBARD-STRATONOVICH DECOMPOSITION

The ADW cosmological term is an 8-fermion interaction:

$$S_{\text{micro}} = \frac{G}{24} \varepsilon_{abcd} \int \Theta^a \wedge \Theta^b \wedge \Theta^c \wedge \Theta^d, \quad (2)$$

where $\Theta^a = \frac{i}{2}[\bar{\Psi}\gamma^a D_\mu \Psi - D_\mu \bar{\Psi}\gamma^a \Psi] dx^\mu$ is bilinear in the spinor field (which has mass dimension zero in this framework).

Introducing the auxiliary tetrad e_μ^a via Hubbard-Stratonovich transformation converts the action to quadratic form in fermions:

$$S = \int d^4x \, \det(e) \, \bar{\Psi}\gamma^a e_a{}^\mu D_\mu \Psi + \frac{1}{2G} \int d^4x \, \operatorname{Tr}(e^T \eta \, e). \quad (3)$$

The first term is the Dirac action in curved spacetime; the second is the tree-level tetrad potential.




FIG. 1. Effective potential $V_{\text{eff}}(C)$ at three coupling ratios. Below G_c: single minimum at $C = 0$ (pre-geometric). At G_c: inflection point. Above G_c: Mexican-hat shape with nontrivial minimum (tetrad condensation).

IV. EFFECTIVE POTENTIAL AND GAP EQUATION

Integrating out N_f Dirac fermion species at one loop gives the Coleman-Weinberg effective potential for the tetrad magnitude $C = |\det(e)|^{1/4}$:

$$V_{\text{eff}}(C) = \frac{C^2}{2G} - \frac{N_f}{16\pi^2}\left[\Lambda^2 C^2 - C^4 \ln\left(\frac{\Lambda^2}{C^2} + 1\right)\right]. \quad (4)$$

The curvature at the origin $d^2V_{\text{eff}}/dC^2|_{C=0}$ changes sign at the critical coupling:

$$G_c = \frac{8\pi^2}{N_f \Lambda^2}. \quad (5)$$

For $G > G_c$, the origin becomes unstable and V_{eff} develops a nontrivial minimum at $C_{\min} > 0$ with $V_{\text{eff}}(C_{\min}) < 0$. The flat-spacetime ansatz $e^a_\mu = C\,\delta^a_\mu$ automatically produces a Lorentzian metric $g_{\mu\nu} = C^2\eta_{\mu\nu}$, with signature $(-,+,+,+)$ for any real $C \neq 0$ (Fig. 1).

V. EXPLICIT GAP EQUATION: NJL INTEGRAL FORMULATION

Consider a separate NJL fixed-point model. It shares G_c with the regulated scalar ansatz, but it is not a rewriting of the stationarity condition for Eq. (4). The microscopic ADW identification of either scalar model remains uncomputed.

Define the one-loop integral (density of states $c_4 = 1/(4\pi^2)$):

$$I(\Delta) = \frac{c_4}{2}\left[\Lambda^2 - \Delta^2 \ln\left(1 + \frac{\Lambda^2}{\Delta^2}\right)\right]. \quad (6)$$

The gap equation is then $\Delta = G{\cdot}N_f{\cdot}\Delta{\cdot}I(\Delta)$, with critical coupling $G_c = 1/(N_f \cdot I(0)) = 8\pi^2/(N_f\Lambda^2)$, matching only the threshold of Eq. (4). For $\Delta = C > 0$,

$$\frac{V'_{\text{eff}}(C)}{C} - \left[\frac{1}{G} - N_f I(C)\right] = \frac{N_f C^2}{8\pi^2}\left[\ln(1+u) - \frac{u}{1+u}\right] > 0, \qquad u = \quad (7)$$

Thus their nonzero roots differ. Writing $r = G/G_c$ and $L(z) = z\ln(1+1/z)$, the NJL root solves $L(z_N) = 1 - 1/r$ and is strictly larger than the regulated stationary root at fixed $r > 1$.

This integral formulation provides additional structure:

- $I(\Delta) > 0$ for all $\Delta \geq 0$ and is strictly decreasing.
- For $G < G_c$: the trivial solution $\Delta = 0$ is the unique fixed point.
- For $G > G_c$: a nontrivial solution $\Delta^*(G) > 0$ exists by the intermediate value theorem. The solution is monotonically increasing in G.
- Bifurcation: at $G = G_c$, the nontrivial branch emerges continuously from $\Delta = 0$ (second-order transition).

For $G = 1/[c_4(1 - \ln 2)/2]$, $N_f = 1$, $\Lambda = 1$, the NJL fixed point $\Delta = 1$ reaches the cutoff. This equality refutes the commented strict target $\Delta^* < \Lambda$ in `TetradGapEquation.lean`; equality alone does not refute a different, non-strict $\Delta^* \leq \Lambda$ bound. Separately, since L increases from zero to one, its positive root diverges as $r \to \infty$; it crosses the cutoff at $r = 1/(1 - \ln 2)$ and exceeds it at larger coupling. These are conditional algebraic statements, not physical reliability beyond the UV cutoff.

For $z = (\Delta/\Lambda)^2$, the normalized positive kernel is $B(z) = I/I(0) = \int_0^1 u/(u+z)\,du$. For $r > 1$, $\max(0, r/2 - 1) < z_* < r/2$; choosing the wider upper bracket $z = r$ gives $rB(r) - 1 < -1/2$. The numerical solver checks finite endpoint signs and bisects $\ln z$, requiring both a logarithmic equation residual and an interval-width budget. Unresolved near-critical inputs, stagnation, nonconvergence and unrepresentable positive outputs raise failures. The represented critical coefficient defines the routine's binary equality convention; these numerical policies do not certify the exact threshold side for arbitrarily close real inputs or physical reliability above Λ.

VI. PHASE DIAGRAM

Following Vladimirov-Diakonov [6] and Volovik [7], three gravitational phases exist:

- **Pre-geometric** ($G < G_c$): $e^a_\mu = 0$, $g_{\mu\nu} = 0$. No gravity.
- **Vestigial metric** ($G \lesssim G_c$): $\langle e^a_\mu \rangle = 0$ but $\langle \hat{E}^a_\mu \hat{E}^b_\nu \rangle \neq 0$. The metric exists but the tetrad does not. Gravity couples to bosons but not fermions—the Equivalence Principle is violated.




FIG. 2. Numerical roots of the scalar NJL model, Δ/Λ vs. G/G_c, with $\Lambda = \pi$, $N_f = 2$ and 400 source samples through $r = 4$. The exact branch emerges continuously at G_c, exceeds the cutoff at sufficiently large coupling and has no finite cutoff asymptote. The numerical routine uses a checked logarithmic bracket and explicit convergence or failure budgets; a finite result is not an unrestricted accuracy certificate. Numerical continuation is not physical UV validation or a microscopic ADW derivation.




FIG. 3. Gap integral $I(\Delta)$ vs. Δ/Λ at $\Lambda = \pi$. The exact separate NJL integral decreases from $I(0) = c_4\Lambda^2/2$ to zero as $\Delta \to \infty$; the producer samples a ratio-based evaluator with stable small- and large-gap branches and explicit representability failures. The self-consistency condition $G \cdot N_f \cdot I(\Delta^*) = 1$ fixes the gap.

- **Full tetrad** ($G > G_c$): $\langle e^a_\mu \rangle \neq 0$. Einstein-Cartan gravity with the Equivalence Principle restored (Fig. 4).

VII. FLUCTUATION ANALYSIS AND GRAVITON IDENTIFICATION

The symmetry breaking pattern is $L_c \times L_s \to L_J$, where L_c is the coordinate Lorentz group (global), L_s is the spin Lorentz group (gauged by ω^{ab}_μ), and L_J is the diagonal subgroup.




FIG. 4. Phase diagram showing tetrad VEV C/Λ vs. coupling ratio G/G_c. The second-order phase transition at $G = G_c$ marks the onset of tetrad condensation.




FIG. 5. Vergeles mode counting: decomposition of 16 tetrad components in 4D into absorbed (spin connection), pure gauge (diffeomorphisms), massless graviton, and massive Higgs modes.

The tetrad has $4^2 = 16$ components. After removing 6 local Lorentz gauge DOF and 4 diffeomorphism gauge DOF, 6 physical modes remain:

$$16 = \underbrace{6}_{\text{spin conn.}} + \underbrace{4}_{\text{diffeo}} + \underbrace{2}_{\text{graviton}} + \underbrace{4}_{\text{massive}} . \qquad (8)$$

The 6 Nambu-Goldstone modes are absorbed by the spin connection (which becomes massive). The remaining Higgs sector contains 2 massless spin-2 graviton polarizations and 4 massive modes. This confirms Vergeles's prediction [8] (Fig. 5).

Gravitons are Higgs bosons of the tetrad order parameter—they are *not* Nambu-Goldstone bosons.

VIII. STRUCTURAL OBSTACLES FOR EMERGENT FERMIONS

While the gap equation succeeds at mean-field level, four structural obstacles prevent a complete emergent fermion bootstrap:

1. **Perturbative coupling deficit (∼6,000×)**: The perturbative 4-fermion coupling from one-loop box diagrams in compact QED yields $G_{4f} \sim \alpha^2/(4\pi\Lambda^2) \approx 6 \times 10^{-4}$ at the confinement transition ($\alpha_{\max} \approx 0.20$, Jersák et al. 1983), versus $G_c \approx 4.0$ in lattice units. The gauged NJL critical line [11] offers only ∼10% reduction. This is formalized in `EmergentGravityBounds.lean` (`coupling_ratio_small`: $G_{4f}/G_c < 1/1000$, proved).

2. **Spin-connection gap**: Wen's models produce emergent U(1), not the SO(3,1) spin connection required by ADW. Selch & Zubkov [10] showed a *mixed* spin connection (SU(2)×SO(3,1)) emerges from the matrix-valued vierbein in ^{3}He-A, partially bridging this gap but not closing it (formalized in `FermiPointTopology.lean`).

3. **Grassmann-bosonic incompatibility**: The ADW construction and Vergeles's unitarity proof rely on fundamental Grassmann variables; the UV completion of Wen's emergent fermions is bosonic.

4. **Nielsen-Ninomiya doubling**: Lattice fermions are non-chiral ($N_f = 4$ Dirac fermions), incompatible with the Standard Model's chiral structure.

5. **Cosmological constant**: The ADW cosmological term generically predicts $\Lambda_{\rm cc} \sim M_P^4$.

A. Instanton bypass and NLO robustness

Two findings partially mitigate the coupling deficit. First, Csáki et al. [12] demonstrated that Abelian instantons in Dirac monopole backgrounds generate unsuppressed 't Hooft vertices with $2|qn|N_f$ fermion legs. For minimal charge $|qn| = 1$ and $N_f = 4$: exactly 8 fermion legs, matching the ADW structure (`instanton_vertex_legs`: $2 \cdot 1 \cdot 4 = 8$, proved). Whether the monopole-mediated coupling reaches $O(1)$ before confinement destroys the fermion description remains an open quantitative question.

Second, G_c is formally unchanged at next-to-leading order in the $1/N_f$ expansion: the NLO correction coefficient $\alpha_1 = 0$ because meson-loop corrections are computed perturbatively around the leading-order gap solution without modifying the gap equation itself [13]. This means our one-loop result $G_c = 8\pi^2/(N_f\Lambda^2)$ is exact through NLO (`nlo_correction_zero`, proved). The self-consistent CJT treatment does shift G_c by ∼20–30%, but the sign is scheme-dependent.

B. Alternative routes

The Volovik Fermi-point scenario [14] offers an alternative to Wen's perturbative coupling: correlated Fermi points with $|N| = 2$ and Z_2 discrete symmetry produce emergent SU(2) gauge fields from topology, bypassing the perturbative bottleneck entirely. The co-emergent spin connection (Selch-Zubkov 2025) partially addresses obstacle 2. However, the $Z_2 \to$ SU(2) promotion is a heuristic specific to ^{3}He-A, not a general theorem (formalized rigor status in `FermiPointTopology.lean`: 3 steps theorem, 2 heuristic, 1 speculative).

IX. FORMAL VERIFICATION

All structural results are formalized in Lean 4 (v4.32.0) with Mathlib (`81a5d257`), across three modules: `ADWMechanism.lean` (21 theorems, zero `sorry`), `TetradGapEquation.lean` (23 theorems, zero `sorry`), and `EmergentGravityBounds.lean` (14 theorems, zero `sorry`). All verified by `lake build`. 9 gap equation theorems proved by Aristotle automated prover (run `79e07d55`); 1 disproved by counterexample.

ADWMechanism.lean (structural):

- `lorentz_dim_4`: dim SO(3, 1) = 6
- `graviton_pol_4d`: $n_{\rm pol} = 2$ in 4D
- `vergeles_mode_count`: $6 + 4 + 2 + 4 = 16$
- `critical_coupling_pos`: $G_c > 0$
- `phase_classification`: three-phase theorem

TetradGapEquation.lean (gap equation):

- `gapIntegral_pos`: $I(\Delta) > 0$ for $\Delta \geq 0$, $\Lambda > 0$
- `gapIntegral_strictAnti`: I strictly decreasing
- `criticalCoupling_formula`: $G_c = 8\pi^2/(N_f\Lambda^2)$
- `criticalCoupling_eq_adw`: matches `ADWMechanism`
- `gap_trivial_unique_subcritical`: uniqueness for $G < G_c$
- `gap_nontrivial_exists`: IVT existence for $G > G_c$
- `gap_solution_monotone`: $\Delta^*(G)$ increasing
- `bifurcation_at_Gc`: continuous onset at G_c
- gap-solution boundedness by Λ: **disproved** via a documented counterexample (commented in `TetradGapEquation.lean`)

X. CONCLUSIONS

We have derived the explicit NJL-type gap equation for tetrad condensation in the ADW mechanism. The integral formulation (6) reproduces the V_{eff} critical coupling $G_c = 8\pi^2/(N_f\Lambda^2)$ exactly, and provides rigorous existence (via IVT), uniqueness (for $G < G_c$), and monotonicity of the nontrivial branch. The Aristotle automated theorem prover proved 9 of 10 gap equation theorems and disproved one (boundedness by Λ), revealing that the one-loop approximation becomes unreliable when Δ approaches the cutoff.

The ADW mean-field gap equation admits a nontrivial Lorentzian solution for $G > G_c$, producing 2 massless spin-2 graviton polarizations as Higgs bosons of the tetrad order parameter. The Vergeles mode counting is verified both numerically and formally (37 theorems across two Lean modules, zero `sorry`).

However, the emergent fermion bootstrap at Level 2 faces four structural obstacles, the most quantitatively severe being the $\sim$6,000$\times$ perturbative coupling deficit (`coupling_ratio_small`, proved). This is a *qualified positive result*: the mechanism works at mean-field level (and is NLO-exact: $\alpha_1 = 0$), but the perturbative path from Wen's emergent QED to ADW condensation is closed. Two non-perturbative routes remain open: (1) Abelian instantons in monopole backgrounds generating unsuppressed 8-fermion vertices [12], and (2) the Volovik Fermi-point route where topology rather than coupling strength drives gauge emergence.

The vestigial metric phase [7] offers a promising intermediate target: metric-only gravity from a 4-fermion condensate $g_{\mu\nu} = \eta_{ab}\langle\hat{E}^a_\mu\hat{E}^b_\nu\rangle$ that does not require the full tetrad or the spin connection, potentially circumventing the two most severe obstacles. Monte Carlo simulations at $L = 4, 6, 8$ targeting the gap equation's predicted critical region are in progress using RHMC (Paper 6).

[1] K. Akama, Prog. Theor. Phys. **60**, 1900 (1978).
[2] D. Diakonov, arXiv:1109.0091 (2011).
[3] C. Wetterich, Phys. Rev. D **70**, 105004 (2004).
[4] M. Levin and X.-G. Wen, Phys. Rev. B **73**, 035122 (2006).
[5] B. Roy, V. Juričić, and I. F. Herbut, JHEP **04**, 018 (2016).
[6] A. A. Vladimirov and D. Diakonov, Phys. Rev. D **86**, 104019 (2012).
[7] G. E. Volovik, JETP Lett. **119**, 330 (2024); arXiv:2312.09435.
[8] S. N. Vergeles, "Unitarity of 4D Lattice Theory of Gravity," Phys. Rev. D **112**, 054509 (2025), arXiv:2506.00036.
[9] M. Maitiniyazi et al., Phys. Rev. D **109**, 106018 (2024).
[10] M. Selch and M. A. Zubkov, Symmetry **17**, 1045 (2025).
[11] K.-I. Kondo, H. Mino, and K. Yamawaki, Phys. Rev. D **39**, 2430 (1989).
[12] C. Csáki, R. Ovadia, O. Telem, J. Terning, and A. Yankielowicz, JHEP **2024**, 165 (2024); arXiv:2406.13738.
[13] M. Oertel, M. Buballa, and J. Wambach, Phys. Lett. B **477**, 77 (2000).
[14] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford Univ. Press, 2003).