

Neutrino Oscillations in Vacuum and Matter ¹

¹ 2015 Summer

Lei Ma

May 15 2015

Notes for neutrino oscillations in vacuum and dense matter.

Contents

<i>Symbols and Definitions</i>	1
<i>Vacuum Oscillations</i>	2
<i>Survival Probability</i>	3
<i>Two Flavor States</i>	4
<i>Density Matrix</i>	6
<i>An Example of Survival Probability</i>	8
<i>Numerical Results for 2 Flavor Oscillations</i>	8
<i>Three Flavor States</i>	10
<i>Ternary Diagram for Neutrino Flavor Oscillation</i>	11
<i>Oscillations in Matter</i>	12
<i>Analytic Solution</i>	13
<i>Constant Electron Number Density</i>	15
<i>Adiabatic Limit</i>	16
<i>First Order Approximation</i>	18
<i>General Discussion for Neutrinos Interacting with Matter</i>	22
<i>Numerical Method and Results</i>	25
<i>Numerical Calculations for Matter Effect in Flavor Basis</i>	28
<i>Ternary Diagrams</i>	28
<i>Decapitated - Solar Density Profile</i>	28

Symbols and Definitions

- $\Delta = \sqrt{2}G_F n(x)$
- $\omega = \frac{\Delta m^2}{2E}$

Vacuum Oscillations

Schrodinger equation is

$$i\partial_t |\Psi\rangle = \mathbf{H} |\Psi\rangle, \quad (1)$$

where for relativistic neutrinos, the energy is²

$$\mathbf{H}^m = \begin{pmatrix} \sqrt{p^2 + m_1^2} & 0 & 0 \\ 0 & \sqrt{p^2 + m_2^2} & 0 \\ 0 & 0 & \sqrt{p^2 + m_3^2} \end{pmatrix},$$

in which the energy terms are simplified using the relativistic condition

$$\sqrt{p^2 + m_i^2} = p \sqrt{1 + \frac{m_i^2}{p^2}} \quad (2)$$

$$\approx p \left(1 + \frac{1}{2} \frac{m_i^2}{p^2}\right). \quad (3)$$

In general the flavor eigenstates are the mixing of the mass eigenstates with a unitary matrix \mathbf{U} , that is

$$|v_\alpha\rangle = U_{\alpha i} |v_i\rangle, \quad (4)$$

where the α s are indices for flavor states while the i s are indices for mass eigenstates.

To find out the equation of motion for flavor states, plugin in the unitary transformation,

$$iU_{\alpha i}\partial_t |v_i\rangle = U_{\alpha i}H_{ij}^m |v_j\rangle. \quad (5)$$

I use index m for representation of Hamiltonian in mass eigenstates. Applying the unitary condition of the transformation,

$$\mathbf{I} = \mathbf{U}^\dagger \mathbf{U}, \quad (6)$$

I get

$$iU_{\alpha i}\partial_t |v_i\rangle = U_{\alpha i}H_{ij}^m U_{j\beta}^\dagger U_{\beta k} |v_k\rangle, \quad (7)$$

which is simplified to

$$i\partial_t |v_\alpha\rangle = H_{\alpha\beta}^f |v_\beta\rangle, \quad (8)$$

since the transformation is time independent.

² They all have the same momentum but different mass. The thing is we assume they have the same velocity since the mass is very small. To have an idea of the velocity difference, I can calculate the distance travelled by another neutrino in the frame of one neutrino.

Assuming the mass of a neutrino is 1eV with energy 10MeV, we will get a speed of $1 - 10^{-14}c$. This $10^{-14}c$ will make a difference about $3\mu\text{m}$ in 1s.

To Be Discussed!

Will decoherence happen due to this? For high energy neutrinos this won't be a problem however for low energy neutrinos this will definitely cause a problem for the wave function approach. Because the different mass eigenstates will decoherent gradually along the path.

The new Hamiltonian in the representations of flavor eigenstates reads

$$H_{\alpha\beta}^f = U_{\alpha i}^\dagger H_{ij}^m U_{j\beta}. \quad (9)$$

Survival Probability

The neutrino states at any time can be written as

$$|\Psi(t)\rangle = X_1 |v_1\rangle e^{-iE_1 t} + X_2 |v_2\rangle e^{-iE_2 t}, \quad (10)$$

where X_1 and X_2 are the initial conditions which are determined using the neutrino initial states.

Survival probability is the square of the projection on an flavor eigenstate,

$$P_\alpha(t) = |\langle v_\alpha | \Psi(t) \rangle|^2. \quad (11)$$

The calculation of this expression requires our knowledge of the relation between mass eigenstates and flavor eigenstates which we have already found out.

Recall that the transformation between flavor and mass states is

$$|v_i\rangle = U_{i\alpha}^{-1} |\nu_\alpha\rangle, \quad (12)$$

which leads to the inner product of mass eigenstates and flavor eigenstates,

$$\langle \nu_\alpha | v_i \rangle = \langle \nu_\alpha | U_{i\beta}^{-1} |\nu_\beta\rangle \quad (13)$$

$$= U_{i\beta}^{-1} \delta_{\alpha\beta} \quad (14)$$

$$U = U_{i\alpha}^{-1}. \quad (15)$$

The survival probability becomes

$$\begin{aligned} P_\alpha(t) &= |\langle \nu_\alpha | X_1 |v_1\rangle e^{-iE_1 t} + X_2 |v_2\rangle e^{-iE_2 t} \rangle|^2 \\ &= |X_1 e^{-iE_1 t} \langle \nu_\alpha | v_1 \rangle + X_2 e^{-iE_2 t} \langle \nu_\alpha | v_2 \rangle|^2 \\ &= |\sum_i X_i e^{-iE_i t} U_{i\alpha}^{-1}|^2 \\ &= \sum_i X_i^* e^{iE_i t} U_{i\alpha}^{\dagger*} \sum_i X_i e^{-iE_i t} U_{i\alpha}^\dagger \\ &= |X_1|^2 U_{1\alpha}^{\dagger*} U_{1\alpha}^\dagger + |X_2|^2 U_{2\alpha}^{\dagger*} U_{2\alpha}^\dagger + X_1^* X_2 U_{1\alpha}^{\dagger*} U_{2\alpha}^\dagger e^{iE_1 t - iE_2 t} + X_2^* X_1 U_{2\alpha}^{\dagger*} U_{1\alpha}^\dagger e^{iE_2 t - iE_1 t} \end{aligned}$$

$U_{i\alpha}^{\dagger*}$ stands for the i th row and the α th column of the matrix $U^{\dagger*}$.

Two Flavor States

For 2 flavor neutrinos the Hamiltonian in the representation of propagation states,

$$\mathbf{H} = \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix} = \begin{pmatrix} p_1 + \frac{1}{2} \frac{m_1^2}{p_1} & 0 \\ 0 & p_2 + \frac{1}{2} \frac{m_1^2}{p_2} \end{pmatrix}.$$

The equation of motion in matrix form is

$$i\partial_t \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} = \begin{pmatrix} p_1 + \frac{1}{2} \frac{m_1^2}{p_1} & 0 \\ 0 & p_2 + \frac{1}{2} \frac{m_1^2}{p_2} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} \quad (16)$$

The flavor eigenstate is a mixing of the propagation eigenstates,

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} \quad (17)$$

Denote the rotation matrix using \mathbf{U} , the transformation can be written as

$$|\nu_\alpha\rangle = \mathbf{U}_{\alpha i} |\nu_i\rangle, \quad (18)$$

where α is for the flavor eigenstates and i is for the mass eigenstates.

The survival probability has been derived in previous section, which is the projection of propagation states onto flavor states.

For arbitrary initial condition,

$$\Psi(t=0) = A |\nu_a\rangle + B |\nu_b\rangle, \quad (19)$$

which can be rewritten into a matrix form,

$$\Psi(t=0) = \begin{pmatrix} A & B \end{pmatrix} \begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} \quad (20)$$

To write down the projection, the relation

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} \quad (21)$$

is needed. BTW, the inverse transformation is the transpose of \mathbf{U} since \mathbf{U} is unitary, thus we have the relation,

$$\begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} = \begin{pmatrix} \cos \theta_v & -\sin \theta_v \\ \sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} \quad (22)$$

Thus in the state can be written as

$$\Psi(t=0) = \begin{pmatrix} A & B \end{pmatrix} \begin{pmatrix} \cos \theta_v & \sin \theta_v \\ -\sin \theta_v & \cos \theta_v \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}. \quad (23)$$

At any t , the state is

$$\Psi(t) = \begin{pmatrix} A \cos \theta_v - B \sin \theta_v & A \sin \theta_v + B \cos \theta_v \end{pmatrix} \begin{pmatrix} \nu_1 e^{-iE_1 t} \\ \nu_2 e^{-iE_2 t} \end{pmatrix} \quad (24)$$

$$= \begin{pmatrix} (A \cos \theta_v - B \sin \theta_v) e^{-iE_1 t} & (A \sin \theta_v + B \cos \theta_v) e^{-iE_2 t} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix} \quad (25)$$

The survival probability which is projection on a flavor state is written as

$$P(\nu_\alpha, t) = |\langle \nu_\alpha | \Psi(t) \rangle|^2. \quad (26)$$

The survival amplitude for ν_a is

$$\begin{aligned} & \langle \nu_a | \Psi(t) \rangle \\ &= \langle \nu_a | \left((A \cos \theta_v - B \sin \theta_v) e^{-iE_1 t} |\nu_1\rangle + (A \sin \theta_v + B \cos \theta_v) e^{-iE_2 t} |\nu_2\rangle \right) \\ &= (\cos \theta_v \langle \nu_1 | + \sin \theta_v \langle \nu_2 |) \left((A \cos \theta_v - B \sin \theta_v) e^{-iE_1 t} |\nu_1\rangle + (A \sin \theta_v + B \cos \theta_v) e^{-iE_2 t} |\nu_2\rangle \right) \end{aligned}$$

This is simple since the transformation matrix is real.

Applying the condition that the propagation eigenstates are orthonormal, the survival probability is

$$\begin{aligned} P(\nu_a, t) &= |\langle \nu_a | \Psi(t) \rangle|^2 \\ &= |\cos \theta_v (A \cos \theta_v - B \sin \theta_v) e^{-iE_1 t} + \sin \theta_v (A \sin \theta_v + B \cos \theta_v) e^{-iE_2 t}|^2 \\ &= |(A \cos^2 \theta_v - B \sin \theta_v \cos \theta_v) e^{-iE_1 t} + (A \sin^2 \theta_v + B \sin \theta_v \cos \theta_v) e^{-iE_2 t}|^2 \end{aligned}$$

In a special limit that $E_1 = E_2 = E$, the probability becomes

$$P(\nu_a, t) = |A|^2 \quad (27)$$

which is the same as initial probability since there is no mixing at all.

There are two kinds of initial conditions.

- The neutrinos are all in ν_a state initially, which means $A = 1, B = 0$. The survival probability simplifies to

$$\begin{aligned} P(\nu_a, t) &= |\cos^2 \theta_v e^{-iE_1 t} + \sin^2 \theta_v e^{-iE_2 t}|^2 \\ &= |\cos^2 \theta_v e^{-i(E_1 - E_2)t} + \sin^2 \theta_v|^2 \end{aligned}$$

As we have already discussed, $E_1 - E_2 = \frac{m_1^2 - m_2^2}{2p}$ assuming the neutrinos have the same momentum.³ Using the notation $\Delta m^2 = m_1^2 - m_2^2$ and the approximation that $E \approx p$, the survival probability can be rewritten as

³ And here is a question.

$$\begin{aligned}
 P(\nu_a, t) &= \cos^4 \theta_v + \sin^4 \theta_v + \cos^2 \theta_v \sin^2 \theta_v \left(e^{-i\Delta m^2 t/E} + e^{i\Delta m^2 t/E} \right) \\
 &= 1 - 2 \cos^2 \theta_v \sin^2 \theta_v + 2 \cos^2 \theta_v \sin^2 \theta_v \cos \left(\frac{\Delta m^2 t}{2E} \right) \\
 &= 1 - 2 \cos^2 \theta_v \sin^2 \theta_v \left(1 - \cos \left(\frac{\Delta m^2 t}{2E} \right) \right) \\
 &= 1 - 4 \cos^2 \theta_v \sin^2 \theta_v \sin^2 \left(\frac{\Delta m^2 t}{4E} \right) \\
 &= 1 - \sin^2(2\theta_v) \sin^2 \left(\frac{\Delta m^2 t}{4E} \right)
 \end{aligned}$$

We always assuming that in the region of interest, all neutrinos are travelling with the same speed, i.e., the speed of light $c = 1$.⁴ Time is related to distance, $L = t$. Survival probability at distance L is

⁴ which is not true obviously

$$P(\nu_a, L) = 1 - \sin^2(2\theta_v) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right) \quad (28)$$

- The neutrinos are all in ν_b state initially. Equivalently, we have $A = 0, B = 1$. Survival probability is

$$\begin{aligned}
 P(\nu_a, t) &= |-\sin \theta_v \cos \theta_v e^{-iE_1 t} + \sin \theta_v \cos \theta_v e^{-iE_2 t}|^2 \\
 &= \sin^2 \theta_v \cos^2 \theta_v |e^{-i(E_1 - E_2)t} - 1|^2 \\
 &= \sin^2 \theta_v \cos^2 \theta_v \left(1 + 1 - e^{-i\Delta m^2 t/2E} - e^{i\Delta m^2 t/2E} \right) \\
 &= 2 \sin^2 \theta_v \cos^2 \theta_v \left(1 - \cos \left(\frac{\Delta m^2 t}{2E} \right) \right) \\
 &= \sin^2(2\theta_v) \sin^2 \left(\frac{\Delta m^2 t}{4E} \right) \\
 &= \sin^2(2\theta_v) \sin^2 \left(\frac{\Delta m^2 L}{4E} \right)
 \end{aligned}$$

Density Matrix

This problem can be solved using density matrix ρ and Von Neumann equation

$$i\partial_t \rho = [H, \rho]. \quad (29)$$

The initial condition for this equation is

$$\begin{aligned}\rho(t=0) &= (A|\nu_a\rangle + B|\nu_b\rangle)(A^*\langle\nu_a| + B^*\langle\nu_b|) \\ &= AA^*|\nu_a\rangle\langle\nu_a| + BB^*|\nu_b\rangle\langle\nu_b| + AB^*|\nu_a\rangle\langle\nu_b| + A^*B|\nu_b\rangle\langle\nu_a|.\end{aligned}$$

To calculate the propagation of the states, we need the Hamiltonian matrix in flavor basis.

This can be done by finding out how the Hamiltonian matrix transforms from one basis to another.

Using propagation basis,

$$i\partial_t|\Psi_p\rangle = H_p|\Psi_p\rangle. \quad (30)$$

The states are $|\Psi\rangle = \mathbf{U}|\Psi_p\rangle$ in flavor basis, which means we could plug in $|\Psi_p\rangle = \mathbf{U}^T|\Psi\rangle$.

$$i\partial_t\mathbf{U}^T|\Psi\rangle = H_p\mathbf{U}^T|\Psi\rangle.$$

Since $\mathbf{U}\mathbf{U}^T = \mathbf{I}$, we have a clean result by multiplying through the equation by \mathbf{U} .

$$i\partial_t|\Psi\rangle = \mathbf{U}H_p\mathbf{U}^T|\Psi\rangle.$$

So we define $H = \mathbf{U}H_p\mathbf{U}^T$ as the Hamiltonian matrix in flavor basis, which is

$$H = \left(p + \frac{m_1^2 + m_2^2}{4p}\right)\mathbf{I} - \frac{1}{4p} \begin{pmatrix} -\Delta m^2 \cos 2\theta & \Delta^2 m \sin 2\theta \\ \Delta m^2 \sin 2\theta & \Delta^2 m \cos 2\theta \end{pmatrix}. \quad (31)$$

The derivation of this is

$$\begin{aligned}\mathbf{H}_\alpha &= \mathbf{U}\hat{H}_j\mathbf{U}^T \\ &= \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \left(p\mathbf{I} + \frac{1}{2p} \begin{pmatrix} m_1^2 & 0 \\ 0 & m_2^2 \end{pmatrix} \right) \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \\ &= p\mathbf{I} + \frac{1}{2p} \begin{pmatrix} \cos^2\theta m_1^2 + \sin^2\theta m_2^2 & -\sin\theta \cos\theta m_1^2 + \sin\theta \cos\theta m_2^2 \\ -\sin\theta \cos\theta m_1^2 + \sin\theta \cos\theta m_2^2 & \sin^2\theta m_1^2 + \cos^2\theta m_2^2 \end{pmatrix} \\ &= p\mathbf{I} + \frac{1}{2p} \begin{pmatrix} m_1^2 - \Delta m^2 \sin^2\theta & -\frac{1}{2}\sin 2\theta \Delta m^2 \\ -\frac{1}{2}\sin 2\theta \Delta m^2 & m_2^2 + \Delta m^2 \sin^2\theta \end{pmatrix} \\ &= p\mathbf{I} + \frac{1}{2p} \left(\frac{1}{2}(m_1^2 + m_2^2)\mathbf{I} - \frac{1}{2} \begin{pmatrix} -\Delta m^2 \cos 2\theta & \Delta^2 m \sin 2\theta \\ \Delta m^2 \sin 2\theta & \Delta^2 m \cos 2\theta \end{pmatrix} \right) \\ &= \left(p + \frac{m_1^2 + m_2^2}{4p} \right) \mathbf{I} - \frac{1}{4p} \begin{pmatrix} -\Delta m^2 \cos 2\theta & \Delta^2 m \sin 2\theta \\ \Delta m^2 \sin 2\theta & \Delta^2 m \cos 2\theta \end{pmatrix}\end{aligned}$$

where $\Delta m^2 = m_1^2 - m_2^2$.

Since identity matrix only shifts the eigenvalues we are only interested in the second term, thus the Hamiltonian we are going to use is

$$H = \frac{\Delta m^2}{4E} \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ -\sin 2\theta & -\cos 2\theta \end{pmatrix}. \quad (32)$$

The equation of motion becomes

$$i\partial_t \begin{pmatrix} u(t) \\ v(t) \end{pmatrix} = \frac{\Delta m^2}{4E} \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ -\sin 2\theta & -\cos 2\theta \end{pmatrix} \begin{pmatrix} u(t) \\ v(t) \end{pmatrix} \quad (33)$$

To solve this we need the eigenvalues and eigenvectors of the Hamiltonian matrix.

An Example of Survival Probability

Suppose the neutrinos are prepared in electron flavor initially, the survival probability of electron flavor neutrinos is calculated using the result I get previously.

Electron neutrinos are the lighter ones, then I have $a = e$ and denote $b = x$.⁵

The survival probability for electron neutrinos is

$$P(\nu_e, L) = 1 - \sin^2(2\theta_v) \sin^2\left(\frac{\Delta m^2 L}{4E}\right).$$

Numerical Results for 2 Flavor Oscillations

To solve a set of first order differential equations, I need the determinant of coefficient matrix. For 2 flavor neutrino oscillations,

$$\partial_t \begin{pmatrix} u(t) \\ v(t) \end{pmatrix} = -i\frac{\Delta m^2}{4E} \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ -\sin 2\theta & -\cos 2\theta \end{pmatrix} \begin{pmatrix} u(t) \\ v(t) \end{pmatrix}.$$

To find the solutions I need the eigenvalues λ I need to find the determinant

$$\begin{aligned} & \det \left(-i\frac{\Delta m^2}{4E} \begin{pmatrix} \cos 2\theta & -\sin 2\theta \\ -\sin 2\theta & -\cos 2\theta \end{pmatrix} - \lambda \mathbf{I} \right) \\ &= \begin{vmatrix} -i\frac{\Delta m^2}{4E} \cos 2\theta - \lambda & i\frac{\Delta m^2}{4E} \sin 2\theta \\ i\frac{\Delta m^2}{4E} \sin 2\theta & i\frac{\Delta m^2}{4E} \cos 2\theta - \lambda \end{vmatrix}. \end{aligned}$$

By defining $\lambda' = \lambda / (-i\Delta m^2/4E)$, the determinant is

$$-\left(\frac{\Delta m^2}{4E}\right)^2 ((\cos 2\theta - \lambda')(-\cos 2\theta - \lambda') - \sin 2\theta \sin 2\theta).$$

⁵ In the small mixing angle limit,

$$\begin{pmatrix} \nu_e \\ \nu_x \end{pmatrix} \rightarrow \begin{pmatrix} 1 & \theta \\ -\theta & 1 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}$$

which is very close to an identity matrix. This implies that electron neutrino is more like mass eigenstate ν_1 . By ν_1 we mean the state with energy $\frac{\delta m^2}{4E}$ in vacuum.

The eigenvalues are the solutions to

$$-\left(\frac{\Delta m^2}{4E}\right)^2 ((\cos 2\theta - \lambda')(-\cos 2\theta - \lambda') - \sin^2 2\theta \sin 2\theta) = 0,$$

whose solution is

$$\lambda' = \pm 1.$$

With the solutions

$$\lambda = \pm i \frac{\Delta m^2}{4E},$$

the eigenvectors can also be solved.

$$\begin{pmatrix} \cos 2\theta - 1 & -\sin 2\theta \\ -\sin 2\theta & -\cos 2\theta - 1 \end{pmatrix} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

gives us $\eta_2 = -\tan \theta \eta_1$, which means the eigenvectors are

$$\begin{pmatrix} 1 \\ -\tan \theta \end{pmatrix}, \begin{pmatrix} 1 \\ \cot \theta \end{pmatrix}. \quad (34)$$

The general solution of the first order differential equations is

$$\begin{pmatrix} 1 \\ -\tan \theta \end{pmatrix} e^{-i\Delta m^2 t/4E} + \begin{pmatrix} 1 \\ \cot \theta \end{pmatrix} e^{i\Delta m^2 t/4E}.$$

Initial condition is

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

and it determines the final solution

$$\begin{aligned} & \cos^2 \theta \begin{pmatrix} 1 \\ -\tan \theta \end{pmatrix} e^{-i\Delta m^2 t/4E} + \sin^2 \theta \begin{pmatrix} 1 \\ \cot \theta \end{pmatrix} e^{i\Delta m^2 t/4E} \\ &= \begin{pmatrix} \cos^2 \theta \\ -\sin \theta \cos \theta \end{pmatrix} e^{-i\Delta m^2 t/4E} + \begin{pmatrix} \sin^2 \theta \\ \sin \theta \cos \theta \end{pmatrix} e^{i\Delta m^2 t/4E} \end{aligned}$$

The survival probability of electron neutrino is

$$\begin{aligned} P &= |\cos^2 \theta e^{-i\Delta m^2 t/4E} + \sin^2 \theta e^{i\Delta m^2 t/4E}|^2 \\ &= |\cos^2 \theta e^{-i\Delta m^2 t/2E} + \sin^2 \theta|^2, \end{aligned}$$

which gets back to the result we had using the previous method.

This problem can also be solved using numerical methods. Here is a comparison between this analytical result and a numerical result.

Three Flavor States

For three flavor neutrinos, the oscillations matrix is 3 by 3 which is called the PMNS matrix.

$$\mathbf{U} = \begin{pmatrix} U_{11} & U_{12} & U_{13} \\ U_{21} & U_{22} & U_{23} \\ U_{31} & U_{32} & U_{33} \end{pmatrix},$$

which acts on the mass eigenstates to give the flavor eigenstates, i.e.,

$$|\nu_\alpha\rangle = \mathbf{U} |\nu_i\rangle.$$

In general, a rotation for 3D with a CP violation phase δ is

$$\mathbf{U} = \begin{pmatrix} \cos(\theta_{12})\cos(\theta_{13}) & \cos(\theta_{13})\sin(\theta_{12}) & e^{-i\delta_{CP}}\sin(\theta_{13}) \\ -\cos(\theta_{23})\sin(\theta_{12}) - e^{i\delta_{CP}}\cos(\theta_{12})\sin(\theta_{13})\sin(\theta_{23}) & \cos(\theta_{12})\cos(\theta_{23}) - e^{i\delta_{CP}}\sin(\theta_{12})\sin(\theta_{13})\sin(\theta_{23}) & \cos(\theta_{13})\sin(\theta_{23}) \\ \sin(\theta_{12})\sin(\theta_{23}) - e^{i\delta_{CP}}\cos(\theta_{12})\cos(\theta_{23})\sin(\theta_{13}) & -e^{i\delta_{CP}}\cos(\theta_{23})\sin(\theta_{12})\sin(\theta_{13}) - \cos(\theta_{12})\sin(\theta_{23}) & \cos(\theta_{13})\cos(\theta_{23}) \end{pmatrix}.$$

However, with the CP violation phase δ this matrix is not unitary thus I'll choose $\delta = 0$ for simplicity.⁶

Now the matrix becomes,

$$\mathbf{U} = \begin{pmatrix} \cos(\theta_{12})\cos(\theta_{13}) & \cos(\theta_{13})\sin(\theta_{12}) & \sin(\theta_{13}) \\ -\cos(\theta_{23})\sin(\theta_{12}) - \cos(\theta_{12})\sin(\theta_{13})\sin(\theta_{23}) & \cos(\theta_{12})\cos(\theta_{23}) - \sin(\theta_{12})\sin(\theta_{13})\sin(\theta_{23}) & \cos(\theta_{13})\sin(\theta_{23}) \\ \sin(\theta_{12})\sin(\theta_{23}) - \cos(\theta_{12})\cos(\theta_{23})\sin(\theta_{13}) & -\cos(\theta_{23})\sin(\theta_{12})\sin(\theta_{13}) - \cos(\theta_{12})\sin(\theta_{23}) & \cos(\theta_{13})\cos(\theta_{23}) \end{pmatrix}.$$

The survival probability is given by the same derivation as the 2 flavor example. First of all we need to find the Hamiltonian in flavor basis from the propagation Hamiltonian, which is

$$\mathbf{H} = \mathbf{U}\mathbf{H}_p\mathbf{U}^{-1}$$

$$= \begin{pmatrix} \frac{1}{4}(\cos^2(\theta_{12})m_1^2 + \sin^2(\theta_{12})m_2^2)\cos^2(\theta_{13}) + \sin^2(\theta_{13})m_3^2 & \frac{1}{4}(\cos(\theta_{12})\cos(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \sin(2\theta_{12})\sin(\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) & \frac{1}{4}(\cos(\theta_{12})\sin(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \cos(\theta_{12})\sin(2\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) \\ \frac{1}{4}(\cos(\theta_{12})\cos(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \sin(2\theta_{12})\sin(\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) & \frac{1}{4}(\cos(\theta_{12})\cos(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \sin(2\theta_{12})\sin(\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) & \frac{1}{4}(\cos(\theta_{12})\sin(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \cos(\theta_{12})\sin(2\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) \\ \frac{1}{4}(\cos(\theta_{12})\sin(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \sin(2\theta_{12})\sin(\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) & \frac{1}{4}(\cos(\theta_{12})\sin(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \sin(2\theta_{12})\sin(\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) & \frac{1}{4}(\cos(\theta_{12})\cos(2\theta_{12})\sin(2\theta_{13})(m_1^2 - m_2^2) + \cos(\theta_{12})\sin(2\theta_{13})(-\cos^2(\theta_{12})m_1^2 - \sin^2(\theta_{12})m_2^2 + m_3^2)) \end{pmatrix}$$

where

$$\mathbf{H}_p = \begin{pmatrix} m_1^2 & 0 & 0 \\ 0 & m_2^2 & 0 \\ 0 & 0 & m_3^2 \end{pmatrix}.$$

The next step is to define




Figure 1: They overlap on all the range completely.

⁶ The effect of the phase can also be studied easily with the help of Mathematica.

$$\Delta m_{12}^2 = m_1^2 - m_2^2$$

$$\Delta m_{23}^2 = m_2^2 - m_3^2$$

so that they simplify the Hermitian Hamiltonian to

$$\mathbf{H} = \begin{pmatrix} H_{11} & H_{12} & H_{13} \\ H_{21} & H_{22} & H_{23} \\ H_{31} & H_{32} & H_{33} \end{pmatrix},$$

where ⁷

$$H_{11} = \cos^2 \theta_{13}(m_2^2 - \sin^2 \theta \Delta m_{12}^2) + \sin^2 \theta_{13} m_3^2$$

⁷ Haven't been copied from Mathematica code.




Figure 2: The overall shape is the same however they differ on small scales.

Ternary Diagram for Neutrino Flavor Oscillation

Since the probability for differential flavors of neutrinos are represented in barycentric coordinates⁸, a ternary plot would be nice to representation the oscillations. An example is shown in figure 3.

⁸ They sum up to 1.




Figure 3: Ternary diagram for neutrino oscillations. The state starts from bottom left, which means that the system has only electron neutrinos. As the neutrino travels, it oscillates in curves. After one period of the beat, it reaches the far end and then oscillates backwards.

Oscillations in Matter

The Hamiltonian should be determined first. We have already derived the Hamiltonian for vacuum oscillation,

$$H_v = \frac{\delta m^2}{2E} \frac{1}{2} \begin{pmatrix} -\cos 2\theta_v & \sin 2\theta_v \\ \sin 2\theta_v & \cos 2\theta_v \end{pmatrix},$$

where we would like to define a new matrix,

$$\mathbf{B} = \frac{1}{2} \begin{pmatrix} -\cos 2\theta_v & \sin 2\theta_v \\ \sin 2\theta_v & \cos 2\theta_v \end{pmatrix},$$



so that the vacuum Hamiltonian can be written as

$$H_v = \frac{\delta m^2}{2E} \mathbf{B}.$$

The effect of matter, adds an extra term to this vacuum Hamiltonian which makes the electron population weighs more,

$$H_m = \sqrt{2} G_F n_e L.$$

Here we have ⁹



⁹ In principle we could shift the Hamiltonian using an identity matrix $\alpha \mathbf{I}$ without change the eigenvectors. We will use an term $\frac{1}{2} \alpha \mathbf{e}_3$ instead in the following sections.

$$L = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Without emphasizing the self-interaction of the neutrinos, the Hamiltonian to be used is

$$H = H_v + H_m. \quad (35)$$

The equation of motion is simply the von Neumann equation

$$i\partial_t \rho = [H, \rho], \quad (36)$$

in which the ∂_t operator is actually ∂_x since we always assume neutrinos travel with speed of light c .

Analytic Solution

This Hamiltonian can be rewritten into a simple form using Pauli matrices,

$$\begin{aligned} \mathbf{H} &= \frac{\Delta m^2}{4E} (-\cos 2\theta_v \sigma_3 + \sin 2\theta_v \sigma_1) + \frac{\Delta}{2} \sigma_3 \\ &= \left(\frac{\Delta}{2} - \frac{\Delta m^2}{4E} \cos 2\theta_v \right) \sigma_3 + \frac{\Delta m^2}{4E} \sin 2\theta_v \sigma_1 \\ &= \left(\frac{\Delta}{2} - \frac{\omega}{2} \cos 2\theta_v \right) \sigma_3 + \frac{\omega}{2} \sin 2\theta_v \sigma_1, \end{aligned}$$

where $\Delta = \sqrt{2}G_F n(x)$ and $n(x)$ is the number density of the electrons.

To solve the equation of motion, this matrix should be diagonalized and its eigenvalues and eigenvectors should be identified. Since we have this Pauli matrices form, this can be done easily.

To see this effect quantitatively, we need to diagonalize this Hamiltonian (Can we actually diagonalize the equation of motion? NO!).

Equivalently, we can rewrite it in the basis of mass eigenstates $\{|\nu_L(x)\rangle, |\nu_H(x)\rangle\}$

¹⁰,

$$\begin{aligned} |\nu_L(x)\rangle &= \cos \theta(x) |\nu_e\rangle - \sin \theta(x) |\nu_\mu\rangle \\ |\nu_H(x)\rangle &= \sin \theta(x) |\nu_e\rangle + \cos \theta(x) |\nu_\mu\rangle. \end{aligned}$$

This new rotation in matrix form is

$$\begin{aligned} \begin{pmatrix} |\nu_L(x)\rangle \\ |\nu_H(x)\rangle \end{pmatrix} &= \begin{pmatrix} \cos \theta(x) & -\sin \theta(x) \\ \sin \theta(x) & \cos \theta(x) \end{pmatrix} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} \\ &= \mathbf{U}_m^{-1} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix}. \end{aligned}$$

¹⁰ This is very different from the vacuum case since this one is local. In principle we can not use this transformation to diagonalize the Hamiltonian because the equation is a differential equation about x .




Figure 4: Due to this rotation, we can find an angle when the electron neutrinos becomes more important in Heavy state than Light state, which is $\theta = \frac{\pi}{4}$. For a mixing that has a larger angle, electron neutrinos actually take part in heavy neutrino state.

To diagonalize it, we need to multiply on both sides the rotation matrix and its inverse, as we have done in the vacuum case,

$$\mathbf{H}_{md} = \mathbf{U}_m^{-1} \mathbf{H} \mathbf{U}_m.$$

The second step is to set the off diagonal elements to zero. By solving the equations we can find the $\sin 2\theta(x)$ and $\cos 2\theta(x)$.¹¹

¹¹ Just remember that $\Delta = \sqrt{2}G_F n(x)$ is a function of position.

$$\begin{aligned} \mathbf{H}_{md} &= \mathbf{U}_m^{-1} (A_1 \sigma_1 + A_3 \sigma_3) \mathbf{U}_m \\ &= \begin{pmatrix} A_3 \cos 2\theta(x) - A_1 \sin 2\theta(x) & A_3 \sin 2\theta(x) + A_1 \cos 2\theta(x) \\ A_3 \sin 2\theta(x) + A_1 \cos 2\theta(x) & -A_3 \cos 2\theta(x) + A_1 \sin 2\theta(x) \end{pmatrix}, \end{aligned}$$

where

$$\begin{aligned} A_3 &= \frac{\Delta}{2} - \frac{\delta^2 m}{4E} \cos 2\theta_v = \frac{\Delta}{2} - \frac{\omega}{2} \cos 2\theta_v \\ A_1 &= \frac{\delta^2 m}{4E} \sin 2\theta_v = \frac{\omega}{2} \sin 2\theta_v. \end{aligned}$$

Set the off-diagonal elements to zero,

$$A_3 \sin 2\theta(x) + A_1 \cos 2\theta(x) = 0$$

So the solutions are

$$\begin{aligned} \sin 2\theta(x) &= \frac{A_1}{\sqrt{A_1^2 + A_3^2}} \\ \cos 2\theta(x) &= \frac{-A_3}{\sqrt{A_1^2 + A_3^2}}. \end{aligned}$$

This diagonalizes the Hamiltonian LOCALLY.¹² It's not possible to diagonalize the Hamiltonian globally if the electron number density is not a constant.

As $\Delta \rightarrow \infty$, we have $A_3 \rightarrow \infty$ and $\sin 2\theta(x)$ vanishes. Thus the neutrino will stay on the defined light and heavy eigenstates if the interaction with matter is much larger than the vacuum energy. On the other hand, $\Delta \rightarrow 0$, everything gets back to the vacuum case.

Now we can obtain an approximate solution by using this approximate diagonalization. This idea is

$$i\partial_x \Psi_m(x) = \text{ExtraMatrixFromLHS} \cdot \mathbf{H}_{md} \Psi_m(x),$$

where the **ExtraMatrixFromLHS** comes from the fact that changing from flavor basis $\Psi(x)$ to heavy-light basis $\Psi_m(x)$ using \mathbf{U}_m

$$i\partial_x (\mathbf{U}_m \Psi_m(x)) = H(\mathbf{U}_m \Psi_m(x))$$

¹² The point is, for equation of motion, we have a differentiation with respect to position x ! So even we diagonalize the Hamiltonian, the equation of motion won't be diagonalized. An extra matrix will occur on the LHS and revert the diagonalization the Hamiltonian on RHS.

only returns

$$i\partial_x \Psi_m(x) = \mathbf{H}_{\mathbf{md}} \Psi_m(x) - i\mathbf{U}_{\mathbf{m}}^{-1}(\partial_x \mathbf{U}_{\mathbf{m}}) \Psi_m(x).$$

Combining the two terms on RHS,

$$i\partial_x \Psi_m(x) = \mathbf{H}_{\mathbf{m}} \Psi_m(x),$$

where

$$\mathbf{H}_{\mathbf{m}} = \mathbf{H}_{\mathbf{md}} - i\mathbf{U}_{\mathbf{m}}^{-1}(\partial_x \mathbf{U}_{\mathbf{m}}).$$

The only part inside $\mathbf{U}_{\mathbf{m}}(x)$ that is space dependent is the number density of the electrons $n(x)$. **Thus we know immediately that the Hamiltonian is diagonalized if the number density is constant.**^{13,14,15}

Constant Electron Number Density

Suppose we have an environment with constant electron number density, the term $-i\mathbf{U}_{\mathbf{m}}^{-1}(\partial_x \mathbf{U}_{\mathbf{m}})$ goes away. All we have is the diagonalized new Hamiltonian $\mathbf{H}_{\mathbf{md}}$ and the eigenvalues are easily obtained which are

$$\begin{aligned} E_1 &= A_3 \cos 2\theta(x) - A_1 \sin 2\theta(x) \\ E_2 &= -A_3 \cos 2\theta(x) + A_1 \sin 2\theta(x). \end{aligned}$$

The final result for these two eigenvalues are

$$\begin{aligned} E_1 &= -\sqrt{A_1^2 + A_3^2} \\ E_2 &= \sqrt{A_1^2 + A_3^2}, \end{aligned}$$

where

$$A_1^2 + A_3^2 = \frac{\Delta^2 + \omega^2}{4} - \frac{\Delta\omega}{2} \cos 2\theta_v.$$

There are two special cases. The first case is $\cos 2\theta_v = 0$ while the other one is $\cos 2\theta_v = 1$ which leads to $\sqrt{A_1^2 + A_3^2} = |\frac{\Delta-\omega}{2}|$ and the two energies becomes $\frac{\Delta-\omega}{2}$ and $\frac{\omega-\Delta}{2}$.¹⁶

As for the survival probability, the result has the same form as the vacuum case, which is

$$P_x(\nu_e, L) = 1 - \sin^2(2\theta_m) \sin^2\left(\frac{\omega L}{2}\right),$$

where $\theta_m = \theta(x)$ is the effective mixing angle which in fact doesn't depend on x if the matter profile is constant.¹⁷

¹³ But even the electrons density is constant, the oscillation is different.

¹⁴ It would be nice to find the expression of $\mathbf{U}_{\mathbf{m}}$ in order to play with the matter effect.

¹⁵ In a two level system, suppose the Hamiltonian is

$$H_2 = H_{d2} + P_2,$$

where

$$\begin{aligned} H_{d2} &= \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix} \\ P_2 &= \begin{pmatrix} 0 & W(r) \\ W^*(r) & 0 \end{pmatrix} \end{aligned}$$

are the diagonalized Hamiltonian and the perturbation.

The diagonalized Hamiltonian will result in a system with two levels which are constant as a function of r , however the perturbation term provides the r dependent term which will cause a r dependent eigen energies.

¹⁶ This means the matter effect in this constant profile situation determines the difference between the eigen energies.

¹⁷ The term $\sin 2\theta_m$ has been derived previously.

$$\sin 2\theta(x) = \frac{\omega \sin 2\theta_v}{\sqrt{\omega^2 + \Delta^2 - 2\omega\Delta \cos 2\theta_v}}.$$

Adiabatic Limit

In some astrophysical environments the electron number density changes very slowly which means the term $\mathbf{U}_m^{-1} \partial_x \mathbf{U}_m$ is much smaller than \mathbf{H}_{md} . By intuition we would expect that this term could be dropped to the lowest order. However to verify that we need to check the Taylor expansion of this gradient term which requires the general form of \mathbf{U}_m .

Before we actually do the expansion, make the quantities dimensionless would greatly help us.

$$\sin 2\theta(x) = \frac{\sin 2\theta_v}{\sqrt{\left(\frac{\Delta}{\omega}\right)^2 + 1 - 2\frac{\Delta}{\omega} \cos 2\theta_v}}$$

$$\cos 2\theta(x) = \frac{\cos 2\theta_v - \frac{\Delta}{\omega}}{\sqrt{\left(\frac{\Delta}{\omega}\right)^2 + 1 - 2\frac{\Delta}{\omega} \cos 2\theta_v}}.$$

Define $\hat{\Delta} = \frac{\Delta}{\omega}$, which represents the matter interaction strength compared to the vacuum oscillation.

$$\sin 2\theta(x) = \frac{\sin 2\theta_v}{\sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}}$$

$$\cos 2\theta(x) = \frac{\cos 2\theta_v - \hat{\Delta}}{\sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}}.$$

$$\sin \theta(x) = \frac{\csc(2\theta_v) \sqrt{\frac{-\hat{\Delta} + \sqrt{\hat{\Delta}^2 - 2\hat{\Delta} \cos(2\theta_v) + 1} + \cos(2\theta_v)}{\sqrt{\hat{\Delta}^2 - 2\hat{\Delta} \cos(2\theta_v) + 1}}} \left(\hat{\Delta} + \sqrt{\hat{\Delta}^2 - 2\hat{\Delta} \cos(2\theta_v) + 1} - \cos(2\theta_v) \right)}{\sqrt{2}}$$

$$\cos \theta(x) = \frac{\sqrt{\frac{-\hat{\Delta} + \sqrt{\hat{\Delta}^2 - 2\hat{\Delta} \cos(2\theta_v) + 1} + \cos(2\theta_v)}{\sqrt{\hat{\Delta}^2 - 2\hat{\Delta} \cos(2\theta_v) + 1}}}}{\sqrt{2}}$$

This is extremely complicated if we take the derivative with respect to x . So we apply Taylor expansion.¹⁸

¹⁸ The i s are there makes sure that $-i\partial_x \mathbf{U}_m(x)$ is real.

$$\begin{aligned} \frac{d \sin \theta(x)}{dx} &= \frac{d \sin \theta(x)}{d \hat{\Delta}} \frac{d \hat{\Delta}}{dx} \\ &= i \frac{d \hat{\Delta}}{dx} \left(-\cos \theta_v \sin^2 \theta_v - \frac{1}{2} ((1 + 5 \cos 2\theta_v) \cos \theta_v \sin^2 \theta_v) \hat{\Delta} + \mathcal{O}(\hat{\Delta}^2) \right) \\ \frac{d \cos \theta(x)}{dx} &= \frac{d \cos \theta(x)}{d \hat{\Delta}} \frac{d \hat{\Delta}}{dx} \\ &= i \frac{d \hat{\Delta}}{dx} \left(\cos^2 \theta_v \sin \theta_v + \frac{1}{2} \cos^2 \theta_v (-1 + 5 \cos 2\theta_v) \sin \theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta}^2) \right) \end{aligned}$$

There is nothing in the coefficients that prohibits us from using the lowest order that $\frac{d\hat{\Delta}/dx}{\text{Energy Differences Between Two Energy Levels}} \ll 1$ since the Taylor series has finite values.¹⁹

Solving the adiabatic limit is just the same as the constant limit but $\theta(x)$ is not constant which ensures changing energies. What would be interesting is the behavior of energies as the density profile is slowing changing.

$$E_1 = -\frac{\omega}{2} \sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}$$

$$E_2 = \frac{\omega}{2} \sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}.$$

When the term $\hat{\Delta}$ is very small $1 - 2\hat{\Delta} \cos 2\theta_v$ will dominate and the whole term decreases. On the other hand as $\hat{\Delta}$ becomes large, $\hat{\Delta}^2$ will dominate and the whole term grows.

The survival probability for the light neutrinos would be

$$P_x(\nu_L, L) = 1 - \sin^2(2\theta(x)) \sin^2\left(\frac{\omega L}{2}\right).$$

The survival probability for electron flavor neutrino is

$$P_x(\nu_e, L) = \frac{1}{2} + \frac{1}{2} \cos 2\theta(x_0) \cos 2\theta_v,$$

if the neutrinos are produced in dense region and the detection happens in vacuum.

Before we move on to higher order corrections, it would be nice to understand this phenomenon.

- The vacuum oscillation length can be extracted from vacuum oscillation survival probability. It is $L_v = \frac{2\pi}{\omega}$.
- In this problem we have another energy scale which is the interaction, Δ . Here we can define another characteristic length $l_m = \frac{2\pi}{\Delta}$.
- MSW resonance happens when the two character lengths are matching to each other. Another way to put it is that the the term $\sin 2\theta(x)$ is minimized so that we have the smallest gap which leads to math $\hat{\Delta} = \cos 2\theta_v$. Equivalently this is the relation

$$l_0 = l_m \cos 2\theta_v.$$

- At resonance, we have

$$\cos 2\theta(x) = 1$$

$$\sin 2\theta(x) = 0.$$

¹⁹ No singularities possible. I assumed that $\sin \theta_v > 0$.




Figure 5: Energy Levels for MSW effect. We have the up-down symmetry since we shifted the energy by a constant to remove the identity matrix in the Hamiltonian.

This is max mixing of the states which means that at the resonance point

$$\begin{pmatrix} \nu_L(x_{reso}) \\ \nu_H(x_{reso}) \end{pmatrix} = \frac{\sqrt{2}}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \nu_e \\ \nu_x \end{pmatrix}$$

- Resonance conditions corresponds to a resonance density which is given by

$$n_e(x) = \frac{\omega}{\sqrt{2}G_F} \cos 2\theta_v \equiv n_0(E, \Delta m^2) \cos 2\theta_v,$$

where $n_0(E, \Delta m^2)$ is a characteristic number density which depends on the energy mixing angles and Δm^2 of the neutrinos.

- One should notice that to get back to the vacuum oscillation survival probability which means $\sin^2 2\theta(x) = \sin^2 2\theta_v$ we have²⁰

$$\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v = 1,$$

which leads to

$$\hat{\Delta} = 0 \quad \text{or} \quad 2 \cos 2\theta_v.$$

The first condition is trivial which corresponds to vacuum however the second condition $\Delta = 2 \cos 2\theta_v \omega$ means the interaction oscillation length is doubled compared to resonance point.

Nevertheless, we should always remember to check what survival probability the expression is describing. Here we have survival probability for $\nu_L(x)$. At $n(x) \rightarrow 0$ the oscillation becomes vacuum oscillation.

²⁰ I should check if this is still true in 3 flavor neutrino scenario. This could be interesting if it is respected in 3 flavor neutrino oscillations.

First Order Approximation

In principle we could use Taylor expansion of \mathbf{U}_m to find the series of the derivatives of $\cos \theta(x)$ and $\sin \theta(x)$, however it doesn't look like the easy way. Now we go back to find the Taylor expansion of the Hamiltonian which only involves the expression $\sin 2\theta(x)$ and $\cos 2\theta(x)$.

The series of them are

$$\begin{aligned} \sin 2\theta(x) &= \sin 2\theta_v + \cos 2\theta_v \sin 2\theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta})^2 \\ \cos 2\theta(x) &= \cos 2\theta_v - \sin^2 2\theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta})^2. \end{aligned}$$

Notice that for small θ_v the first order can be neglected since they are second or higher order small quantities.

$\sin \theta(x)$ and $\cos \theta(x)$ can be found using these results.

$$\begin{aligned}\sin \theta(x) &= \sin \theta_v + \cos^2 \theta_v \sin \theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta}^2) \\ \cos \theta(x) &= \cos \theta_v - \cos \theta_v \sin^2 \theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta}^2).\end{aligned}$$

To find out the solutions the explicit expression for \mathbf{U}_m is required. The rotation matrix is

$$\begin{aligned}\mathbf{U}_m &= \begin{pmatrix} \cos \theta(x) & \sin \theta(x) \\ -\sin \theta(x) & \cos \theta(x) \end{pmatrix} \\ &\approx \begin{pmatrix} \cos \theta_v - \cos \theta_v \sin^2 \theta_v \hat{\Delta} & \sin \theta_v + \cos^2 \theta_v \sin \theta_v \hat{\Delta} \\ -\sin \theta_v - \cos^2 \theta_v \sin \theta_v \hat{\Delta} & \cos \theta_v - \cos \theta_v \sin^2 \theta_v \hat{\Delta} \end{pmatrix}.\end{aligned}$$

The derivative of this rotation matrix with respect to x is

$$\partial_x \mathbf{U}_m = \frac{d\hat{\Delta}}{dx} \begin{pmatrix} -\cos \theta_v \sin^2 \theta_v & \cos^2 \theta_v \sin \theta_v \\ -\cos^2 \theta_v \sin \theta_v & -\cos \theta_v \sin^2 \theta_v \end{pmatrix}$$

The effective potential is²¹

²¹ This expression comes from the result of the first order approximation of $\sin \theta(x)$ and $\cos \theta(x)$.

$$\begin{aligned}\mathbf{V}_m &= -i\mathbf{U}_m^{-1}(\partial_x \mathbf{U}_m) \\ &= -i \sin \theta_v \cos \theta_v \frac{d\hat{\Delta}}{dx} \begin{pmatrix} \cos \theta_v \sin \theta_v \hat{\Delta} & 1 \\ -1 & \cos \theta_v \sin \theta_v \hat{\Delta} \end{pmatrix}.\end{aligned}$$

Since $\frac{d\hat{\Delta}}{dx} \hat{\Delta}$ is one order higher than $\hat{\Delta}$, the effective potential can be simplified to first order, which is

$$\mathbf{V}_m = -i \sin \theta_v \cos \theta_v \frac{d\hat{\Delta}}{dx} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

The equation of motion up to first order of $\hat{\Delta}$ becomes

$$i\partial_x |\Psi_m\rangle = (\mathbf{H}_{md} + \mathbf{V}_m) |\Psi_m\rangle.$$

We have already solved

$$i\partial_x |\Psi_m\rangle = \mathbf{H}_{md} |\Psi_m\rangle,$$

where the eigenstates are $|\nu_L\rangle$ and $|\nu_H\rangle$.

In general we define

$$v = -\sin \theta_v \cos \theta_v \frac{d\hat{\Delta}}{dx}$$

so that

$$\mathbf{V}_m = \begin{pmatrix} 0 & iv \\ -iv & 0 \end{pmatrix}.$$

The general solution to the equation we need to solve can be written as

$$|\Psi_m\rangle = C_L(x)e^{-i \int \omega_{m1} dx} |v_L\rangle + C_H(x)e^{-i \int \omega_{m2} dx} |v_H\rangle,$$

where

$$\begin{aligned} \omega_{m1} &= -\sqrt{\frac{\Delta^2 + \omega^2}{4} - \frac{\Delta\omega}{2} \cos 2\theta_v} \\ &= -\omega \sqrt{\left(\frac{\hat{\Delta}^2 + 1}{4} - \frac{\hat{\Delta}}{2} \cos 2\theta_v\right)}, \\ \omega_{m2} &= -\omega_{m1} \equiv \frac{\omega_m}{2}. \end{aligned}$$

Hamiltonian applied to this state results in

$$\begin{aligned} \mathbf{H}_m |\Psi_m\rangle &= \omega_{m1} C_L(x) e^{-i \int \omega_{m1} dx} |v_L\rangle - iv C_L(x) e^{-i \int \omega_{m1} dx} |v_H\rangle \\ &\quad + \omega_{m2} C_H(x) e^{-i \int \omega_{m2} dx} |v_H\rangle + iv C_H(x) e^{-i \int \omega_{m2} dx} |v_L\rangle. \end{aligned}$$

Plug the state $|\Psi_m\rangle$ into the Schrödinger equation, we have

$$\begin{aligned} \dot{C}_L(x) &= v C_H(x) e^{-i \int \delta\omega_m dx} \\ \dot{C}_H(x) &= -v C_L(x) e^{i \int \delta\omega_m dx}, \end{aligned}$$

in which $\delta\omega_m$ is defined as

$$\delta\omega_m = \omega_{m2} - \omega_{m1}.$$

The first order differential equations of $C_L(x)$ and $C_H(x)$ can be combined and produce a second order differential equation.

$$\ddot{C}_L - \left(\frac{\dot{v}}{v} - i\delta\omega_m\right) \dot{C}_L + v^2 C_L = 0.$$

In our linear approximation that $\frac{d\hat{\Delta}}{dx}$ is a constant.²² The equation simplifies to

$$\ddot{C}_L + i\delta\omega_m \dot{C}_L + v^2 C_L = 0,$$

where $v = -\sin\theta_v \cos\theta_v \frac{d\hat{\Delta}}{dx}$ is constant.

²² We are assuming that $n(x)$ is linearly depending on x which means $\hat{\Delta}$ is a linear function of x . Thus $\frac{d\hat{\Delta}}{dx}$ is a constant.

Assuming one of the solutions is e^{rx} , we could insert it to the second order differential equation and get the characteristic equation,

$$r^2 + i\delta\omega_m r + v^2 = 0,$$

which has two solutions

$$r_1 = i \frac{-\delta\omega_m + \sqrt{\delta\omega_m^2 + 4v^2}}{2}$$

$$r_2 = i \frac{-\delta\omega_m - \sqrt{\delta\omega_m^2 + 4v^2}}{2}.$$

The solutions to $C_L(x)$ and $C_H(x)$ will be

$$C_L(x) = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

$$C_H(x) = (c_1 e^{r_1 x} + c_2 e^{r_2 x}) \frac{e^{i \int \delta\omega_m dx}}{v}.$$

These constants are determined by the initial condition. The state of the system at any x is

$$|\Psi_m(x)\rangle = (c_1 e^{r_1 x} + c_2 e^{r_2 x}) e^{-i \int_0^x \omega_{m1}(x') dx'} |\nu_L\rangle$$

$$+ (c_1 e^{r_1 x} + c_2 e^{r_2 x}) \frac{e^{i \int \delta\omega_m dx}}{v} e^{-i \int_0^x \omega_{m2}(x') dx'} |\nu_H\rangle.$$

Suppose we have the initial condition as $|\Psi_m(x=0)\rangle = |\nu_L\rangle$, the system can jump to $|\nu_H\rangle$ since the state at arbitrary position x is a mixing of the two states.²³ The rate of jumping is given by²⁴

$$\Gamma = \frac{(2v)^2}{4 \left(\frac{d\omega_m}{dx} \right)} = \frac{v^2}{\frac{d\omega_m}{dx}}.$$

The probability jumping to state $|\nu_H\rangle$ after the level crossing (at infinite x) is

$$P(x \rightarrow \infty, |\nu_L\rangle \rightarrow |\nu_H\rangle) = e^{-2\pi\Gamma}$$

$$= \exp \left(-2\pi \frac{v^2}{\frac{d\omega_m}{dx}} \right)$$

$$= \exp \left(-2\pi \frac{\left(-\sin\theta_v \cos\theta_v \frac{d\hat{\Delta}}{dx} \right)^2}{\frac{d\omega_m}{dx}} \right).$$

The survival probability can be calculated by counting the probability left on the initial state.

²³ In fact this approximation is valid at the level crossing.

²⁴ Amar C Vutha. A simple approach to the Landau-Zener formula. 389: 4, 2010. ISSN 0143-0807. DOI: 10.1088/0143-0807/31/2/016. URL <http://arxiv.org/abs/1001.3322>; C. Zener. Non-Adiabatic Crossing of Energy Levels. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 137 (833):696–702, 1932. ISSN 1364-5021. DOI: 10.1098/rspa.1932.0165; and Jan R. Rubbmark, Michael M. Kash, Michael G. Littman, and Daniel Kleppner. Dynamical effects at avoided level crossings: A study of the Landau-Zener effect using Rydberg atoms. *Physical Review A*, 23(6):3107–3117, 1981. ISSN 10502947. DOI: 10.1103/PhysRevA.23.3107

To be clear, if electron neutrinos are produced inside core of our sun, it will be the light state. Since the interaction with matter is very strong, it transfers to $|\nu_L\rangle$ with probability $P(x \rightarrow \infty, |\nu_L\rangle \rightarrow |\nu_H\rangle)$ due to the gradient of the matter profile which works as the perturbation. Thus the final state will be a mixing of $|\nu_L\rangle$ and $|\nu_H\rangle$.

General Discussion for Neutrinos Interacting with Matter

This part is a very general discussion of the matter effect.²⁵

To work in flavor basis, we use the subscript $_{mf}$ to denote the flavor basis representation with mass effect. The equation of motion in flavor basis can be written down as²⁶

$$i\partial_x \Psi_{mf}(x) = \mathbf{H}_{mf} \Psi_{mf}(x)$$

where

$$\mathbf{H}_{mf} = \left(\frac{\Delta}{2} - \frac{\omega}{2} \cos 2\theta_v \right) \sigma_3 + \frac{\omega}{2} \sin 2\theta_v \sigma_1.$$

As we have seen in adiabatic situation, the states will stay in heavy and light states all along the evolution if the system starts from heavy or light state,

$$\begin{aligned} |\nu_{a1}(x)\rangle &= \exp(-i \int_0^x \frac{\omega_m(x)}{2} dx) |\nu_L(x)\rangle \\ |\nu_{a2}(x)\rangle &= \exp(i \int_0^x \frac{\omega_m(x)}{2} dx) |\nu_H(x)\rangle, \end{aligned}$$

where the heavy and light states are defined in the adiabatic situation previously. [This is what happens before the passing through of the resonance.](#)

However, when it comes to the non-adiabatic transitions, the evolution of the states will be the mixture of the heavy and light state since transitions occurs $|\nu_{a1}\rangle \rightarrow |\nu_1(x)\rangle$ and $|\nu_{a2}\rangle \rightarrow |\nu_2(x)\rangle$,

$$\begin{aligned} |\nu_1(x)\rangle &= a_L \exp(-i \int_{x_r}^x \omega_m(x')/2 dx') |\nu_L(x)\rangle + a_H \exp(i \int_{x_r}^x \omega_m(x')/2 dx') |\nu_H(x)\rangle \\ |\nu_2(x)\rangle &= b_L \exp(-i \int_{x_r}^x \omega_m(x')/2 dx') |\nu_L(x)\rangle + b_H \exp(i \int_{x_r}^x \omega_m(x')/2 dx') |\nu_H(x)\rangle, \end{aligned}$$

where the relations between the constants are determined using the condition that $|\nu_1(x)\rangle$ and $|\nu_2(x)\rangle$ are orthonormal²⁷, which leads to the conclusion that²⁸

$$\begin{aligned} b_L &= -a_H^* \\ b_H &= a_L^* \\ |a_L|^2 &= -|a_H|^2. \end{aligned}$$

²⁵ Stephen J. Parke. Nonadiabatic Level Crossing in Resonant Neutrino Oscillations. *Physical Review Letters*, 57(10):1275–1278, September 1986. ISSN 0031-9007. DOI: 10.1103/PhysRevLett.57.1275. URL <http://link.aps.org/doi/10.1103/PhysRevLett.57.1275>

²⁶ Writing down the dimensionless equation, I have

$$i\partial_x \Psi_{mf} = \frac{R_S \omega}{2} ((\hat{\Delta} - \cos 2\theta_v) \sigma_3 + \sin 2\theta_v \sigma_1) \Psi_{mf}.$$

²⁷ I don't have a good argument for this requirement.

²⁸ It is easy to verify the results.

Electron neutrinos are produced in a dense region as $|\nu_e\rangle$, which are partially transformed to other the other neutrinos due to matter and the resonance then it propagates as if it satisfies the adiabatic condition again. The initial state in terms of light and heavy state is²⁹

²⁹ The relation between θ_m and θ_v is given by

$$\omega_m \sin 2\theta_m = \omega \sin 2\theta_v.$$

$$|\Psi_m(x_0)\rangle = |\nu_e\rangle = \cos \theta_m(x_0) |\nu_L(x_0)\rangle + \sin \theta_m(x_0) |\nu_H(x_0)\rangle.$$

The final state right before the resonance is

$$|\Psi_m(x_{r-})\rangle = \cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) |\nu_L(x_{r-})\rangle + \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) |\nu_H(x_{r-})\rangle$$

After the resonance the state is described by the general jumping

$$\begin{aligned} |\Psi_m(x)\rangle = & \cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) \left(a_L \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_L(x)\rangle + a_H \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_H(x)\rangle \right) \\ & + \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) \left(-a_H^* \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_L(x)\rangle + a_L^* \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_H(x)\rangle \right) \end{aligned}$$

in which the x_{r-} is actually x_r thus

$$\begin{aligned} |\Psi_m(x)\rangle = & \cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_r} \frac{\omega_m(x)}{2} dx\right) \left(a_L \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_L(x)\rangle + a_H \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_H(x)\rangle \right) \\ & + \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_r} \frac{\omega_m(x)}{2} dx\right) \left(-a_H^* \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_L(x)\rangle + a_L^* \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') |\nu_H(x)\rangle \right) \end{aligned}$$

To calculate the survival probability it is easier to use flavor basis, hence we have another form of $|\Psi_m(x)\rangle$ which is

$$\begin{aligned} |\Psi_m(x)\rangle = & \left[\cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx'\right) a_L \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right. \\ & \left. - \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x')}{2} dx'\right) a_H^* \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right] |\nu_L(x)\rangle \\ & + \left[\cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx'\right) a_H \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right. \\ & \left. + \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x')}{2} dx'\right) a_L^* \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right] |\nu_H(x)\rangle \\ = & \left[\cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_r} \frac{\omega_m(x)}{2} dx\right) a_L \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right. \\ & \left. - \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) a_H^* \exp(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right] (\cos \theta_m(x) |\nu_e\rangle - \sin \theta_m(x) |\nu_x\rangle) \\ & + \left[\cos \theta_m(x_0) \exp\left(-i \int_{x_0}^{x_r} \frac{\omega_m(x)}{2} dx\right) a_H \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right. \\ & \left. + \sin \theta_m(x_0) \exp\left(i \int_{x_0}^{x_{r-}} \frac{\omega_m(x)}{2} dx\right) a_L^* \exp(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx') \right] (\sin \theta_m(x) |\nu_e\rangle + \cos \theta_m(x) |\nu_x\rangle) \end{aligned}$$

Survival amplitude of electron neutrinos is given by³⁰

³⁰ $\cos \theta_m$, $\sin \theta_m$ and ω_m are real but a_L and a_H are complex.

$$\begin{aligned}
& \langle \Psi_m(0) | \Psi_m(x) \rangle \\
&= \langle \nu_e | \Psi_m(x) \rangle \\
&= \left[\cos \theta_m(x_0) \exp \left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) a_L \exp \left(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) \right. \\
&\quad \left. - \sin \theta_m(x_0) \exp \left(i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) a_H^* \exp \left(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) \right] \cos \theta_m(x) \\
&\quad + \left[\cos \theta_m(x_0) \exp \left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) a_H \exp \left(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) \right. \\
&\quad \left. + \sin \theta_m(x_0) \exp \left(i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) a_L^* \exp \left(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) \right] \sin \theta_m(x) \\
&= A_L \exp \left(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) + A_H \exp \left(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right),
\end{aligned}$$

where the coefficients are

$$\begin{aligned}
A_L(x) &= \cos \theta_m(x) \left[a_L \cos \theta_m(x_0) \exp \left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) - a_H^* \sin \theta_m(x_0) \exp \left(i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) \right] \\
A_H(x) &= \sin \theta_m(x) \left[a_H \cos \theta_m(x_0) \exp \left(-i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) + a_L^* \sin \theta_m(x_0) \exp \left(i \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx' \right) \right].
\end{aligned}$$

The detection is in a region where matter density is very small, thus we use $x \rightarrow \infty$ which means the effective mixing angle becomes vacuum mixing angle. The probability is the square of the amplitude,

$$\begin{aligned}
P(\nu_e, x) &= |\langle \Psi_m(0) | \Psi_m(x) \rangle|^2 \\
&= |A_L(x) \exp \left(-i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right) + A_H(x) \exp \left(i \int_{x_r}^x \frac{\omega_m(x')}{2} dx' \right)|^2 \\
&= |A_L(x)|^2 + |A_H(x)|^2 + A_L^*(x) A_H(x) \exp(2i\phi) + A_H^*(x) A_L(x) \exp(-2i\phi) \\
&= |A_L(x)|^2 + |A_H(x)|^2 + 2\text{Re}(A_L^*(x) A_H(x) \exp(2i\phi)),
\end{aligned}$$

where ϕ is a constant

$$\phi = \int_{x_0}^{x_r} \frac{\omega_m(x')}{2} dx'$$

Note that for any complex number $(a + ib)e^{i\phi} \equiv \rho e^{i\psi}$,

$$(a + ib)e^{i\phi} + c.c. = 2\rho \cos(\psi + \phi),$$

which means that the previous result can be simplified to

$$\begin{aligned} P(\nu_e, x) &= |A_L(x)|^2 + |A_H(x)|^2 + 2\text{Re}(A_L^*(x)A_H(x)\exp(2i\phi)) \\ &= |A_L(x)|^2 + |A_H(x)|^2 + 2|A_L^*(x)A_H(x)|\cos(2\phi + \psi_{LH}), \end{aligned}$$

with the definition that $\psi_{LH}(x)$ is the argument of $A_L^*(x)A_H(x)$.

However the coefficients a_L and a_H are still not known yet. The trick is to average over the detection and production. The average over x removes the cos term and averages $\cos^2 \theta_m(x)$ to $\frac{1}{2}$, which results in

$$\begin{aligned} \langle P(\nu_e, x) \rangle_x &= \cos^2 \theta_m(x) (|a_H|^2 \cos^2 \theta_m(x_0) + |a_L|^2 \sin^2 \theta_m(x_0)) \\ &\quad + \sin^2 \theta_m(x) (|a_H|^2 \cos^2 \theta_m(x_0) + |a_L|^2 \sin^2 \theta_m(x_0)) \\ &\quad + (-\cos^2 \theta_m(x) + \sin^2 \theta_m(x)) \cos \theta_m(x_0) \sin \theta_m(x_0) (a_H a_L e^{-2i\phi} + \text{c.c.}). \end{aligned}$$

Applying the condition that $|a_L|^2 + |a_H|^2 = 1$, the probability becomes

$$\langle P(\nu_e, x) \rangle_x = \frac{1}{2} + \frac{1}{2}(1 - 2|a_H|^2) \cos 2\theta_m(x_0) \cos 2\theta_v - |a_H a_L| \sin 2\theta_m(x_0) \cos 2\theta_v \cos(2\phi + \psi_{LH}),$$

where ψ_{LH} is the argument of $a_H a_L$.

Notice that in fact the detection happens in vacuum, which means $\theta_m(x) = \theta_v$. The average over production removes the last part,³¹

$$\langle \langle P(\nu_e, x) \rangle_x \rangle_{x_0} = \frac{1}{2} + \frac{1}{2}(1 - 2|a_H|^2) \cos 2\theta_m(x_0) \cos 2\theta_v.$$

Recall that the adiabatic result is of the form

$$P(\nu_e, x)_{\text{adiabatic}} = \frac{1}{2}(1 + \cos 2\theta_m \cos 2\theta_v).$$

Define a transition probability at resonance

$$P_r(\nu_L \rightarrow \nu_H) = |a_2|^2,$$

which is determined by the Landau-Zener transition to the first order.³²

Numerical Method and Results

To simplify the codes, I prefer to write down all quantities dimensionless. In this spirit, the Hamiltonian becomes

$$\mathbf{H}_m \mathbf{d} = \begin{pmatrix} \omega_{m1} & 0 \\ 0 & \omega_{m2} \end{pmatrix} + \mathbf{V}_m,$$

³¹ In these averages it is important to keep the angles not averaged because these are the densities around the production and detection. The average only gets rid of the integral formed phase.

³² As paper by Parke, the probability is

$$P_r(\nu_L \rightarrow \nu_H) = \exp \left(-\frac{\pi}{2} \frac{\sin^2 2\theta_v}{\cos 2\theta_v} \frac{\omega}{\left| \frac{1}{N} \frac{dN}{dx} x_r \right|} \right)$$

where

$$\omega_{m2} = -\omega_{m1} \equiv \frac{\omega_m}{2} = \omega \sqrt{\frac{\hat{\Delta}^2 + 1}{4} - \frac{\hat{\Delta}}{2} \cos 2\theta_v}$$

while the effective potential \mathbf{V}_m is

$$\begin{aligned} \mathbf{V}_m &= -i\mathbf{U}_m^{-1}\partial_x\mathbf{U}_m \\ &= -i \begin{pmatrix} \cos\theta(x) & -\sin\theta(x) \\ \sin\theta(x) & \cos\theta(x) \end{pmatrix} \partial_x \begin{pmatrix} \cos\theta(x) & \sin\theta(x) \\ -\sin\theta(x) & \cos\theta(x) \end{pmatrix} \end{aligned}$$

in which

$$\begin{aligned} \sin 2\theta(x) &= \frac{\sin 2\theta_v}{\sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}} \\ \cos 2\theta(x) &= \frac{\cos 2\theta_v - \hat{\Delta}}{\sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}}. \end{aligned}$$

and

$$\hat{\Delta} = \frac{\sqrt{2}G_F n(x)}{\omega}.$$

To compare with the theory, the first order effective potential is

$$\begin{aligned} \mathbf{V}_m &= -i \sin\theta_v \cos\theta_v \frac{d\hat{\Delta}}{dx} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ &= -i \sin\theta_v \cos\theta_v \sqrt{2} \frac{G_F}{\omega} \frac{dn(\hat{x})}{d\hat{x}} \frac{1}{R_S} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \end{aligned}$$

Now we could write down the dimensionless equation

$$i\partial_{\hat{x}}\Psi(x) = (R_S\mathbf{H}_m + R_S\mathbf{V}_m)\Psi(x),$$

which becomes a matrix equation given a set of basis

$$i\partial_{\hat{x}} \begin{pmatrix} \nu_L \\ \nu_H \end{pmatrix} = \left(\frac{R_S\omega}{2} \begin{pmatrix} -\hat{\omega}_m & 0 \\ 0 & \hat{\omega}_m \end{pmatrix} - i \sin\theta_v \cos\theta_v \sqrt{2} \frac{G_F}{\omega} \frac{dn(\hat{x})}{d\hat{x}} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right) \begin{pmatrix} \nu_L \\ \nu_H \end{pmatrix}$$

The solar electron number density from standard solar model is given in arXiv:astro-ph/001034.³³

The data file³⁴ is listed in units of R_S and cm^{-3} while the density is actually $\log(n(x)/N_A)$ where N_A is Avogadro number.

Before any numerical calculations, I would like to do a simple analysis of the equation. First thing to do is to see if the second term in Hamiltonian is too small.

³³ Data from this work is listed <http://www.sns.ias.edu/jnb/SNdata/sndata.html> while the data file is actually in this file here <http://www.sns.ias.edu/jnb/SNdata/Export/BP2000/neordered.output>.

³⁴ <http://www.sns.ias.edu/jnb/SNdata/Export/BP2000/neordered.output>

The radius of the sun is about $R_S = 695800\text{km}$ which can be approximated as³⁵

$$R_S = 7 \times 10^{23}\text{fm} = \frac{7 \times 10^{23}}{0.2\text{GeV}} = 3.5 \times 10^{24}\text{GeV}^{-1}.$$

The frequency $\omega = \Delta m^2/2E$ is approximately

$$\omega = \frac{10^{-4}\text{eV}^2}{10^{-3}\text{GeV}} = 10^{-19}\text{GeV}.$$

Thus

$$R_S \omega \approx 10^5$$

Notice that the dimensionless frequency ω_m is

$$\hat{\omega}_m = \sqrt{\hat{\Delta}^2 + 1 - 2\hat{\Delta} \cos 2\theta_v}.$$

In the situation of the sun, where $\hat{\Delta} \ll 13^6$, we keep the first order of this quantity,

$$\begin{aligned} \frac{\hat{\omega}_m}{2} &= \sqrt{\frac{\hat{\Delta}^2 + 1}{4} - \frac{\hat{\Delta}}{2} \cos 2\theta_v} \\ &= \frac{1}{2} - \frac{1}{2} \cos 2\theta_v \hat{\Delta} + \mathcal{O}(\hat{\Delta})^3. \end{aligned}$$

The derivative of number density of the sun $\frac{dn(\hat{x})}{d\hat{x}}$ is shown to be of the order -10^{26}cm^{-3} ³⁷, i.e.,

$$\frac{dn(\hat{x})}{d\hat{x}} \approx -10^{26}\text{cm}^{-3} \approx -10^{-23}\text{GeV}^3.$$

We also have

$$\frac{G_F}{\omega} \approx \frac{10^{-5}\text{GeV}^{-2}}{10^{-19}\text{GeV}} = 10^{14}\text{GeV}^{-3}$$

The second term is of order³⁸

$$\frac{G_F}{\omega} \frac{dn(\hat{x})}{d\hat{x}} \approx -10^{14}\text{GeV}^{-3} \times 10^{-23}\text{GeV}^3 = 10^{-9}.$$

So this means that I can not use the data from this paper since the data starts from too far away from the center.

Another way of doing the estimation is to find the resonance density, which means a maximum mixing of the two states, as the diagonal elements disappears,³⁹

$$\Delta - \omega \cos 2\theta_v = 0.$$

³⁵ The conversion between SI and natural units is done by the relation $1\text{fm} = 1/0.2\text{GeV}$.

³⁶ Put in the numbers,

$$\hat{\Delta} \approx 10^{-19} \sim 10^{-11}$$

³⁷ $1\text{cm}^{-3} = 8 \times 10^{-51}\text{GeV}^3$

³⁸ $G_F = 1.17 \times 10^{-5}\text{GeV}^{-2}$

³⁹ The Hamiltonian in flavor basis is

$$\mathbf{H} = \frac{\omega}{2} (-\cos 2\theta_v \sigma_3 + \sin 2\theta_v \sigma_1) + \frac{\Delta}{2} \sigma_3 + \Delta \mathbf{I}$$

By solving this equation, the number density of electrons should be of the order

$$n \approx \frac{\omega}{G_F} \approx 10^{-14} \text{GeV}^3.$$

In terms of cm^{-3} , the results would be 10^{36}cm^{-3} .

Combining the two relations between density and pressure, we can find the density profile with respect to radius.

Numerical Calculations for Matter Effect in Flavor Basis

The previous method is somewhat more complicated in the calculations. Hence we work in flavor basis.

The equation of motion in flavor basis is

$$i\partial_x \Psi_{mf}(x) = \mathbf{H}_{mf} \Psi_{mf}(x)$$

where

$$\mathbf{H}_{mf} = \left(\frac{\Delta}{2} - \frac{\omega}{2} \cos 2\theta_v \right) \sigma_3 + \frac{\omega}{2} \sin 2\theta_v \sigma_1.$$

Writing down the dimensionless equation, I have

$$i\partial_{\hat{x}} \Psi_{mf} = \frac{R_S \omega}{2} ((\hat{\Delta} - \cos 2\theta_v) \sigma_3 + \sin 2\theta_v \sigma_1) \Psi_{mf}.$$

As for the data of the sun I use a simple exponential distribution. The data is also from the paper by Bahcall which is shown in figure 6.

The model using just exponential is not accurate however it is enough to make the point in MSW resonance. **Unfortunately, using this model the center density is still not enough for a MSW resonance.**

Thus I choose to use an artificial sun which is dense enough for MSW resonance, in which the core density is $n(0) = 10^{-14} \text{GeV}^{-3}$. The distribution is

$$n = 10^{-14-4.3\hat{x}} \text{GeV}^{-3}.$$

The numerical results can be obtained by plugging this result into the differential equation solver. The result is shown in figure 7.

Ternary Diagrams

Decapitated - Solar Density Profile

The solar density profile can be determined using simple gravitational theory and nuclear physics models.




Figure 6: Solar electron density from Bahcall. Horizontal axis is the distance from the core of the sun normalized by the radius of the sun while the vertical axis is the number density of electrons in $\log_{10}(n/N_A)$. The best fit for the line is

$$2.3 - 4.3\hat{x}.$$

So the equation of the number density distribution is

$$n = N_A 10^{2.3-4.3\hat{x}}.$$




Figure 7: Numerical results for electron flavor neutrino when the electron density profile is $10^{-14-4.3\hat{x}}\text{GeV}^{-3}$.




Figure 8: Numerical results for electron flavor neutrino probability and the other flavor neutrino probability when the electron density profile is $10^{-14-4.3\hat{x}}\text{GeV}^{-3}$.




Figure 9: $\theta_{12} = 33.36/180^\circ\pi$;
 $\theta_{13} = 8.66/180^\circ\pi$;
 $\theta_{23} = 40/180^\circ\pi$;
 $\delta_{\text{acp}} = 0$;
 $m_{1s} = 0.01$;
 $m_{2s} = m_{1s} + 0.000079$;
 $\text{oneFourE} = 100$;




Figure 10: Reduce θ_{12} to half of the original value
 $\theta_{12} = 33.36/180^\circ\pi \cdot 1/2$;
 $\theta_{13} = 8.66/180^\circ\pi$;
 $\theta_{23} = 40/180^\circ\pi$;
 $\delta_{\text{acp}} = 0$;
 $m_{1s} = 0.01$;
 $m_{2s} = m_{1s} + 0.000079$;
 $\text{oneFourE} = 100$;


Neutrino Flavor Oscillations for $x \in [0,1000]$ 

Figure 11: Reduce θ_{13}
 $\theta_{12} = 33.36/180^\circ\text{Pi}$;
 $\theta_{13} = 8.66/180^\circ\text{Pi} \cdot 1/2$;
 $\theta_{23} = 40/180^\circ\text{Pi}$;
 $\delta_{\text{tcp}} = 0$;
 $m_{1\text{sq}} = 0.01$;
 $m_{2\text{sq}} = m_{1\text{sq}} + 0.000079$;
 $\text{oneFourE} = 100$;


Neutrino Flavor Oscillations for $x \in [0,1000]$ 

Figure 12: Reduce θ_{23}
 $\theta_{12} = 33.36/180^\circ\text{Pi}$;
 $\theta_{13} = 8.66/180^\circ\text{Pi}$;
 $\theta_{23} = 40/180^\circ\text{Pi} \cdot 1/2$;
 $\delta_{\text{tcp}} = 0$;
 $m_{1\text{sq}} = 0.01$;
 $m_{2\text{sq}} = m_{1\text{sq}} + 0.000079$;
 $\text{oneFourE} = 100$;




Figure 13: Increase $\delta m_{22}^2 - \delta m_{11}^2$
 $\theta_{12} = 33.36/180^\circ \text{Pi}$;
 $\theta_{13} = 8.66/180^\circ \text{Pi}$;
 $\theta_{23} = 40/180^\circ \text{Pi}$;
 $\delta m_{cp} = 0$;
 $m_{1sq} = 0.01$;
 $m_{2sq} = m_{1sq} + 0.000079 \times 10$;
 $\text{oneFourE} = 100$;




Figure 14: Starting from all ν_μ
 $\theta_{12} = 33.36/180^\circ \text{Pi}$;
 $\theta_{13} = 8.66/180^\circ \text{Pi}$;
 $\theta_{23} = 40/180^\circ \text{Pi}$;
 $\delta m_{cp} = 0$;
 $m_{1sq} = 0.01$;
 $m_{2sq} = m_{1sq} + 0.000079$;
 $\text{oneFourE} = 100$;

The mass contained in a spherical shell dr is

$$dm = 4\pi r^2 \rho dr, \quad (37)$$

which is assuming spherical symmetry.

The gravitational potential for each unit of mass can generate force as gravitational attraction, which is

$$\frac{dv}{dr} = \frac{Gm}{r^2}.$$

The balance between pressure and gravity shows that

$$\frac{dv}{dr} \rho + \frac{dp}{dr} = 0.$$

The result of these relations is a rather intuitive relation between density and pressure, which is not solvable until we find another relation between them.

$$\frac{1}{r^2} \frac{d}{dr} \left(\frac{r^2}{\rho} \frac{dp}{dr} \right) = -4\pi G \rho.$$

To be simple, we will use Eddington model to solve the density profile, which gives us the relation

$$p = K \rho^{4/3},$$

where

$$K = \left(\left(\frac{k}{m_p} \frac{3}{\alpha} \frac{\beta}{\mu^4 (1-\beta)^4} \right)^4 \right)^{1/3}.$$

References

Stephen J. Parke. Nonadiabatic Level Crossing in Resonant Neutrino Oscillations. *Physical Review Letters*, 57(10):1275–1278, September 1986. ISSN 0031-9007. DOI: 10.1103/PhysRevLett.57.1275. URL <http://link.aps.org/doi/10.1103/PhysRevLett.57.1275>.

Jan R. Rubbmark, Michael M. Kash, Michael G. Littman, and Daniel Kleppner. Dynamical effects at avoided level crossings: A study of the Landau-Zener effect using Rydberg atoms. *Physical Review A*, 23(6):3107–3117, 1981. ISSN 10502947. DOI: 10.1103/PhysRevA.23.3107.

Amar C Vutha. A simple approach to the Landau-Zener formula. 389:4, 2010. ISSN 0143-0807. DOI: 10.1088/0143-0807/31/2/016. URL <http://arxiv.org/abs/1001.3322>.

C. Zener. Non-Adiabatic Crossing of Energy Levels. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 137(833):696–702, 1932. ISSN 1364-5021. DOI: 10.1098/rspa.1932.0165.