

# **Synthesized b0 for diffusion distortion correction (Synb0-DisCo)**

Kurt G. Schilling, Justin Blaber, Yuankai Huo, Allen Newton, Colin Hansen, Vishwesh Nath, Andrea T. Shafer, Owen Williams, Susan M. Resnick, Baxter Rogers, Adam W. Anderson, Bennett A. Landman

Magnetic Resonance Imaging  
2019

# Table of content

1. Goal of the paper
2. Acquisition of dataset
3. Methodology
4. Results
5. Conclusion

## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)


- Goal of the paper:

Enabling TOPUP processing with limited diffusion imaging data



Proposing to synthesize an undistorted EPI image from the structural image and use the non-distorted synthetic image as an anatomical target

- ✓ Using a database of pairs of structural images and multi-shot EPI images



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ▪ Acquisition of dataset:

- ✓ 586 pairs of T1-weighted and diffusion b0 EPI brain images from healthy controls
- ✓ Acquiring T1-weighted images using an MPRAGE sequence
- ✓ Acquiring diffusion acquisition using a single shot EPI sequence
- ✓ To obtain a non-distorted image with b0 contrast



Acquiring a high resolution, multi-shot EPI image

## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

- **Methodology:**

- Acquiring b0 synthesis:

- ✓ Registering the multi-shot EPI b0 scans to the paired T1-weighted MPAGE scan

- ➡ Transforming the paired multi-shot EPI scans to MNI space


- ✓ Registering the MPAGE scans for each subject to the MNI-152 1mm isotropic T1-weighted atlas




Reducing the variation in the data

## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ■ Training:


Three separate GANs  One network learning mapping of sagittal 2D slices  
The second learning mapping of coronal 2D slices  
The third learning mapping of axial 2D slices



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)



### ■ Pre-Training:

- ✓ Pre-training three separate orthogonal pix2pix networks
- ✓ Using all subjects' datasets to create sets of three consecutive slices in a paired manner



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ■ Pre-Training:






## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ▪ Application of the network:

- ✓ Registering MPAGE scans to align with the MNI space
- ✓ Processing MPAGE to creating overlapping sets of three contiguous slices
- ✓ Applying the corresponding generative models to each stack independently




## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ▪ Application of the network:

- ✓ Producing three output channels for each stack of images by the generative model
- ✓ Averaging the channels for each reconstruction to reconstruct an estimated b0 volume
- ✓ Combining the reconstructions from the three orientations by applying median filtering

Resulting in the synthesis of a b0 image for the subject



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ▪ Distortion correction:

- ✓ Registering the synthesized b0 image to the real (actual) b0 image



Aligning the synthetic and real b0 images

- ✓ Concatenating the registered synthesized b0 image and the real b0 image




Using combined dataset as input for TOPUP

- ✓ Further processing steps

**TOPUP:** estimating and correcting for susceptibility-induced distortions in the MRI data

## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)


- **Results:** Test on healthy subject scanned at Vanderbilt University



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

- **Results:** Test on healthy subject scanned at Vanderbilt University

Anatomically-corrected T1 scans







PE-P

PE-A

TOPUP

Synthetic b0 Correction  
(using only PE-A)



## ❖ Synthesized b0 for diffusion distortion correction (Synb0-DisCo)

### ■ Conclusion:

- ✓ Utilizing the proposed network in the absence of additional reverse PE acquisitions, and only requiring a standard T1-weighted acquisition
- ✓ Reasonable distortion corrections similar to state-of-the art methods that require blip-up blip-down acquisitions
- ✗ Creating a slight blurring of fine details and structures by median filtering of sagittal, axial, and coronal slices
- ✗ Incapability to predict the appropriate b0 contrast in certain regions of non-healthy populations (for example, tumors)

# Thank you