

# Diagnostic Biomarkers for Posttraumatic Stress Disorder: Promising Horizons from Translational Neuroscience Research

Vasiliki Michopoulos, Seth Davin Norrholm, and Tanja Jovanovic

## ABSTRACT

Posttraumatic stress disorder (PTSD) is a heterogeneous disorder that affects individuals exposed to trauma (e.g., combat, interpersonal violence, and natural disasters). Although its diagnostic features have been recently reclassified with the emergence of the Diagnostic and Statistical Manual for Mental Disorders, Fifth Edition, the disorder remains characterized by hyperarousal, intrusive reminders of the trauma, avoidance of trauma-related cues, and negative cognition and mood. This heterogeneity indicates the presence of multiple neurobiological mechanisms underlying the etiology and maintenance of PTSD. Translational research spanning the past few decades has revealed several potential avenues for the identification of diagnostic biomarkers for PTSD. These include, but are not limited to, monoaminergic transmitter systems, the hypothalamic-pituitary-adrenal axis, metabolic hormonal pathways, inflammatory mechanisms, psychophysiological reactivity, and neural circuits. The current review provides an update to the literature with regard to the most promising putative PTSD biomarkers, with specific emphasis on the interaction between neurobiological influences on disease risk and symptom progression. Such biomarkers will most likely be identified by multi-dimensional models derived from comprehensive descriptions of molecular, neurobiological, behavioral, and clinical phenotypes.

**Keywords:** Biomarkers, Inflammation, Neuroendocrinology, Neuroimaging, Psychophysiology, PTSD

<http://dx.doi.org/10.1016/j.biopsych.2015.01.005>

Posttraumatic stress disorder (PTSD) is a severe psychiatric disorder that occurs after a psychological traumatic life event and increases individual vulnerability to adverse health outcomes (1). PTSD is heterogeneous, often presenting across different symptom domains, including re-experiencing, avoidance/numbing, and hyperarousal symptoms (2). While extensive work has successfully identified psychological, genomic, and biological risk factors that are associated with PTSD in trauma survivors (3–5), the identification of discrete diagnostic biomarkers for PTSD remains elusive. The lack of diagnostic biomarkers for PTSD is not due to a lack of intensive study but rather likely due to the complexity of PTSD and the complex set of rules by which we classify individuals according to the 5th edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5), as illustrated by the recent description of 636,120 different ways in which an individual can be diagnosed with PTSD (6). Furthermore, PTSD is associated with significant mental health (e.g., major depression, substance and alcohol abuse, panic disorder, suicide) and general medical (e.g., diabetes, cardiovascular disease) comorbidities (7,8), which can obscure the search for diagnostic biomarkers for PTSD. Given that DSM-5 criteria are not based on the underlying biology, PTSD research could benefit significantly from the new approach to mental health diagnoses using the Research Domain Criteria (RDoC) (9). One of the tenets of this approach is dimensional analyses of neurobiological metrics and symptoms, rather than diagnostic classification. The putative

biomarkers listed in this review are reflective of the extant literature but can also serve RDoC objectives in future studies by linking PTSD symptoms to relevant biological underpinnings.

The vast heterogeneity inherent in PTSD symptom presentation makes it highly unlikely that a valid, singular biomarker will be identified for PTSD (10,11). However, comprehensive biological phenotyping of the factors associated with PTSD may yield a parsimonious diagnostic model with which to diagnose PTSD in the future. The current review will highlight several biomarkers associated with PTSD symptomatology and vulnerability, in addition to underscoring how individual factors, such as one's comorbid diagnoses and gender, must be considered, as they can profoundly influence biology and thus influence our search for true biomarkers of PTSD. Specifically, we will emphasize monoamine, neuroendocrine, inflammatory, genetic, epigenetic, psychophysiological, neuroanatomical, and neuroactivation phenotypes associated with PTSD to illustrate the potential efficacy of using multidimensional phenotypic data to characterize unique profiles of PTSD.

## MONOAMINE SYSTEMS IN PTSD

PTSD is characterized by increased sympathetic nervous system tone that is coincident with augmented levels of catecholamine secretion (12). Urinary and central levels of norepinephrine (NE) are heightened in individuals with PTSD (13) and in child trauma victims (14), and peripheral and central

levels of NE in response to threatening stimuli are also elevated in PTSD (15,16). Recent evidence suggests that this increase in NE in PTSD is due to attenuated levels of the NE transporter within the brainstem locus coeruleus (17). PTSD has also been associated with decreased expression of peripheral  $\alpha$ 2-adrenergic receptors, receptors that underlie an auto-receptor-driven mechanism that serves to inhibit synaptic transmitter release (18). Further, facilitation of NE release via blockade of presynaptic  $\alpha$ 2-adrenergic receptors with the antagonist yohimbine can produce panic attacks and an increase in anxiety- and trauma-related symptoms in individuals with PTSD (19,20). A prospective study of motor vehicle accident survivors indicated that urinary levels of NE were associated with increased development of PTSD 1 month following trauma but only in men (21), indicating that gender may be important for characterizing catecholaminergic biomarkers of PTSD. Increased catecholamines, however, are also coincident with panic attacks and other fear-related psychopathology (22), indicating that increased sympathetic activation is not a specific biomarker of PTSD but rather of a common neurobiological feature of fear- and anxiety-related disorders.

Alterations in the serotonergic system have also been implicated in the pathophysiology of PTSD. Individuals with PTSD show decreased levels of paroxetine binding, suggesting that levels of the serotonin transporter are attenuated in PTSD (23) and involved in the manifestation of arousal and avoidance symptoms (24). Empirical evidence has shown that serotonin transporter expression within the amygdala is attenuated in PTSD and is significantly associated with higher anxiety and depressive symptoms (25). Brainstem and forebrain levels of the serotonin 1A receptor are higher in individuals with PTSD (26), similar to what has been described in depression (27). Likewise, reductions in central serotonin 1B receptors in trauma-exposed individuals are associated with increased PTSD and depression symptoms (25). Taken together, these data indicate that alterations within the serotonergic system could reveal putative biomarkers for depressive symptoms common to both PTSD and major depression (26). The effectiveness of selective serotonin reuptake inhibitors (e.g., sertraline) for reducing the symptoms of PTSD (28–30), major depression, and other psychiatric conditions with which PTSD is highly comorbid (2,22) further suggests that more careful examination of serotonergic phenotypes is warranted to better disentangle the specificity of biomarkers for PTSD- and depression-specific phenotypes.

One way in which to elucidate the specificity of monoaminergic biomarkers on PTSD symptomology is to concurrently characterize sympathetic and serotonergic function within the same individuals. Using a repeated-measures design, Southwick *et al.* (20) found that both yohimbine and meta-chlorophenylpiperazine treatment increased panic attacks, anxiety, and trauma-related symptoms in veterans diagnosed with PTSD in a manner that suggested at least two different biological subtypes of PTSD, thus underscoring the need for more robust phenotyping of biological factors including the monoaminergic transmitter systems.

## NEUROENDOCRINE BIOMARKERS OF PTSD


Dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis is present in PTSD and has been extensively characterized (Figure 1) [for review see (31)]. Evidence suggests that

individuals with PTSD have attenuated levels of basal cortisol (31) and that a low level of cortisol in trauma survivors is associated with increased risk for subsequent development of PTSD (32,33). However, findings on baseline cortisol levels have been mixed, and a recent meta-analysis concluded that there are no consistent differences between PTSD and control subjects (34). Similarly, equivocal results exist surrounding the cortisol response to acute cognitive stressors, as reports show heightened or no differences in cortisol response to a stressor (35,36). In part, these discordant HPA results appear to be due to different sampling methods, the diurnal rhythm of cortisol release, and confounding analyses that have disregarded the influence of sex on HPA activity (37).

Rather than focus on baseline cortisol, a more promising approach is to measure cortisol reactivity to a challenge. Blunted cortisol reactivity to acute stress exposure is associated with increased prospective risk for PTSD (38). Low cortisol levels in PTSD have been coupled to enhanced glucocorticoid negative feedback inhibition of the HPA axis as evidenced by increased suppression of cortisol levels following a dexamethasone suppression test (39). This enhanced HPA negative feedback in PTSD is coincident with 1) augmented levels of peripheral and central corticotropin-releasing hormone (40,41); 2) elevated glucocorticoid receptor (GR) levels (42); 3) increased glucocorticoid sensitivity (43); and 4) decreased levels of FKBP5 (44), a co-chaperone of GR that inhibits ligand binding and nuclear translocation of GRs. A recent prospective study indicates that augmented baseline GR levels and diminished FKBP5 messenger RNA levels are associated with increased risk for PTSD symptoms following trauma (45).

While extensive work has alluded to HPA-based biomarkers of PTSD, it is clear that additional neuroendocrine factors influence PTSD vulnerability and symptomology (Figure 1; Table 1). For example, menstrual cycle phase (46,47) and pregnancy (48) influence PTSD symptom expression profile and psychophysiology in women, suggesting that ovarian steroid hormones are important modulators of PTSD susceptibility and symptom presentation. Indeed, low levels of estradiol are associated with impaired fear extinction in PTSD (49), and high levels of pituitary adenylate cyclase-activating polypeptide, a peptide implicated in stress-related behavior and physiology (50–52), are associated with PTSD only in women (53). Furthermore, central levels of the anxiolytic neuroactive steroid allopregnanolone, a potent modulator of gamma-aminobutyric acidergic inhibition, are decreased in women with PTSD (54). Low levels of testosterone in men, on the other hand, have prospectively been associated with increased rates of PTSD (55) and increased risk for PTSD (56). These data, along with epidemiologic studies strongly suggesting that female sex is a risk factor for psychopathology (including PTSD) (57) and reinforce the need to better understand the influence of gonadal steroid hormones in men and women with PTSD.

An additional avenue of exploration with regard to PTSD and putative biomarkers is the expression and regulation of metabolic hormones in individuals with PTSD. Neuropeptide Y (NPY) is an orexigenic peptide neurotransmitter (58) that also shows anxiolytic properties via antagonism of corticotropin-releasing hormone and noradrenergic systems (59). Trauma



**Figure 1.** A summary of key biomarkers that are associated with posttraumatic stress disorder (PTSD), highlighting the interactions between different biological systems that influence and complicate biological phenotypes within PTSD. Gonadal steroid hormones and the hypothalamic-pituitary-adrenal (HPA) axis modulate neurotransmitter and neuropeptide systems (145), influence amygdala activity (146,147), and influence inflammatory responses (93). HPA activity, via cortisol and corticotropin-releasing hormone (CRH), alters sensitivity to gonadal hormones (148). Inflammation alters HPA activity and has adverse effects on cardiovascular function (149). Taken together, these data indicate that as a field we must begin to study these physiological systems in concert with one another to begin to characterize comprehensive biological phenotypes of PTSD. ALLO, allopregnanolone; AVP, arginine vasopressin; CRP, C-reactive protein; E2, estradiol; HR, heart rate; HRV, heart rate variability; 5HT1A R, serotonin 1A receptor; 5HTT, serotonin transporter; IL1beta, interleukin 1 beta; IL2, interleukin 2; IL6, interleukin 6; NE, norepinephrine; NFkB, nuclear factor-κB; NPY, neuropeptide Y, T4, testosterone.

exposure (60) and PTSD (59) are associated with attenuated peripheral levels of NPY, and conversely, resilience to trauma is associated with increased NPY levels (61). Ghrelin, an orexigenic peptide secreted from the stomach (58), displays fear-enhancing effects in rodents (62) and could serve as a biomarker of trauma exposure and PTSD. More recently, individuals with PTSD have shown a hyperinsulinemic response to an oral glucose challenge (63). Finally, peripheral endocannabinoid levels (64) are reduced and central cannabinoid receptor type 1 receptors (65) are increased in PTSD.

In summary, it is clear that significant progress has been made in identifying and characterizing PTSD-related neuroendocrine perturbations. However, the majority of these neuroendocrine factors have been studied in isolation in traumatized

populations exhibiting PTSD signs and symptoms, and as such, it is important to characterize multilevel neuroendocrine profiles of PTSD accounting for parallel trauma-related neuroendocrinologic changes, their interaction, and the relationship to stress exposure or resilience. For example, increases in dehydroepiandrosterone and dehydroepiandrosterone-sulfate have been linked to PTSD symptom expression but are also associated with decreased levels of affective symptoms and PTSD severity (66,67). Thus, it has been suggested that the ratio of these adrenal hormones to cortisol might be important for resilience to stress and recovery from PTSD (68,69). Furthermore, elucidating the complex interaction of neuroendocrine factors (i.e., allopregnanolone/estradiol/NPY effects on cortisol) on the regulation of the HPA axis will likely expand

**Table 1. Neuroendocrine Biological Factors Associated with PTSD**

Neuroendocrine Biomarkers	Relationship to PTSD	References
<b>HPA-Axis</b>		
Glucocorticoid negative feedback	Augmented in PTSD	(39)
Baseline cortisol	Attenuated in PTSD	(31,32)
Acute cortisol following trauma	Lower levels increase risk for PTSD	(33,38)
Pituitary adenylate cyclase-activating polypeptide	Increased in women with PTSD	(53)
<b>Steroid Hormones</b>		
Estradiol	Reduced levels increase risk for PTSD and are associated with impaired fear extinction	(49)
Allopregnanolone	Decreased in women with PTSD	(54)
Dehydroepiandrosterone	Increased in PTSD	(68)
Dehydroepiandrosterone sulfate	Increased in PTSD; high DHEAS increases risk for PTSD	(68)
Testosterone	Low levels increase risk for PTSD	(56)
<b>Metabolic Hormones</b>		
NPY	Decreased in PTSD	(59)
Ghrelin	Increases fear in rodents	(62)
Insulin	Increased response to glucose in PTSD	(63)
Endocannabinoids	Decreased in PTSD	(64)

DHEAS, dehydroepiandrosterone sulfate; HPA, hypothalamic-pituitary-adrenal; NPY, neuropeptide Y; PTSD, posttraumatic stress disorder.

our ability to further describe PTSD-specific and may prove beneficial in characterizing biological subprofiles of PTSD (Figure 1). For instance, avoidance symptoms in male veterans with PTSD (70) may be related to arginine vasopressin levels and, as such, may serve as a biomarker for increased aggression in men with PTSD (71).

### BIOMARKERS OF HEIGHTENED INFLAMMATION IN PTSD

The high comorbidity between PTSD, physical illness (7), and inflammation spanning cardiovascular (72) and metabolic disease (73) has led to investigations of the relationship between inflammatory markers and PTSD symptomology (Table 2). Pro-inflammatory cytokines (i.e., proteins), including interleukin (IL)-6 (74), IL-1 $\beta$  (75), and IL-2 (76), are elevated in individuals with PTSD and peripheral levels of inflammatory markers correlate positively with PTSD symptomology (Figure 1) (77). C-reactive protein (CRP) levels are also elevated in individuals with PTSD (78–80). More specifically, increased CRP levels have been reported with exacerbated PTSD symptoms and impaired inhibition of fear-potentiated startle in the presence of a safety signal (79), a

psychophysiological biomarker for PTSD described in a later section of this review (81).

In addition, individuals with PTSD also show altered immune cell sensitivity to glucocorticoids that results in increased inflammation (82). Lysozyme enzyme activity is more sensitive to dexamethasone in PTSD (43), indicating that innate immune efficiency is higher in individuals with PTSD. Enhanced monocyte sensitivity to glucocorticoids in individuals with PTSD is also coincident with hypocortisolemia and can lead to increased cytokine production (83). The transcriptional factor, nuclear factor- $\kappa$ B (NF- $\kappa$ B), lays upstream of cytokine activation (84) and is activated by exposure to psychosocial stress (85), as well as noradrenergic activity (85), and thus may be critically sensitive to immune changes following trauma exposure. Individuals with PTSD show augmented NF- $\kappa$ B gene expression (86) and NF- $\kappa$ B activity (87).

Overall, the cross-sectional data linking PTSD to a pro-inflammatory state further support the notion that PTSD is associated with chronic inflammation (Table 2) and suggest that inflammation may serve as a possible therapeutic target for alleviating PTSD symptoms. However, increased inflammation is a hallmark of depression (88) and other adverse health outcomes that are comorbid with PTSD (7,72,73,89), thus complicating the view that immune factors may serve as

**Table 2. Immunological Factors Associated with PTSD**

Immune Biomarkers	Relationship to PTSD	References
Interleukin-6	Increased in PTSD	(74)
Interleukin-1 $\beta$	Increased in PTSD	(75)
Interleukin-2	Increased in PTSD	(76)
C-Reactive Protein	Increased in PTSD; increases risk for PTSD	(78,79,94)
Nuclear Factor- $\kappa$ B	Increased in PTSD	(86,87)
Tumor Necrosis Factor- $\alpha$	Increased in PTSD	(144)
Immune Cell Sensitivity to Glucocorticoids	Enhanced in PTSD	(43)

PTSD, posttraumatic stress disorder.



diagnostic biomarkers for PTSD specifically. This point is further highlighted by other reports that have described no differences or decreases in pro-inflammatory markers, such as CRP, in individuals with PTSD (90–92). Factors such as gender should also be considered in our examination of immunological biomarkers, as there are clear sex differences in immune system function and risk for infection (93). Finally, it is still unclear whether increased inflammation is a consequence of trauma exposure and PTSD or whether a baseline pro-inflammatory state increases individual vulnerability to PTSD after trauma exposure. As such, baseline inflammation may serve as a biomarker of PTSD vulnerability, as recent evidence from a prospective study indicates that predeployment levels of CRP significantly predict postdeployment PTSD (94).

### GENETIC AND EPIGENETIC BIOMARKERS OF PTSD

Genetic loci within genes critical for the neuroendocrine regulation of the HPA axis and emotional behavior have been associated with increased risk for PTSD [see review (5)]. However, these genetic loci have been associated with other psychiatric conditions as well, indicating that these genetic polymorphisms are not specific to PTSD but rather may serve as biomarkers for stress-induced psychopathology in general or common underlying symptoms. There are several recent genomic reviews of PTSD [e.g., (95)] and the disorders with which it is comorbid and, as such, will not be discussed at length in the current review. We will simply note that the emerging genetic and epigenetic findings related to PTSD risk versus resilience have focused on modulators of HPA axis function (before and following trauma), e.g., *FKBP5*, *PACAP*.

### PSYCHOPHYSIOLOGICAL BIOMARKERS OF PTSD

Hyperarousal symptoms, which include some of the long-standing, hallmark symptoms of PTSD, can be strongly influenced by an individual's autonomic response following trauma; the output of the autonomic nervous system can be indexed noninvasively via psychophysiological assessments of peripheral targets, such as heart rate (HR), blood pressure, skin conductance (SC), respiration rate, muscle contractions using electromyography (EMG) (e.g., startle), and body temperature. However, the use of these psychophysiological measures as biomarkers of PTSD may rely heavily on the timing and context in which they are collected. For instance, while some reports indicate that HR in the immediate aftermath of trauma exposure is predictive of later PTSD development (96), others suggest this is not the case (97,98). These equivocal findings suggest that a more robust and controlled measurement of psychophysiological data may be necessary (99). Indeed, HR and SC changes in response to a challenge have been repeatedly associated with a diagnosis of PTSD (100–104).

Exaggerated startle response, a hyperarousal symptom that remains central to DSM-5 based PTSD diagnosis, is readily assessed by psychophysiology. Increased HR (SC and EMG less so) reactivity to startling loud tones has been found to reliably differentiate PTSD from non-PTSD (105). Heightened HR reactivity to loud tones does not appear to be pre-existing but rather is acquired with the development of PTSD (106,107). Whereas heightened HR reactivity to loud tones appears to be

an acquired marker, there is accumulating evidence that heightened SC reactivity to loud tones is a pretrauma risk marker for posttraumatic stress (108,109). Exaggerated arousal can manifest as sleep disturbances, which are frequently observed in PTSD (110,111); however, the diagnostic specificity of these disturbances are not yet understood.

To examine and further explore hyperreactivity following trauma, Pitman *et al.* (112) modified an imagery procedure originally developed to study phobias (113). In this method, psychophysiological data are recorded from participants while listening to a script of their actual traumatic event. This method has been used with divergent PTSD populations, including several combat populations (114–116) and a heavily traumatized civilian population (117). In all trauma survivors, PTSD patients exhibit a stronger HR and SC response to scripts than non-PTSD trauma survivors. In studies using script-driven imagery, SC was found to be the most sensitive measure of hyperarousal in PTSD. In 1998, Keane *et al.* (101) published the results of the largest study (multisite Veterans Affairs Cooperative Study with Vietnam veterans) to date examining the utility of psychophysiological measures in diagnosing PTSD. The study employed script-driven imagery coupled with psychophysiological recordings. While this study did not find a perfect correlation between interview-based PTSD diagnosis and psychophysiological reactivity, they concluded that psychophysiological data did provide useful and objective assessment of the disorder. Recent re-analyses of script-driven imagery data collected in the 1990s have shown high specificity for PTSD (i.e., 90% of individuals without PTSD classified correctly) (102) and high concordance with subjective distress (118), but sensitivity to PTSD diagnosis remained at approximately 60% (102). Simply talking about autobiographical trauma appears to have similar effects as script-driven imagery in increasing physiological arousal (118). These methods are currently being standardized as common data elements to promote generation of large datasets using the same approach. Technological advances have afforded the opportunity to employ physiological indices that can be easily obtained in most clinical settings and may prove beneficial in the diagnosis and treatment evaluation of PTSD. A recent application of these methods using virtual reality techniques to provide immersive trauma-related imagery during recording of psychophysiological responses showed utility of this approach in tracking treatment outcomes (119).

The findings described previously support the notion that the etiology and maintenance of the fear-related symptoms of PTSD can be characterized according to the principles of fear conditioning (120,121). Given the richness of the translational literature, the neural underpinnings of fear conditioning are well understood, and PTSD research can capitalize on these findings (122). Fear conditioning is based on a simple Pavlovian conditioning model in which a neutral conditioned stimulus (CS) (for example, a light) is paired with an aversive unconditioned stimulus (US) (for example, electric shock). After a number of pairings, an association is formed such that the CS alone elicits a conditioned response (for example, a fear response). Following initial acquisition, conditioned fear is subject to consolidation, extinction, and reconsolidation, all of which may be disturbed in PTSD (123–125). Fear conditioned responses can be measured with peripheral outcomes

such as SC (123) or EMG startle responses (81). These psychophysiological measures can be used to index both the increase in fear during conditioning, as well as the reduction of fear during extinction, or the repeated presentation of the CS without the US. In addition, these measures can be used in differential conditioning studies using a CS+ cue predicting danger (US) and a CS− predicting safety from the US; these have shown that PTSD, but not depression, is associated with a reduced ability to inhibit fear-potentiated startle responses to safety signals (81). Similarly, retention of the extinction memory has been tested using SC 24 hours after fear extinction, and PTSD subjects have exhibited reduced levels of extinction recall (125). Taken together, these studies indicate the fear responses to traumatic memories may be serve as biomarkers specific for dysregulated fear in PTSD.

### NEUROANATOMICAL AND NEUROACTIVATIONAL BIOMARKERS OF PTSD

Neuroimaging data gathered during the last decade demonstrate that PTSD is associated with greater amygdala activation compared with control subjects (126). Functional magnetic resonance imaging (fMRI) studies have shown that trauma-relevant words increase amygdala activation in PTSD subjects more than in control subjects (127–130). Exaggerated fear responses observed in PTSD may be due to a weakened inhibitory control of the amygdala by the medial prefrontal cortex (mPFC). A large number of imaging studies have indicated that this inhibitory neurocircuit is dysregulated in patients with PTSD (126,128,130). A recent meta-analysis of imaging studies during emotion processing in PTSD, social anxiety, and specific phobia indicated that the rostral anterior cingulate cortex (ACC) is less active in PTSD patients relative to control subjects, an effect not found in other anxiety disorders (131).

Neuroimaging studies using fear conditioning paradigms demonstrate that fear acquisition and extinction of fear activate the prefrontal cortex, specifically the ventromedial prefrontal cortex (vmPFC) (132). For example, activation of the vmPFC (which includes the rostral ACC) is decreased in PTSD patients during an extinction recall in fMRI task (133). The vmPFC also differs in shape and size in PTSD patients (134). To date, one of the most replicated neuroanatomical findings in PTSD has been reduced hippocampal volume (135,136). Early studies of twins discordant for trauma exposure suggested that smaller hippocampal volume likely confers individual vulnerability to PTSD (137); however, a recent prospective study found that hippocampal reductions were acquired with trauma exposure (136). Finally, methods using higher resolution imaging techniques have indicated that reductions in specific subregions of the hippocampus, such as the cornu ammonis 3 and dentate gyrus, are associated with PTSD symptoms (138). Studies of neural activation have used several fMRI paradigms to activate the mPFC; the simplest and most commonly used tasks involve response inhibition. In such tasks, the participant is presented with a stimulus indicating that a response is required, for example, to press a button when a letter appears on the monitor. This is referred to as a Go signal. On a minority of trials, however, the

participant is required to withhold a response during a NoGo signal (the Go/NoGo task). The Go/NoGo task has been used in subjects with PTSD with fMRI and it reliably indicates decreased activation in the rostral vmPFC and rostral ACC in PTSD subjects compared with control subjects (139,140). Weakened mPFC control of the amygdala may be a risk factor for trauma-related psychopathology: a recent study of children with depressed parents found a lack of ACC activation to the emotional Stroop, using both fear-relevant words depicting physical threat as well as social threat (141).

### SUMMARY AND CONCLUSIONS

To date, an array of putative biomarkers associated with PTSD risk and symptom progression have been identified across distinct biological domains, including, but not limited to, alterations and differences in monoaminergic systems, neuroendocrinology, inflammation, genomics, psychophysiology, and neuroanatomy. However, the heterogeneity inherent in PTSD symptom presentation and the common comorbidity with other psychiatric and general medical conditions represent formidable obstacles in the identification of valid biomarkers specifically for PTSD when considered as a diagnostic categorization (10,11). Indeed, the likelihood of characterizing one biological marker associated with the suggested 636,120 different ways in which an individual can present with PTSD (6) is vanishingly small. Rather, it is more prudent that future studies develop a cross-dimensional, comprehensive biological and psychological phenotypic profile in individuals with PTSD to 1) characterize biomarkers for specific clusters of symptoms; and/or 2) uncover divergent biological profiles of PTSD using more complex statistical techniques (142). To be compatible with the RDoC approach, biomarkers should be dimensional as well as transdiagnostic—in effect, not biomarkers specific to PTSD as a DSM-5 disorder but biomarkers of features associated with PTSD. For example, physiological measures of fear responses would be relevant to other fear-related disorders such as phobias in addition to PTSD. Similarly, deficient prefrontal activity could be associated with PTSD symptoms, as well as addiction, and could clarify common bases for comorbid disorders.

To begin collecting comprehensive phenotypes necessary for such analyses, the importance of studying the interaction between biological factors (e.g., cellular, molecular, genetic, neurotransmitter, endocrine) (Figure 1) needs to be emphasized, most notably as they relate to physiology and behaviors underlying complex biological phenotypes within PTSD. It is important to note that biology is dynamic. Thus, it is critical for the field to understand that biomarkers might be relevant at one time point (HR immediately following trauma exposure) and not at another (11). Lastly, the implications of characterizing diagnostic biomarkers for PTSD must be carefully considered to ensure that the benefits outweigh the costs (143).

In summary, the available biological and translational data point to promising new horizons for diagnostic biomarkers of PTSD symptoms. It is most likely that such biomarkers will represent a panel of several measures that will combine molecular with behavioral and clinical information to increase specificity and sensitivity of these tools.

## ACKNOWLEDGMENTS AND DISCLOSURES

This work was funded in part by the Brain and Behavior Foundation (formerly National Alliance for Research on Schizophrenia and Depression) (SDN and TJ); the Department of Defense/Congressionally Directed Medical Research Program (Award W81XWH-08-2-0170) (SDN); the Emory University Research Committee; a Public Health Service Grant (UL1 RR025008) from the Clinical and Translational Science Award program, National Institutes of Health, National Center for Research Resources (SDN); and National Institute of Mental Health (MH098212 and MH100122) (TJ). This material is the result of work supported with resources and the use of facilities at the Atlanta Veterans Affairs Medical Center, Decatur, Georgia. The contents of this manuscript do not represent the views of the United States Department of Veterans Affairs or the United States Government.

Dr. Michopoulos, Dr. Norrholm, and Dr. Jovanovic reported no biomedical financial interests or potential conflicts of interest.

## ARTICLE INFORMATION

From the Department of Psychiatry and Behavioral Sciences (VM, SDN, TJ), Emory University School of Medicine, Atlanta; Yerkes National Primate Research Center (VM), Atlanta; and Atlanta Veterans Affairs Medical Center (SDN), Mental Health Service Line, Decatur, Georgia.

Address correspondence to Tanja Jovanovic, Ph.D., Emory University School of Medicine, Department of Psychiatry and Behavioral Sciences, 49 Jesse Hill Jr NE, Atlanta, GA 30303; E-mail: tjovano@emory.edu.

Received Oct 30, 2014; revised Jan 7, 2015; accepted Jan 15, 2015.

## REFERENCES

- Dedert EA, Calhoun PS, Watkins LL, Sherwood A, Beckham JC (2010): Posttraumatic stress disorder, cardiovascular, and metabolic disease: A review of the evidence. *Ann Behav Med* 39:61–78.
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995): Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060.
- Breslau N (2001): The epidemiology of posttraumatic stress disorder: What is the extent of the problem? *J Clin Psychiatry* 62:16–22.
- Roberts AL, Galea S, Austin SB, Cerda M, Wright RJ, Rich-Edwards JW, Koenen KC (2012): Posttraumatic stress disorder across two generations: Concordance and mechanisms in a population-based sample. *Biol Psychiatry* 72:505–511.
- Pitman RK, Rasmusson AM, Koenen KC, Shin LM, Orr SP, Gilbertson MW, et al. (2012): Biological studies of post-traumatic stress disorder. *Nat Rev Neurosci* 13:769–787.
- Galatzer-Levy IR, Bryant RA (2013): 636,120 ways to have post-traumatic stress disorder. *Perspect Psychol Sci* 8:651–662.
- Boscarino JA (2004): Posttraumatic stress disorder and physical illness: Results from clinical and epidemiologic studies. *Ann N Y Acad Sci* 1032:141–153.
- Jacobsen LK, Southwick SM, Kosten TR (2001): Substance use disorders in patients with posttraumatic stress disorder: A review of the literature. *Am J Psychiatry* 158:1184–1190.
- Cuthbert BN, Insel TR (2013): Toward the future of psychiatric diagnosis: The seven pillars of RDoC. *BMC Med* 11:126.
- Zoladz PR, Diamond DM (2013): Current status on behavioral and biological markers of PTSD: A search for clarity in a conflicting literature. *Neurosci Biobehav Rev* 37:860–895.
- Schmidt U, Kaltwasser SF, Wotjak CT (2013): Biomarkers in post-traumatic stress disorder: Overview and implications for future research. *Dis Markers* 35:43–54.
- Southwick SM, Bremner JD, Rasmusson A, Morgan CA 3rd, Arnsten A, Charney DS (1999): Role of norepinephrine in the pathophysiology and treatment of posttraumatic stress disorder. *Biol Psychiatry* 46:1192–1204.
- Geraciotti TD Jr, Baker DG, Ekhtor NN, West SA, Hill KK, Bruce AB, et al. (2001): CSF norepinephrine concentrations in posttraumatic stress disorder. *Am J Psychiatry* 158:1227–1230.
- Delahanty DL, Nugent NR, Christopher NC, Walsh M (2005): Initial urinary epinephrine and cortisol levels predict acute PTSD symptoms in child trauma victims. *Psychoneuroendocrinology* 30:121–128.
- Blanchard EB, Kolb LC, Prins A, Gates S, McCoy GC (1991): Changes in plasma norepinephrine to combat-related stimuli among Vietnam veterans with posttraumatic stress disorder. *J Nerv Ment Dis* 179:371–373.
- Geraciotti TD Jr, Baker DG, Kasckow JW, Strawn JR, Mulchahey JJ, Dashevsky BA, et al. (2008): Effects of trauma-related audiovisual stimulation on cerebrospinal fluid norepinephrine and corticotropin-releasing hormone concentrations in post-traumatic stress disorder. *Psychoneuroendocrinology* 33:416–424.
- Pietrzak RH, Gallezot JD, Ding YS, Henry S, Potenza MN, Southwick SM, et al. (2013): Association of posttraumatic stress disorder with reduced in vivo norepinephrine transporter availability in the locus coeruleus. *JAMA Psychiatry* 70:1199–1205.
- Perry BD, Giller EL Jr, Southwick SM (1987): Altered platelet alpha 2-adrenergic binding sites in posttraumatic stress disorder. *Am J Psychiatry* 144:1511–1512.
- Southwick SM, Krystal JH, Morgan CA, Johnson D, Nagy LM, Nicolaou A, et al. (1993): Abnormal noradrenergic function in posttraumatic stress disorder. *Arch Gen Psychiatry* 50:266–274.
- Southwick SM, Krystal JH, Bremner JD, Morgan CA 3rd, Nicolaou AL, Nagy LM, et al. (1997): Noradrenergic and serotonergic function in posttraumatic stress disorder. *Arch Gen Psychiatry* 54:749–758.
- Hawk LW, Dougall AL, Ursano RJ, Baum A (2000): Urinary catecholamines and cortisol in recent-onset posttraumatic stress disorder after motor vehicle accidents. *Psychosom Med* 62:423–434.
- Martin EI, Ressler KJ, Binder E, Nemeroff CB (2009): The neurobiology of anxiety disorders: Brain imaging, genetics, and psychoneuroendocrinology. *Psychiatr Clin North Am* 32:549–575.
- Arora RC, Fichtner CG, O'Connor F, Crayton JW (1993): Paroxetine binding in the blood platelets of post-traumatic stress disorder patients. *Life Sci* 53:919–928.
- Maes M, Lin AH, Verkerk R, Delmeire L, Van Gastel A, Van der Planken M, Scharpe S (1999): Serotonergic and noradrenergic markers of post-traumatic stress disorder with and without major depression. *Neuropsychopharmacology* 20:188–197.
- Murrough JW, Huang Y, Hu J, Henry S, Williams W, Gallezot JD, et al. (2011): Reduced amygdala serotonin transporter binding in posttraumatic stress disorder. *Biol Psychiatry* 70:1033–1038.
- Sullivan GM, Ogden RT, Huang YY, Oquendo MA, Mann JJ, Parsey RV (2013): Higher in vivo serotonin-1a binding in posttraumatic stress disorder: A PET study with [<sup>11</sup>C]WAY-100635. *Depress Anxiety* 30:197–206.
- Parsey RV, Oquendo MA, Ogden RT, Olvet DM, Simpson N, Huang YY, et al. (2006): Altered serotonin 1A binding in major depression: A [carbonyl-<sup>11</sup>C]WAY100635 positron emission tomography study. *Biol Psychiatry* 59:106–113.
- McDougle CJ, Southwick SM, Charney DS, St James RL (1991): An open trial of fluoxetine in the treatment of posttraumatic stress disorder. *J Clin Psychopharmacol* 11:325–327.
- Nagy LM, Morgan CA, Southwick S, Charney D (1993): Open prospective trial of fluoxetine for posttraumatic stress disorder. *J Clin Psychopharmacol* 13:107–113.
- Van der Kolk BA, Dreyfuss D, Michaels M, Shera D, Berkowitz R, Fiszler R, Saxe G (1994): Fluoxetine in posttraumatic stress disorder. *J Clin Psychiatry* 55:517–522.
- Yehuda R (2005): Neuroendocrine aspects of PTSD. *Handb Exp Pharmacol* 169:371–403.
- Walsh K, Nugent NR, Kotte A, Amstadter AB, Wang S, Guille C, et al. (2013): Cortisol at the emergency room rape visit as a predictor of PTSD and depression symptoms over time. *Psychoneuroendocrinology* 38:2520–2528.
- Mouthaan J, Sijbrandij M, Luitse JS, Goslings JC, Gersons BP, Olf M (2014): The role of acute cortisol and DHEAS in predicting acute and chronic PTSD symptoms. *Psychoneuroendocrinology* 45:179–186.

34. Meewisse M, Reitsma JB, De Vries GJ, Gersons BPR, Olff M (2007): Cortisol and post-traumatic stress disorder in adults: Systematic review and meta-analysis. *Br J Psychiatry* 191:387–392.
35. Bremner JD, Vythilingam M, Vermetten E, Adil J, Khan S, Nazeer A, *et al.* (2003): Cortisol response to a cognitive stress challenge in posttraumatic stress disorder (PTSD) related to childhood abuse. *Psychoneuroendocrinology* 28:733–750.
36. Kolassa I-T, Eckart C, Ruf M, Neuner F, de Quervain DJF, Elbert T (2007): Lack of cortisol response in patients with posttraumatic stress disorder (PTSD) undergoing a diagnostic interview. *BMC Psychiatry* 7:54.
37. Freidenberg BM, Gusmano R, Hickling EJ, Blanchard EB, Bremner JD, Frye C (2010): Women with PTSD have lower basal salivary cortisol levels later in the day than do men with PTSD: A preliminary study. *Physiol Behav* 99:234–236.
38. Galatzer-Levy IR, Steenkamp MM, Brown AD, Qian M, Inslicht S, Henn-Haase C, *et al.* (2014): Cortisol response to an experimental stress paradigm prospectively predicts long-term distress and resilience trajectories in response to active police service. *J Psychiatr Res* 56:36–42.
39. Yehuda R, Boisoineau D, Lowy MT, Giller EL Jr (1995): Dose-response changes in plasma cortisol and lymphocyte glucocorticoid receptors following dexamethasone administration in combat veterans with and without posttraumatic stress disorder. *Arch Gen Psychiatry* 52:583–593.
40. de Kloet CS, Vermetten E, Geuze E, Lentjes EG, Heijnen CJ, Stalla GK, Westenberg HG (2008): Elevated plasma corticotrophin-releasing hormone levels in veterans with posttraumatic stress disorder. *Prog Brain Res* 167:287–291.
41. Baker DG, Ekhtor NN, Kasckow JW, Dashevsky B, Horn PS, Bednarik L, Geraciotti TD Jr (2005): Higher levels of basal serial CSF cortisol in combat veterans with posttraumatic stress disorder. *Am J Psychiatry* 162:992–994.
42. Matic G, Milutinovic DV, Nestorov J, Elakovic I, Jovanovic SM, Perisic T, *et al.* (2013): Lymphocyte glucocorticoid receptor expression level and hormone-binding properties differ between war trauma-exposed men with and without PTSD. *Prog Neuropsychopharmacol Biol Psychiatry* 43:238–245.
43. Yehuda R, Golier JA, Yang RK, Tischler L (2004): Enhanced sensitivity to glucocorticoids in peripheral mononuclear leukocytes in posttraumatic stress disorder. *Biol Psychiatry* 55:1110–1116.
44. Yehuda R, Cai G, Golier JA, Sarapas C, Galea S, Ising M, *et al.* (2009): Gene expression patterns associated with posttraumatic stress disorder following exposure to the World Trade Center attacks. *Biol Psychiatry* 66:708–711.
45. van Zuiden M, Geuze E, Willemen HL, Vermetten E, Maas M, Amarouchi K, *et al.* (2012): Glucocorticoid receptor pathway components predict posttraumatic stress disorder symptom development: A prospective study. *Biol Psychiatry* 71:309–316.
46. Bryant RA, Felmingham KL, Silove D, Creamer M, O'Donnell M, McFarlane AC (2011): The association between menstrual cycle and traumatic memories. *J Affect Disord* 131:398–401.
47. Glover EM, Mercer KB, Norrholm SD, Davis M, Duncan E, Bradley B, *et al.* (2013): Inhibition of fear is differentially associated with cycling estrogen levels in women. *J Psychiatry Neurosci* 38:341–348.
48. Michopoulos V, Rothbaum AO, Corwin E, Bradley B, Ressler KJ, Jovanovic T (2014): Psychophysiology and posttraumatic stress disorder symptom profile in pregnant African-American women with trauma exposure [published online ahead of print October 3]. *Arch Womens Ment Health*.
49. Glover EM, Jovanovic T, Mercer KB, Kerley K, Bradley B, Ressler KJ, Norrholm SD (2012): Estrogen levels are associated with extinction deficits in women with posttraumatic stress disorder. *Biol Psychiatry* 72:19–24.
50. Hammack SE, Cheung J, Rhodes KM, Schutz KC, Falls WA, Braas KM, May V (2009): Chronic stress increases pituitary adenylate cyclase-activating peptide (PACAP) and brain-derived neurotrophic factor (BDNF) mRNA expression in the bed nucleus of the stria terminalis (BNST): Roles for PACAP in anxiety-like behavior. *Psychoneuroendocrinology* 34:833–843.
51. Hashimoto H, Shintani N, Baba A (2006): New insights into the central PACAPergic system from the phenotypes in PACAP- and PACAP receptor-knockout mice. *Ann N Y Acad Sci* 1070:75–89.
52. Ghzili H, Grumolato L, Thouennon E, Tanguy Y, Turquier V, Vaudry H, Anouar Y (2008): Role of PACAP in the physiology and pathology of the sympathoadrenal system. *Front Neuroendocrinol* 29:128–141.
53. Ressler KJ, Mercer KB, Bradley B, Jovanovic T, Mahan A, Kerley K, *et al.* (2011): Post-traumatic stress disorder is associated with PACAP and the PAC1 receptor. *Nature* 470:492–497.
54. Rasmusson AM, Pinna G, Paliwal P, Weisman D, Gottschalk C, Charney D, *et al.* (2006): Decreased cerebrospinal fluid allopregnanolone levels in women with posttraumatic stress disorder. *Biol Psychiatry* 60:704–713.
55. Mulchahey JJ, Ekhtor NN, Zhang H, Kasckow JW, Baker DG, Geraciotti TD Jr (2001): Cerebrospinal fluid and plasma testosterone levels in post-traumatic stress disorder and tobacco dependence. *Psychoneuroendocrinology* 26:273–285.
56. Reijnen A, Geuze E, Vermetten E (2015): The effect of deployment to a combat zone on testosterone levels and the association with the development of posttraumatic stress symptoms: A longitudinal prospective Dutch military cohort study. *Psychoneuroendocrinology* 51:525–533.
57. Bromet E, Sonnega A, Kessler RC (1998): Risk factors for DSM-III-R posttraumatic stress disorder: Findings from the National Comorbidity Survey. *Am J Epidemiol* 147:353–361.
58. Keen-Rhinehart E, Ondek K, Schneider JE (2013): Neuroendocrine regulation of appetitive ingestive behavior. *Front Neurosci* 7:213.
59. Rasmusson AM, Hauger RL, Morgan CA, Bremner JD, Charney DS, Southwick SM (2000): Low baseline and yohimbine-stimulated plasma neuropeptide Y (NPY) levels in combat-related PTSD. *Biol Psychiatry* 47:526–539.
60. Morgan CA 3rd, Rasmusson AM, Winters B, Hauger RL, Morgan J, Hazlett G, Southwick S (2003): Trauma exposure rather than posttraumatic stress disorder is associated with reduced baseline plasma neuropeptide-Y levels. *Biol Psychiatry* 54:1087–1091.
61. Morgan CA 3rd, Rasmusson AM, Wang S, Hoyt G, Hauger RL, Hazlett G (2002): Neuropeptide-Y, cortisol, and subjective distress in humans exposed to acute stress: Replication and extension of previous report. *Biol Psychiatry* 52:136–142.
62. Meyer RM, Burgos-Robles A, Liu E, Correia SS, Goossens KA (2014): A ghrelin-growth hormone axis drives stress-induced vulnerability to enhanced fear. *Mol Psychiatry* 19:1284–1294.
63. Rao MN, Chau A, Madden E, Inslicht S, Talbot L, Richards A, *et al.* (2014): Hyperinsulinemic response to oral glucose challenge in individuals with posttraumatic stress disorder. *Psychoneuroendocrinology* 49:171–181.
64. Hill MN, Bierer LM, Makotkine I, Golier JA, Galea S, McEwen BS, *et al.* (2013): Reductions in circulating endocannabinoid levels in individuals with post-traumatic stress disorder following exposure to the World Trade Center attacks. *Psychoneuroendocrinology* 38:2952–2961.
65. Neumeister A, Normandin MD, Pietrzak RH, Piomelli D, Zheng MQ, Gujarr-Anton A, *et al.* (2013): Elevated brain cannabinoid CB1 receptor availability in post-traumatic stress disorder: A positron emission tomography study. *Mol Psychiatry* 18:1034–1040.
66. Sondergaard HP, Hansson LO, Theorell T (2002): Elevated blood levels of dehydroepiandrosterone sulphate vary with symptom load in posttraumatic stress disorder: Findings from a longitudinal study of refugees in Sweden. *Psychother Psychosom* 71:298–303.
67. Spivak B, Maayan R, Kotler M, Mester R, Gil-Ad I, Shtaif B, Weizman A (2000): Elevated circulatory level of GABA(A)-antagonistic neurosteroids in patients with combat-related post-traumatic stress disorder. *Psychol Med* 30:1227–1231.
68. Yehuda R, Brand SR, Golier JA, Yang RK (2006): Clinical correlates of DHEA associated with post-traumatic stress disorder. *Acta Psychiatr Scand* 114:187–193.
69. Morgan CA 3rd, Southwick S, Hazlett G, Rasmusson A, Hoyt G, Zimolo Z, Charney D (2004): Relationships among plasma dehydroepiandrosterone sulfate and cortisol levels, symptoms of dissociation, and objective performance in humans exposed to acute stress. *Arch Gen Psychiatry* 61:819–825.



70. de Kloet CS, Vermetten E, Geuze E, Wiegant VM, Westenberg HG (2008): Elevated plasma arginine vasopressin levels in veterans with posttraumatic stress disorder. *J Psychiatr Res* 42:192–198.
71. Stappenbeck CA, Hellmuth JC, Simpson T, Jakupcak M (2014): The effects of alcohol problems, PTSD, and combat exposure on non-physical and physical aggression among Iraq and Afghanistan war veterans. *Psychol Trauma* 6:65–72.
72. Boscarino JA, Chang J (1999): Electrocardiogram abnormalities among men with stress-related psychiatric disorders: Implications for coronary heart disease and clinical research. *Ann Behav Med* 21: 227–234.
73. Weiss T, Skelton K, Phifer J, Jovanovic T, Gillespie CF, Smith A, *et al.* (2011): Posttraumatic stress disorder is a risk factor for metabolic syndrome in an impoverished urban population. *Gen Hosp Psychiatry* 33:135–142.
74. Maes M, Lin AH, Delmeire L, Van Gastel A, Kenis G, De Jongh R, Bosmans E (1999): Elevated serum interleukin-6 (IL-6) and IL-6 receptor concentrations in posttraumatic stress disorder following accidental man-made traumatic events. *Biol Psychiatry* 45:833–839.
75. Spivak B, Shohat B, Mester R, Avraham S, Gil-Ad I, Bleich A, *et al.* (1997): Elevated levels of serum interleukin-1 beta in combat-related posttraumatic stress disorder. *Biol Psychiatry* 42:345–348.
76. Smith AK, Conneely KN, Kilaru V, Mercer KB, Weiss TE, Bradley B, *et al.* (2011): Differential immune system DNA methylation and cytokine regulation in post-traumatic stress disorder. *Am J Med Genet B Neuropsychiatr Genet* 156B:700–708.
77. von Kanel R, Hepp U, Kraemer B, Traber R, Keel M, Mica L, Schnyder U (2007): Evidence for low-grade systemic proinflammatory activity in patients with posttraumatic stress disorder. *J Psychiatr Res* 41:744–752.
78. Miller RJ, Sutherland AG, Hutchison JD, Alexander DA (2001): C-reactive protein and interleukin 6 receptor in post-traumatic stress disorder: A pilot study. *Cytokine* 13:253–255.
79. Michopoulos V, Rothbaum AO, Jovanovic T, Almli LM, Bradley B, Rothbaum B, *et al.* (2015): CRP levels are associated with increased PTSD symptoms and influenced by CRP genetic variation in a highly traumatized civilian population [published online ahead of print]. *Am J Psychiatry*.
80. Plantinga L, Bremner JD, Miller AH, Jones DP, Veledar E, Goldberg J, Vaccarino V (2013): Association between posttraumatic stress disorder and inflammation: A twin study. *Brain Behav Immun* 30:125–132.
81. Jovanovic T, Norrholm SD, Blanding NQ, Davis M, Duncan E, Bradley B, Ressler KJ (2010): Impaired fear inhibition is a biomarker of PTSD but not depression. *Depress Anxiety* 27:244–251.
82. Rohleder N, Wolf JM, Wolf OT (2010): Glucocorticoid sensitivity of cognitive and inflammatory processes in depression and posttraumatic stress disorder. *Neurosci Biobehav Rev* 35:104–114.
83. Rohleder N, Joksimovic L, Wolf JM, Kirschbaum C (2004): Hypocortisolism and increased glucocorticoid sensitivity of pro-inflammatory cytokine production in Bosnian war refugees with posttraumatic stress disorder. *Biol Psychiatry* 55:745–751.
84. Liang Y, Zhou Y, Shen P (2004): NF-kappaB and its regulation on the immune system. *Cell Mol Immunol* 1:343–350.
85. Bierhaus A, Wolf J, Andrassy M, Rohleder N, Humpert PM, Petrov D, *et al.* (2003): A mechanism converting psychosocial stress into mononuclear cell activation. *Proc Natl Acad Sci U S A* 100:1920–1925.
86. O'Donovan A, Sun B, Cole S, Rempel H, Lenoci M, Pulliam L, Neylan T (2011): Transcriptional control of monocyte gene expression in post-traumatic stress disorder. *Dis Markers* 30:123–132.
87. Pace TW, Wingenfeld K, Schmidt I, Meinlschmidt G, Hellhammer DH, Heim CM (2012): Increased peripheral NF-kappaB pathway activity in women with childhood abuse-related posttraumatic stress disorder. *Brain Behav Immun* 26:13–17.
88. Raison CL, Capuron L, Miller AH (2006): Cytokines sing the blues: Inflammation and the pathogenesis of depression. *Trends Immunol* 27:24–31.
89. Boscarino JA (1997): Diseases among men 20 years after exposure to severe stress: Implications for clinical research and medical care. *Psychosom Med* 59:605–615.
90. Söndergaard HP, Hansson L-O, Theorell T (2004): The inflammatory markers C-reactive protein and serum amyloid A in refugees with and without posttraumatic stress disorder. *Clin Chim Acta* 342:93–98.
91. von Kanel R, Hepp U, Kraemer B, Traber R, Keel M, Mica L, Schnyder U (2007): Evidence for low-grade systemic proinflammatory activity in patients with posttraumatic stress disorder. *J Psychiatr Res* 41:744–752.
92. McCanlies EC, Araia SK, Joseph PN, Mnatsakanova A, Andrew ME, Burchfiel CM, Violanti JM (2011): C-reactive protein, interleukin-6, and posttraumatic stress disorder symptomatology in urban police officers. *Cytokine* 55:74–78.
93. Klein SL (2000): Hormones and mating system affect sex and species differences in immune function among vertebrates. *Behav Processes* 51:149–166.
94. Eraly SA, Nievergelt CM, Maihofer AX, Barkauskas DA, Biswas N, Agorastos A, *et al.* (2014): Assessment of plasma C-reactive protein as a biomarker of posttraumatic stress disorder risk. *JAMA Psychiatry* 71:423–431.
95. Skelton K, Ressler KJ, Norrholm SD, Jovanovic T, Bradley-Davino B (2012): PTSD and gene variants: New pathways and new thinking. *Neuropharmacology* 62:628–637.
96. Shalev A, Peri T, Brandes D, Freedman S, Orr S, Pitman R (2000): Auditory startle response in trauma survivors with posttraumatic stress disorder: A prospective study. *Am J Psychiatry* 157:255–261.
97. Buckley B, Nugent N, Sledjeski E, Raimonde AJ, Spoonster E, Bogart LM, Delahanty DL (2004): Evaluation of initial posttrauma cardiovascular levels in association with acute PTSD symptoms following a serious motor vehicle accident. *J Trauma Stress* 17: 317–324.
98. Blanchard EB, Hickling EJ, Galovski T, Veazey C (2002): Emergency room vital signs and PTSD in a treatment seeking sample of motor vehicle accident survivors. *J Trauma Stress* 15:199–204.
99. Shalev AY, Sahar T, Freedman S, Peri T, Glick N, Brandes D, *et al.* (1998): A prospective study of heart rate response following trauma and the subsequent development of posttraumatic stress disorder. *Arch Gen Psychiatry* 55:553–559.
100. Blanchard EB, Kolb LC, Pallmeyer TP, Gerardi RJ (1982): A psychophysiological study of post-traumatic stress disorder in Vietnam veterans. *Psychiatr Q* 54:220–229.
101. Keane TM, Kolb LC, Kaloupek D, Blanchard EB, Thomas RG, Hsieh FY, Lavori PW (1998): Utility of psychophysiological measurement in the diagnosis of posttraumatic stress disorder: Results from a Department of Veterans Affairs Cooperative Study. *J Consult Clin Psychol* 66:914–923.
102. American Psychiatric Association (2013): *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. Washington, DC: American Psychiatric Association.
103. Orr SP, Metzger LJ, Lasko NB, Macklin ML, Hu FB, Shalev AY, Pitman RK (2003): Physiologic responses to sudden, loud tones in monozygotic twins discordant for combat exposure: Association with posttraumatic stress disorder. *Arch Gen Psychiatry* 60:283–288.
104. Shalev AY, Peri T, Brandes D, Freedman S, Orr SP, Pitman RK (2000): Auditory startle response in trauma survivors with posttraumatic stress disorder: A prospective study. *Am J Psychiatry* 157:255–261.
105. Orr SP, Metzger LJ, Miller MW, Kaloupek DG (2004): Psychophysiological assessment of PTSD. In: Wilson JP, Keane TN, editors. *Assessing Psychological Trauma and PTSD: A Handbook for Practitioners*, 2nd ed. New York: Guilford Publications, 289–343.
106. Griffin MG (2008): A prospective assessment of auditory startle alterations in rape and physical assault survivors. *J Trauma Stress* 21:91–99.
107. Orr SP, Roth WT (2000): Psychophysiological assessment: Clinical applications for PTSD. *J Affect Disord* 61:225–240.
108. Pole N, Neylan TC, Best SR, Orr SP, Marmar CR (2003): Fear-potentiated startle and posttraumatic stress symptoms in urban police officers. *J Trauma Stress* 15:471–479.
109. Orr SP, Lasko NB, Macklin ML, Pineles SL, Chang Y, Pitman RK (2012): Predicting post-trauma stress symptoms from pre-trauma psychophysiological reactivity, personality traits and measures of psychopathology. *Biol Mood Anxiety Disord* 18:8.

110. Neylan TC, Otte C, Yehuda R, Marmar CR (2006): Neuroendocrine regulation of sleep disturbances in PTSD. *Ann N Y Acad Sci* 1071: 203–215.
111. Kobayashi I, Boarts JM, Delahanty DL (2007): Polysomnographically measured sleep abnormalities in PTSD: A meta-analytic review. *Psychophysiology* 44:660–669.
112. Pitman RK, Orr SP, Forgue DF, de Jong JB, Claiborn JM (1987): Psychophysiological assessment of posttraumatic stress disorder imagery in Vietnam combat veterans. *Arch Gen Psychiatry* 44:970–975.
113. Lang PJ, Levin DN, Miller GA, Kozak MJ (1983): Fear behaviour, fear imagery, and the psychophysiology of emotion: The problem of affective response integration. *J Abnorm Psychol* 92:276–306.
114. Pitman RK, Orr SP, Forgue DF, Altman B, de Jong JB, Herz LR (1990): Psychophysiological responses to combat imagery of Vietnam veterans with posttraumatic stress disorder versus other anxiety disorders. *J Abnorm Psychol* 99:49–54.
115. Orr SP, Pitman RK, Lasko NB, Herz LR (1993): Psychophysiological assessment of posttraumatic stress disorder imagery in World War II and Korean combat veterans. *J Abnorm Psychol* 102:152–159.
116. Shalev AY, Orr SP, Pitman RK (1992): Psychophysiological response during script-driven imagery as an outcome measure in posttraumatic stress disorder. *J Clin Psychiatry* 53:324–326.
117. McTeague LM, Lang PJ, Laplante M-C, Cuthbert BN, Shumen JR, Bradley MM (2010): Aversive imagery in posttraumatic stress disorder: Trauma recurrence, comorbidity, and physiological reactivity. *Biol Psychiatry* 67:346–356.
118. Marx BP, Bovin MJ, Suvak MK, Monson CM, Sloan DM, Fredman SJ, *et al.* (2012): Concordance between physiological arousal and subjective distress among Vietnam combat veterans undergoing challenge testing for PTSD. *J Trauma Stress* 25:416–425.
119. Rothbaum BO, Price ML, Jovanovic T, Norrholm SD, Gerardi M, Dunlop B, *et al.* (2014): A randomized, double-blind evaluation of D-cycloserine or alprazolam combined with virtual reality exposure therapy for posttraumatic stress disorder in Iraq and Afghanistan war veterans. *Am J Psychiatry* 171:640–648.
120. Amstедler AB, Nugent NR, Koenen KC (2009): Genetics of PTSD: Fear conditioning as a model for future research. *Psychiatr Ann* 39:338–369.
121. Foa EB, Steketee G, Rothbaum BO (1989): Behavioral/cognitive conceptualizations of post-traumatic stress disorder. *Behav Ther* 20:155–176.
122. Jovanovic T, Ressler KJ (2010): How the neurocircuitry and genetics of fear inhibition may inform our understanding of PTSD. *Am J Psychiatry* 167:648–662.
123. Orr SP, Metzger LJ, Lasko NB, Macklin ML, Peri T, Pitman RK (2000): De novo conditioning in trauma-exposed individuals with and without posttraumatic stress disorder. *J Abnorm Psychol* 109:290–298.
124. Norrholm SD, Jovanovic T, Olin IW, Sands LA, Karapanou I, Bradley B, Ressler KJ (2011): Fear extinction in traumatized civilians with posttraumatic stress disorder: Relation to symptom severity. *Biol Psychiatry* 69:556–563.
125. Milad MR, Pitman RK, Ellis CB, Gold AL, Shin LM, Lasko NB, *et al.* (2009): Neurobiological basis of failure to recall extinction memory in posttraumatic stress disorder. *Biol Psychiatry* 66:1075–1082.
126. Hughes KC, Shin LM (2011): Functional neuroimaging studies of post-traumatic stress disorder. *Expert Rev Neurother* 11:275–285.
127. Protopopescu X, Pan H, Tuescher O, Cloitre M, Goldstein M, Engelien W, *et al.* (2005): Differential time courses and specificity of amygdala activity in posttraumatic stress disorder subjects and normal control subjects. *Biol Psychiatry* 57:464–473.
128. Rauch SL, Whalen PJ, Shin LM, McInerney SC, Macklin ML, Lasko NB, *et al.* (2000): Exaggerated amygdala response to masked facial stimuli in posttraumatic stress disorder: A functional MRI study. *Biol Psychiatry* 47:769–776.
129. Shin LM, Wright CI, Cannistraro PA, Wedig MM, McMullin K, Martis B, *et al.* (2005): A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Arch Gen Psychiatry* 62:273–281.
130. Stevens JS, Jovanovic T, Fani N, Ely TD, Glover EM, Bradley B, Ressler KJ (2013): Disrupted amygdala-prefrontal functional connectivity in civilian women with posttraumatic stress disorder. *J Psychiatr Res* 47:1469–1478.
131. Etkin A, Wager T (2007): Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *Am J Psychiatry* 164:1476–1488.
132. Phelps EA, Delgado MR, Nearing KI, LeDoux JE (2004): Extinction learning in humans: Role of the amygdala and vmPFC. *Neuron* 43: 897–905.
133. Rougemont-Bücking A, Linnman C, Zeffiro TA, Zeidan MA, Lebron-Milad K, Rodriguez-Romaguera J, *et al.* (2010): Altered processing of contextual information during fear extinction in PTSD: An fMRI study. *CNS Neurosci Ther* 17:227–236.
134. Corbo V, Clément M-H, Armony JL, Pruessner JC, Brunet A (2005): Size versus shape differences: Contrasting Voxel-based and volumetric analyses of the anterior cingulate cortex in individuals with acute posttraumatic stress disorder. *Biol Psychiatry* 58:119–124.
135. Bremner JD, Elzinga B, Schmahl C, Vermetten E (2008): Structural and functional plasticity of the human brain in posttraumatic stress disorder. *Prog Brain Res* 167:171–186.
136. Admon R, Milad MR, Hendler T (2013): A causal model of posttraumatic stress disorder: Disentangling predisposed from acquired neural abnormalities. *Trends Cogn Sci* 17:337–347.
137. Gilbertson MW, Shenton ME, Ciszewski A, Kasai K, Lasko NB, Orr SP, Pitman RK (2002): Smaller hippocampal volume predicts pathological vulnerability to psychological trauma. *Nat Neurosci* 5: 1242–1247.
138. Neylan TC, Mueller SG, Wang Z, Metzler TJ, Lenoci M, Truran D, *et al.* (2010): Insomnia severity is associated with a decreased volume of the CA3/dentate gyrus hippocampal subfield. *Biol Psychiatry* 68:494–496.
139. Carrion VG, Garrett A, Menon V, Weems CF, Reiss AL (2008): Posttraumatic stress symptoms and brain function during a response-inhibition task: An fMRI study in youth. *Depress Anxiety* 25:514–526.
140. Jovanovic T, Ely T, Fani N, Glover EM, Gutman D, Tone EB, *et al.* (2013): Reduced neural activation during an inhibition task is associated with impaired fear inhibition in a traumatized civilian sample. *Cortex* 49:1884–1891.
141. Mannie ZN, Norbury R, Murphy SE, Inkster B, Harmer CJ, Cowen PJ (2008): Affective modulation of anterior cingulate cortex in young people at increased familial risk of depression. *Br J Psychiatry* 192:356–361.
142. Neylan TC, Schadt EE, Yehuda R (2014): Biomarkers for combat-related PTSD: Focus on molecular networks from high-dimensional data. *Eur J Psychotraumatol* 5.
143. Lehrner A, Yehuda R (2014): Biomarkers of PTSD: Military applications and considerations. *Eur J Psychotraumatol* 5.
144. Gill J, Vythilingam M, Page GG (2008): Low cortisol, high DHEA, and high levels of stimulated TNF-alpha, and IL-6 in women with PTSD. *J Trauma Stress* 21:530–539.
145. Bethea CL, Lu NZ, Gundlach C, Streicher JM (2002): Diverse actions of ovarian steroids in the serotonin neural system. *Front Neuroendocrinol* 23:41–100.
146. Zeidan MA, Igoe SA, Linnman C, Vitalo A, Levine JB, Klibanski A, *et al.* (2011): Estradiol modulates medial prefrontal cortex and amygdala activity during fear extinction in women and female rats. *Biol Psychiatry* 70:920–927.
147. Goetz SMM, Tang L, Thomason ME, Diamond MP, Hariri AR, Carré JM (2014): Testosterone rapidly increases neural reactivity to threat in healthy men: A novel two-step pharmacological challenge paradigm. *Biol Psychiatry* 76:324–331.
148. Michopoulos V, Berga SL, Kaplan JR, Wilson ME (2009): Social subordination and polymorphisms in the gene encoding the serotonin transporter enhance estradiol inhibition of luteinizing hormone secretion in female rhesus monkeys. *Biol Reprod* 81:1154–1163.
149. Nijm J, Jonasson L (2009): Inflammation and cortisol response in coronary artery disease. *Ann Med* 41:224–233.