

Dissecting the processing of color vision in the primary visual cortex (V1) at the center-of-gaze

 Felix Bartsch^{1,2,3}, Jacob L. Yates^{1,4}, Cole Saborio¹, Daniel A. Butts^{1,2}, Bevil R. Conway³
¹Department of Biology, ²Program in Neuroscience and Cognitive Science, University of Maryland, College Park, MD, USA


³Laboratory of Sensorimotor Research, National Eye Institute, National Institutes of Health, Bethesda, MD, USA

⁴Herbert Wertheim School of Optometry and Vision Science, University of California, Berkeley, CA, USA

Background


How does the cortex process high-acuity vision?

High-acuity occurs in the **fovea**, within the center 1° of visual angle. Perception is at cone resolution (1 arcmin or 1/60°). Pre-cortical circuitry is at this resolution.




Foveal V1 is all but unstudied

V1 neurons are typically studied in parafovea (>2°) and higher to avoid problems with knowing the true eye position, which must be at ~arcmin resolution for fovea V1.




How is spatial and chromatic information combined in foveal V1?

V1 neurons are typically nonlinear, and mixing of spatial and chromatic information will only be clear with nonlinear models.




Spatio-chromatic stimuli




Large-scale recordings from foveal V1

Recording using 3 multielectrode arrays. Animal rewarded for fixation within 0.5°.





Computation across the cortical column

We sample a different cortical column with the V-probe in each experiment, and can use population models to understand how spatio-chromatic selectivity is constructed across the cortical column.




Model-based eye tracking

To infer the eye-position precisely, we use the population activity of foveal V1 neurons [4] combined with deep neural network models of each neuron.




STAs before and after ET correction




Spatio-chromatic selectivity in foveal V1

Utah array RF center locations




Relationship between color selectivity and RF size in fovea




Other preliminary analyses

Binocular GLMs can isolate each eye's S-C selectivity



Deep neural networks to trace the construction of S-C selectivity across (different) cortical columns



References

- [1] Hubel, D. H., & Wiesel, T. N. (2004). Brain and visual perception: the story of a 25-year collaboration. Oxford Univ Press.
- [2] Conway, B. R. (2001). Spatial structure of cone inputs to color cells in alert macaque V-1. *J Neurosci*, 21(8), 2768-83.
- [3] Hofer, H., Carroll, J., Neitz, J., Mitz, M., & Williams, D. R. (2005). Organization of the human trichromatic cone mosaic. *J Neurosci*, 25(42), 9669-9679.
- [4] McFarland JM, Cui Y, Butts DA (2013) Inferring nonlinear neuronal computation based on physiologically plausible inputs. *PLoS Comput Biol* 9(7), e1003143.
- [5] McFarland, J. M., Bondy, A. G., Cumming, B. G., & Butts, D. A. (2014). High-resolution eye tracking using V1 neuron activity. *Nat Comm*, 5(1), 1-12.
- [6] Horwitz, G. D. (2020). Signals related to color in the early visual cortex. *Annual Review of Vision Science*, 6, 287-311.
- [7] Lurz K-K, Bashiri M, Willeke K, et al. Generalization in data-driven models of V1. *bioRxiv*: 10.1101/2020.10.05.326256
- [8] Yates JL, Coop S, Wu RJ, Butts DA, Rucci M, Mitchell JF (2021) Beyond fixation: detailed characterization of neural selectivity in free-viewing primates. *bioRxiv*: 10.1101/2021.11.06.467566.

This work supported by NSF IIS-2113197 (FB, JLY, CS, DAB) and the NIH intramural program (BRC, FB).