

Methods of proof – 4

Invariants, monovariants and double counting

13 сентября 2026 года

Theoretical minimum.

Fact 1 (invariant). A quantity unchanged by the allowed operations separates the reachable states from the unreachable ones. The classical discrete invariants are parity, a residue, a sum with weights, the sign of a permutation, a colouring; the continuous ones are a first integral of a differential equation and a conserved energy (Functional and differential equations – 3); the algebraic ones are the rank, the determinant, the trace and the characteristic polynomial, which do not change under similarity (Linear algebra – 3).

Fact 2 (monovariant). A quantity that changes monotonically proves that a process terminates or that a configuration stabilises; a bounded integer monovariant gives termination, and a Lyapunov function is its continuous counterpart. The two are used in the same way: find the quantity, prove monotonicity, look at what happens at the end.

Theorem 1 (double counting). Counting the same set in two ways gives an identity; counting incidences in two ways gives an inequality. The systematic form is $\sum_v \deg v = 2|E|$ for graphs and $\sum_i |A_i| = \sum_x \#\{i : x \in A_i\}$ for set systems, and the linear-algebraic strengthening is the rank bound of Linear algebra – 6.

Fact 3 (a generating function is a ledger). Encoding a family by a polynomial or a power series turns a counting statement into an algebraic identity; the coefficient extraction by roots of unity is in Trig-2, and the analytic machinery in Series and infinite products – 4.

-
1. Prove that the 15-puzzle position obtained by swapping two neighbouring tiles is unreachable, by using the sign of the permutation together with

the parity of the position of the empty cell, and formulate the general principle for a group acting on configurations.

2. On a board the numbers may be replaced by $a, b \mapsto \frac{a+b}{\sqrt{2}}, \frac{a-b}{\sqrt{2}}$. Find an invariant, and decide which multisets are reachable from a given one; then do the same for the operation $a, b \mapsto a + b, ab$ over the reals.
3. Prove that the process in which two numbers $a \geq b$ are replaced by $a - b$ and b terminates, by exhibiting a monovariant; deduce the Euclidean algorithm and its complexity, and give a continuous analogue where the monovariant is an integral.
4. Prove $\sum_v \deg v = 2|E|$ and use double counting to prove that a graph on n vertices with no 4-cycle has at most $\frac{1}{2}(1 + \sqrt{4n - 3}) \cdot \frac{n}{2}$ edges; then prove the Oddtown theorem by the rank bound and compare the two methods (Linear algebra – 6).
5. (Putnam 2018) Let S be the set of sequences of length 2018 whose terms are in $\{1, 2, 3, 4, 5, 6, 10\}$ and sum to 3860. Prove that the cardinality of S is at most $2^{3860} \cdot \left(\frac{2018}{2048}\right)^{2018}$.