

Limits – 3

Asymptotics of sequences

1 сентября 2026 года

Theoretical minimum.

Def. Knowing the limit is not enough: an olympiad problem usually asks for the rate. The standard answer has the form $x_n \sim Cn^{-\alpha}$ or $x_n = L + Cq^n + o(q^n)$, and the choice between the two is decided by the behaviour of f at the fixed point.

Theorem 1 (neutral fixed point). Let $x_n \rightarrow 0$, $x_n > 0$ and $x_{n+1} = x_n - ax_n^p + o(x_n^p)$ with $a > 0$, $p > 1$. Then

$$x_n \sim \left(\frac{1}{a(p-1)n} \right)^{1/(p-1)}.$$

In particular $x_{n+1} = x_n - x_n^2$ gives $nx_n \rightarrow 1$, while $x_{n+1} = \sin x_n$ (where $p = 3$, $a = \frac{1}{6}$) gives $x_n \sim \sqrt{3/n}$.

Theorem 2 (Stolz–Cesàro). If b_n is strictly increasing and unbounded and $\lim \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = L$, then $\lim \frac{a_n}{b_n} = L$. This is the discrete analogue of l'Hôpital's rule, and it is exactly what proves Theorem 1: apply it to $y_n = x_n^{1-p}$.

Fact 1. The telescoping trick in powers of x_n : from $x_{n+1} = x_n - ax_n^p$ one gets $x_{n+1}^{1-p} - x_n^{1-p} \rightarrow a(p-1)$, and then Stolz finishes the job. This trick solves almost every problem of the sheet.

1. (IMC 2014) Consider the sequence $(a_n)_{n \geq 1} = (1, 1, 2, 1, 2, 3, 1, 2, 3, 4, 1, 2, 3, 4, 5, 1, \dots)$. Find all pairs (α, β) of positive reals such that $\lim_{n \rightarrow \infty} \frac{\sum_{k=1}^n a_k}{n^\alpha} = \beta$.
2. (Putnam 1966) Let $0 < x_0 < x_1$ and $x_{n+1} = x_n - x_n^2$ for $n \geq 0$. Compute $\lim_{n \rightarrow \infty} nx_n$.

3. For an arbitrary $x_0 \in (0, \pi)$ define $x_{n+1} = \sin x_n$. Compute $\lim_{n \rightarrow \infty} \sqrt{n} x_n$, and find the next term of the asymptotics.
4. (AKHIMO 2022) Let $x_1 \in (0, \frac{\pi}{2})$ and $x_{n+1} = \sqrt[3]{\frac{\sin^3 x_1 + \dots + \sin^3 x_n}{n}}$ for $n \geq 1$. Find $\lim_{n \rightarrow \infty} x_n^3 \ln n$.
5. Let $f: [a, b] \rightarrow [a, b]$ be an increasing function. Show that there exists $\xi \in [a, b]$ with $f(\xi) = \xi$, and give an example of a decreasing f for which the iterations $x_{n+1} = f(x_n)$ do not converge although the fixed point is unique.