arXiv:2611.10883v1 [math.CO] 7 Nov 2026

THE BRUALDI–QUINN CONJECTURE ON STRONG EDGE-COLORING HOLDS FRACTIONALLY AND ASYMPTOTICALLY

RONG LUO[1], GEXIN YU[2], XIANGQIAN ZHOU[3]

[1]*Department of Mathematics, West Virginia University, Morgantown, WV 26506*
[2]*Department of Mathematics, William & Mary, Williamsburg, VA 23185*
[3]*Department of Mathematics and Statistics, Wright State University, Dayton, Ohio 45435*

Abstract. Let G be a bipartite graph with parts A and B, and let Δ_A and Δ_B be the maximum degrees of the vertices in A and in B, respectively. Brualdi and Quinn conjectured that the strong chromatic index of G is at most $\Delta_A\Delta_B$. We prove that the fractional strong chromatic index of G is at most $\Delta_A\Delta_B$, and that the conjecture holds asymptotically: $\chi'_s(G) \le (1+o(1))\Delta_A\Delta_B$ as $\max\{\Delta_A,\Delta_B\} \to \infty$. The error term is $12(\ln\Delta)^{1/2}\Delta^{-1/4}$ with $\Delta = \max\{\Delta_A,\Delta_B\}$. Both results rest on a smart partition of edges and Lovász Local Lemma.

Mathematics Subject Classification (2020): 05C15, 05C70, 05D40.

Keywords: strong edge-coloring, bipartite graph, induced matching, Lovász Local Lemma.

1. Introduction

All graphs in this paper are finite and simple. A *strong edge-coloring* of a graph G is an assignment of colors to the edges of G in which every color class is an induced matching; equivalently, two edges receive distinct colors whenever they share an endpoint or are joined by an edge. The *strong chromatic index* $\chi'_s(G)$ is the minimum number of colors in a strong edge-coloring of G. The notion was introduced by Fouquet and Jolivet [11]. For a graph of maximum degree Δ, a greedy argument gives $\chi'_s(G) \le 2\Delta^2 - 2\Delta + 1$. Erdős and Nešetřil [8] conjectured that

$$\chi'_s(G) \le \begin{cases} \frac{5}{4}\Delta^2, & \text{if } \Delta \text{ is even,} \\ \frac{1}{4}(5\Delta^2 - 2\Delta + 1), & \text{if } \Delta \text{ is odd.} \end{cases}$$

Molloy and Reed [16] established the first asymptotic improvement over the greedy bound on $\chi'_s(G)$ where they proved that $\chi'_s(G) \le 1.998\Delta^2$ for sufficiently large Δ. This was successively improved to $1.93\Delta^2$ by Bruhn and Joos [4], to $1.835\Delta^2$ by Bonamy, Perrett and Postle [2], and to $1.772\Delta^2$ by Hurley, de Joannis de Verclos and Kang [14]. More recently, Davey, Hurley, de Joannis de Verclos, Kang and Volec [6] obtained the bound $1.73\Delta^2$ by way of flag algebra.

For bipartite graphs, Faudree, Gyárfás, Schelp and Tuza [9] proposed the following conjecture.

Conjecture 1 (Faudree, Gyárfás, Schelp and Tuza [9])**.** *Every bipartite graph G with maximum degree $\Delta(G)$ satisfies $\chi'_s(G) \le \Delta(G)^2$.*

Brualdi and Quinn [3] proposed a stronger, asymmetric version. Let G be bipartite with parts A and B, and let

$$\Delta_A = \max_{x\in A} d(x), \qquad \Delta_B = \max_{b\in B} d(b).$$

Conjecture 2 (Brualdi and Quinn [3])**.** *Every bipartite graph G satisfies $\chi'_s(G) \le \Delta_A\Delta_B$.*

E-mail address: rluo@mail.wvu.edu, gyu@wm.edu, xiangqian.zhou@wright.edu.

Both conjectures are tight: if $|A| = \Delta_B$ and $|B| = \Delta_A$, the complete bipartite graph on $A \cup B$ has $\Delta_A\Delta_B$ edges and all edges need distinct colors.

Steger and Yu [19] proved Conjecture 1 for $\Delta(G) = 3$. Conjecture 2 is confirmed when one of Δ_A, Δ_B is at most 2 (Nakprasit [17]) or at most 3 (Huang, Yu and Zhou [13]). For large degrees, Hao, Yang and Yu [12] proved $\chi'_s(G) \le 1.676\,\Delta_A\Delta_B$ when $\Delta_A\Delta_B$ is sufficiently large. Davey, Hurley, de Joannis de Verclos, Kang and Volec [6] proved $\chi'_s(G) \le 1.6255\,\Delta(G)^2$ for bipartite G with $\Delta(G)$ sufficiently large, $\chi'_s(G) \le 1.6633\,\Delta_A\Delta_B$ when Δ_B/Δ_A is a fixed rational number and Δ_A is sufficiently large, and Conjecture 2 holds asymptotically almost surely for random bipartite graphs whose two parts have comparable sizes. Before [6, 12], no general improvement over the bounds for arbitrary graphs was known in the bipartite setting [6].

Let $\mathcal{M}(G)$ be the set of induced matchings of G. The *fractional strong chromatic index* of G is

$$\chi'_{s,f}(G) = \min\Big\{ \sum_{M\in\mathcal{M}(G)} w_M \;:\; w_M \ge 0 \text{ for all } M,\ \sum_{M\in\mathcal{M}(G),\, M\ni e} w_M \ge 1 \text{ for every } e \in E(G)\Big\}.$$

Requiring $w_M \in \{0,1\}$ gives $\chi'_s(G)$, so $\chi'_{s,f}(G)$ is the optimum of the linear programming relaxation of this integer program, and $\chi'_{s,f}(G) \le \chi'_s(G)$. Equivalently, $\chi'_{s,f}(G)$ is the fractional chromatic number of $L(G)^2$, whose independent sets are exactly the induced matchings of G; see Śleszyńska-Nowak [18]. Cames van Batenburg, Kang and Pirot [5] showed that $\chi'_{s,f}(G) \le 1.5\,\Delta(G)^2$ for bipartite G; Dębski [7] improved it to $1.4762\Delta^2$.

Our first result is the fractional version of Conjecture 2.

Theorem 1.1. *Every bipartite graph G with parts A and B satisfies $\chi'_{s,f}(G) \le \Delta_A\Delta_B$.*

Since an induced matching in the complete bipartite graph K_{Δ_A,Δ_B} contains exactly one edge, $\chi'_{s,f}(K_{\Delta_A,\Delta_B}) = \Delta_A\Delta_B$ and Theorem 1.1 is tight. Our second result is an asymptotic version of Conjecture 2 with an explicit error term.

Theorem 1.2. *Let G be a bipartite graph with parts A and B and $\Delta = \max\{\Delta_A, \Delta_B\}$. Then*

$$\chi'_s(G) \le (1 + o(1))\,\Delta_A\Delta_B,$$

where $o(1)$ is a quantity depending only on Δ that tends to 0 as $\Delta \to \infty$. More precisely,

$$\chi'_s(G) \le \Big(1 + 12\,\frac{(\ln\Delta)^{1/2}}{\Delta^{1/4}}\Big)\Delta_A\Delta_B.$$

The main ingredient in our proofs is a simple coloring lemma based on linear orders of one part of the bipartition. Given a linear order of A, let each vertex $b \in B$ select the edge joining it to its first neighbor in that order. Lemma 2.3 shows that the selected edges can be strongly colored with at most Δ_A colors. Choosing the order at random then yields the fractional bound in Theorem 1.1.

For the ordinary strong chromatic index, we use several linear orders of A and their reverses. These orders produce a collection of columns to which the edges are assigned. Lemma 2.3 reduces the problem to balancing the number of edges assigned to each column around each vertex of A. We obtain the required balance using the Lovász Local Lemma, which leads to Theorem 1.2.

The paper is organized as follows. Section 2 collects the probabilistic tools and proves the column lemma; both are key ingredients in our proof. The proof of Theorem 1.1 is presented in Section 3. Finally, in Section 4, we first construct balanced families of orders and develop the randomized rounding argument, and then use them to prove Theorem 1.2.

2. Preliminaries

For a vertex v we write $N(v)$ for its neighborhood and $d(v) = |N(v)|$ for its degree; ln is the natural logarithm and $[n] = \{1, \dots, n\}$. Throughout, G is a bipartite graph with parts A and B;

we write each edge as xb with $x \in A$ and $b \in B$. Vertices of degree 0 play no role, and we assume that every vertex of B has degree at least 1.

Two distinct edges of a graph *conflict* if they share an endpoint or are joined by an edge. Since G is bipartite, two distinct edges xb and $x'b'$, where $x, x' \in A$ and $b, b' \in B$, conflict if and only if $x = x'$, $b = b'$, $xb' \in E(G)$, or $x'b \in E(G)$. Hence a set of edges is an induced matching if and only if no two of its edges conflict, and a strong edge-coloring is a coloring of $E(G)$ in which conflicting edges receive distinct colors.

We use two probabilistic tools. The first is a form of the Chernoff and Bernstein inequalities, see [15, Theorems 2.1 and 2.8].

Lemma 2.1. *Let S be a sum of finitely many independent random variables with values in $\{0,1\}$, and let μ be the expected value of S. Then for every $t \geq 0$,*

$$\Pr(S \geq \mu + t) \leq \exp\Big(-\frac{t^2}{2(\mu + t/3)}\Big) \quad \textit{and} \quad \Pr(S \leq \mu - t) \leq \exp\Big(-\frac{t^2}{2\mu}\Big).$$

The second is the symmetric Lovász Local Lemma, see [1, Corollary 5.1.2].

Lemma 2.2 (Lovász Local Lemma)**.** *Let $\mathcal{O}_1, \ldots, \mathcal{O}_n$ be events, each having probability at most p. Suppose that each $\mathcal{O}_i$ is mutually independent of all but at most d of the other events. If $e \cdot p \cdot (d+1) \leq 1$, then $\Pr\left(\bigcap_{i=1}^n \overline{\mathcal{O}_i}\right) > 0$. In particular, with positive probability none of the events $\mathcal{O}_1, \ldots, \mathcal{O}_n$ occurs.*

For our purposes, the underlying probability space is a product of finitely many independent random variables, and each event $\mathcal{O}_i$ is determined by a set S_i of these variables. In this situation $\mathcal{O}_i$ is mutually independent of the family of all events $\mathcal{O}_j$ with $S_j \cap S_i = \emptyset$, which is how the hypothesis of Lemma 2.2 will be verified.

Let $K > 0$ be an integer and let $\sigma_1, \ldots, \sigma_K$ be linear orders of A. For each $i \in [K]$, associate two columns, denoted by $(i,-)$ and $(i,+)$, corresponding to rules to assign the first and last neighbors of $b \in B$ in the order σ_i, respectively. Let $J = \{(i,-), (i,+) : i \in [K]\}$, and for each $b \in B$ and $j \in J$, the *center* $c_j(b) \in N(b)$ is defined by

$$c_j(b) = \begin{cases} \min_{\sigma_i} N(b), & j = (i,-) \\ \max_{\sigma_i} N(b), & j = (i,+). \end{cases}$$

For $y \in N(b)$, let $M_y(b) = \{j \in J : c_j(b) = y\}$ be the set of columns in which y is the center of b, and let $m_y(b) = |M_y(b)|$. For fixed b, $M_y(b) \cap M_{y'}(b) = \emptyset$, where $y \neq y' \in N(b)$, and $\cup_{y \in N(b)} M_y(b) = J$. If $d(b) \geq 2$, then $m_y(b)$ is the number of indices i for which y is the σ_i-smallest or the σ_i-largest vertex of $N(b)$; if $d(b) = 1$, then $m_y(b) = 2K$.

Lemma 2.3 (Column lemma)**.** *Let G be a bipartite graph with parts A and B. Let $V_j \subseteq B$ for $j \in J$, and let $E_j = \{c_j(b)b : b \in V_j\}$.*

(a) *For every $j \in J$, the edge set E_j can be partitioned into at most $\max_{x \in A} |N(x) \cap V_j|$ induced matchings of G.*

(b) *If every edge of G lies in at least one of the sets E_j, then*

$$\chi'_s(G) \leq \sum_{j \in J} \max_{x \in A} |N(x) \cap V_j|.$$

Proof. (a) Let $j = (i,-)$; for $j = (i,+)$ replace σ_i by its reverse in the following argument. Put $\lambda = \max_{x \in A} |N(x) \cap V_j|$. Every $b \in V_j$ contributes exactly one edge to E_j, namely $c_j(b)b$. Color the edges of E_j one by one, processing their centers in increasing σ_i-order and the edges with a common center in any order, and give each edge a color in $[\lambda]$ that is not used on an already colored edge conflicting with it.



FIGURE 1. The order σ of A is $x_1 < x_2 < x_3 < x_4 < x_5$. Each b_i keeps the edge to its σ-first neighbor (thick edges). Lemma 2.3(a), with one column and $V_j = B$, guarantees a strong coloring of the thick edges with $\max_x d(x) = 3$ colors; here the procedure of its proof, always taking the smallest available color, uses two: solid edges x_1b_1, x_2b_2 and dashed edges x_1b_4, x_3b_3. Note that x_3b_3 and x_1b_1 conflict (via the edge x_3b_1), as do x_3b_3 and x_2b_2 (via x_3b_2), while x_3b_3 and x_1b_4 do not.

We show that such a color always exists. Let $xb \in E_j$, so $x = c_j(b)$, and let $x'b' \in E_j$ be an already colored edge that conflicts with xb. Then $b' \neq b$, since b contributes only one edge to E_j, and $x' \leq_{\sigma_i} x$, since centers are processed in increasing order. If $x' \neq x$ and $x'b \in E(G)$, then x' is a neighbor of b with $x' <_{\sigma_i} x = \min_{\sigma_i} N(b)$, which is impossible. As $b \neq b'$, the definition of conflict leaves two possibilities: $x' = x$, or $xb' \in E(G)$. In both cases $xb' \in E(G)$, and hence, $b' \in N(x)$. Therefore, every already colored edge conflicting with xb is of the form $c_j(b')b'$ with $b' \in (N(x) \cap V_j) \setminus \{b\}$. There are at most $\lambda - 1$ such edges, so a color in $[\lambda]$ is available. In the resulting coloring no two conflicting edges have the same color, so every color class is an induced matching of G.

(b) Assign each edge to one of the sets E_j containing it. Color the edges assigned to E_j as in (a), using a palette of $\max_x |N(x) \cap V_j|$ colors, with pairwise disjoint palettes for different columns. Since distinct columns use disjoint palettes, any two edges receiving the same color belong to the same E_j, and hence do not conflict by part (a). □

Remark 2.4. The choice of the first neighbor is used exactly once in the proof: it excludes conflicts through an edge $x'b$ with x' preceding x. Figure 1 shows a small example. The bound in Lemma 2.3(a) is attained, for instance, by a complete bipartite graph with a single column and $V_j = B$.

3. PROOF OF THEOREM 1.1: THE FRACTIONAL BOUND

The proof of Theorem 1.1 has two parts. First, a fractional strong edge-coloring can be obtained from a suitable probability distribution on the induced matchings of G (Lemma 3.1). Second, a single random order of A produces a random induced matching that contains every edge with probability at least $1/(\Delta_A\Delta_B)$.

Lemma 3.1. *Suppose there exists a probability distribution on the induced matchings of G such that a random induced matching M drawn from this distribution satisfies $\Pr(e \in M) \geq p$ for every edge $e \in E(G)$. Then $\chi'_{s,f}(G) \leq \frac{1}{p}$.*

Proof. For every induced matching M' of G put $w_{M'} = \Pr(M = M')/p$. These weights are nonnegative, and since the probabilities $\Pr(M = M')$ add up to 1, the weights add up to $1/p$. For every edge e,

$$\sum_{e \in M'} w_{M'} = \frac{1}{p} \sum_{e \in M'} \Pr(M = M') = \frac{\Pr(e \in M)}{p} \geq 1.$$

So the weights $w_{M'}$ are admissible in the definition of $\chi'_{s,f}(G)$, and $\chi'_{s,f}(G) \leq \sum_{M'} w_{M'} = 1/p$. □

Proof of Theorem 1.1. By Lemma 3.1, it suffices to construct a probability distribution on the induced matchings of G such that a random induced matching M drawn from this distribution satisfies

$$\Pr(e \in M) \geq \frac{1}{\Delta_A \Delta_B}$$

for every edge $e \in E(G)$. We construct such a distribution in four steps.

Step 1: a random order. Let σ be a linear order of A chosen uniformly at random from all linear orders of A.

Step 2: every vertex of B keeps one edge. Every $b \in B$ keeps the edge that joins b to its σ-first neighbor. Let

$$E_\sigma = \big\{(\min\nolimits_\sigma N(b))\, b : \ b \in B\big\}$$

be the set of kept edges; it contains exactly one edge at each vertex of B.

Step 3: splitting the kept edges. Apply Lemma 2.3(a) with $K = 1$, $\sigma_1 = \sigma$, the column $j = (1, -)$ and $V_j = B$. Then $E_j = E_\sigma$ and $\max_{x \in A} |N(x) \cap V_j| = \max_{x \in A} d(x) = \Delta_A$. Hence E_σ can be partitioned into induced matchings $M_{\sigma,1}, \ldots, M_{\sigma,\Delta_A}$ of G, some of which may be empty. We fix such a partition for every linear order σ of A.

Step 4: choosing one part. Choose $t \in [\Delta_A]$ uniformly at random and independently of σ, and put $M = M_{\sigma,t}$. Then M is an induced matching of G.

Now fix an edge yb with $y \in A$ and $b \in B$. The edge yb belongs to M exactly when the following two events occur:

(i) $yb \in E_\sigma$, that is, y is the σ-first vertex of $N(b)$;
(ii) t is the index of the part of the partition of E_σ that contains yb.

The order that σ induces on $N(b)$ is uniformly random, so each of the $d(b)$ vertices of $N(b)$ is the σ-first one with the same probability, and (i) has probability $1/d(b)$. If (i) occurs, then yb lies in exactly one of the parts $M_{\sigma,1}, \ldots, M_{\sigma,\Delta_A}$, and since t is uniform on $[\Delta_A]$ and independent of σ, the conditional probability of (ii) is $1/\Delta_A$. Therefore

$$\Pr(yb \in M) = \frac{1}{d(b)} \cdot \frac{1}{\Delta_A} \geq \frac{1}{\Delta_A \Delta_B},$$

since $d(b) \leq \Delta_B$. Lemma 3.1 with $p = 1/(\Delta_A \Delta_B)$ gives $\chi'_{s,f}(G) \leq \Delta_A \Delta_B$. □

4. The proof of Theorem 1.2

First, suppose that $k \geq 2$ and $\Delta \geq 1$ are integers, and G is a bipartite graph with parts A and B such that $d(x) \leq \Delta$ for every $x \in A$ and $1 \leq d(b) \leq k$ for every $b \in B$. A vertex $y \in N(b)$ is the center of b in exactly $m_y(b)$ of the $2K$ columns. On average this number is $2K/d(b) \geq 2K/k$; the next lemma shows that all of them can be made close to this value simultaneously.

Lemma 4.1. *Let $\delta \in (0, 1)$ and let K be an integer with*

$$K \geq \frac{k}{\delta^2} \ln(3k^2\Delta).$$

There are linear orders $\sigma_1, \ldots, \sigma_K$ of A such that

$$m_y(b) \geq (1 - \delta)\, \frac{2K}{k} \qquad \text{for every } b \in B \text{ and every } y \in N(b).$$

Proof. If $d(b) = 1$, then $m_y(b) = 2K$. If $d(b) = 2$, then in every order one vertex of $N(b)$ is the smallest and the other is the largest, so $m_y(b) = K \geq 2K/k$. These vertices b satisfy the conclusion for any orders.

Choose independent random labels $\ell_i(x)$, uniform in $[0, 1]$, for $i \in [K]$ and $x \in A$, and let σ_i order A by increasing ℓ_i; ties occur with probability 0. Let $b \in B$ with $d = d(b) \geq 3$ and let $y \in N(b)$.

For each i, the events "y is the σ_i-smallest vertex of $N(b)$" and "y is the σ_i-largest vertex of $N(b)$" are disjoint with probability $1/d$ each, and those events with different is are independent. So $m_y(b)$ is a sum of K independent indicator variables with mean $\mu = 2K/d \geq 2K/k$. By Lemma 2.1 with $t = \delta\mu$,

$$\Pr\Big(m_y(b) < (1-\delta)\frac{2K}{k}\Big) \leq \Pr\big(m_y(b) \leq (1-\delta)\mu\big) \leq \exp\Big(-\frac{\delta^2\mu}{2}\Big) \leq \exp\Big(-\frac{\delta^2 K}{k}\Big).$$

Let $\mathcal{D}_b$ be the event that $m_y(b) < (1-\delta)2K/k$ for some $y \in N(b)$. Then $\Pr(\mathcal{D}_b) \leq k\exp(-\delta^2 K/k) =: p$. The event $\mathcal{D}_b$ is determined by the labels of the vertices in $N(b)$, so it is mutually independent of the family of events $\mathcal{D}_{b'}$ with $N(b')\cap N(b) = \emptyset$. The number of vertices $b' \neq b$ with $N(b')\cap N(b) \neq \emptyset$ is at most $\sum_{x\in N(b)}(d(x)-1) \leq k(\Delta-1)$. Since $K \geq (k/\delta^2)\ln(3k^2\Delta)$,

$$e\cdot p\cdot\big(k(\Delta-1)+1\big) \leq e\cdot k\cdot k\Delta\cdot\exp\Big(-\frac{\delta^2 K}{k}\Big) \leq \frac{e}{3} < 1.$$

By Lemma 2.2, with positive probability none of the events $\mathcal{D}_b$ occurs. □

The next result may be regarded as a version of Theorem 1.2 with free parameters.

Theorem 4.2. *Let $\delta \in (0,1)$, let K be an integer with $K \geq (k/\delta^2)\ln(3k^2\Delta)$, and put $L = \ln(9Kk\Delta)$. Then*

$$\chi'_s(G) \leq \frac{k\Delta}{1-\delta} + 2\sqrt{\frac{Kk\Delta L}{1-\delta}} + 2KL.$$

Proof. Fix orders $\sigma_1,\dots,\sigma_K$ as in Lemma 4.1. For every $b \in B$ and every $y \in N(b)$, $m_y(b) \geq (1-\delta)\frac{2K}{k} > 0$ by Lemma 4.1. Hence $M_y(b) \neq \emptyset$. Choose a column $j_{b,y} \in M_y(b)$ uniformly at random and all choices are independent. For $j \in J$ put

$$V_j = \{b \in B : j_{b,c_j(b)} = j\},$$

and let $E_j = \{c_j(b)b : b \in V_j\}$ as in Lemma 2.3. Since $j_{b,y} \in M_y(b)$, every edge yb lies in E_j with $j = j_{b,y}$.

Fix $x \in A$ and $j \in J$, and let

$$X_{x,j} = |N(x)\cap V_j| = \sum_{b\in N(x)} \mathbf{1}\big[j_{b,c_j(b)} = j\big].$$

The summand corresponding to b depends only on the choice $j_{b,c_j(b)}$, and the choices corresponding to different vertices $b \in N(x)$ are made independently. Since $j \in M_{c_j(b)}(b)$, the summand of b equals 1 with probability $1/m_{c_j(b)}(b)$, which is at most $k/(2(1-\delta)K)$ by Lemma 4.1 (and equals $1/(2K)$ if $d(b) = 1$). As $d(x) \leq \Delta$, the expected value $\mu_{x,j}$ of $X_{x,j}$ is at most

$$\lambda := \frac{k\Delta}{2(1-\delta)K}.$$

If $\mu_{x,j} < \lambda$, add to $X_{x,j}$ the sum of $r = \lceil\lambda - \mu_{x,j}\rceil$ indicator variables, each with mean $(\lambda-\mu_{x,j})/r$, independent of each other and of all choices $j_{b,y}$. The result $X'_{x,j}$ is a sum of independent indicators with $X'_{x,j} \geq X_{x,j}$ and expected value exactly λ; if $\mu_{x,j} = \lambda$, put $X'_{x,j} = X_{x,j}$. Let $t = \sqrt{2\lambda L} + L$. Then

$$t^2 - 2\lambda L - \tfrac{2}{3}Lt = \tfrac{4}{3}L\sqrt{2\lambda L} + \tfrac{1}{3}L^2 \geq 0,$$

that is, $t^2/(2(\lambda + t/3)) \geq L$. By Lemma 2.1,

$$\Pr(X_{x,j} \geq \lambda + t) \leq \Pr(X'_{x,j} \geq \lambda + t) \leq \exp(-L) = \frac{1}{9Kk\Delta}.$$

Let $\mathcal{O}_{x,j}$ be the event $X_{x,j} \geq \lambda + t$. It is determined by the choices $j_{b,y}$ with $b \in N(x)$, so it is mutually independent of the family of events $\mathcal{O}_{x',j'}$ with $N(x')\cap N(x) = \emptyset$. The number of vertices

x' with $N(x') \cap N(x) \neq \emptyset$, including x itself, is at most $1 + \sum_{b \in N(x)} (d(b) - 1) \leq 1 + \Delta(k-1) \leq k\Delta$; hence the number of pairs (x', j') with $N(x') \cap N(x) \neq \emptyset$, including (x, j), is at most $2Kk\Delta$. Since

$$e \cdot \frac{1}{9Kk\Delta} \cdot 2Kk\Delta = \frac{2e}{9} < 1,$$

Lemma 2.2 shows that for some choice of the columns $j_{b,y}$ we have $X_{x,j} < \lambda + t$ for all $x \in A$ and $j \in J$. For this choice, Lemma 2.3(b) gives

$$\chi'_s(G) \leq \sum_{j \in J} \max_{x \in A} X_{x,j} \leq 2K(\lambda + t) = \frac{k\Delta}{1-\delta} + 2K\sqrt{2\lambda L} + 2KL,$$

and $2K\sqrt{2\lambda L} = 2\sqrt{Kk\Delta L/(1-\delta)}$. □

Now we are ready to prove Theorem 1.2

Proof of Theorem 1.2. Let

$$u = \frac{(\ln \Delta)^{1/2}}{\Delta^{1/4}}, \qquad \text{so that} \qquad u^2 = \frac{\ln \Delta}{\Delta^{1/2}}.$$

Choose Δ large enough so that $u < 1/10$. Put

$$\delta = u, \qquad K = \left\lceil \frac{k}{u^2} \ln(3k^2\Delta) \right\rceil, \qquad L = \ln(9Kk\Delta).$$

Then $\delta \in (0,1)$ and K satisfies the hypothesis of Theorem 4.2. Dividing the bound of Theorem 4.2 by $k\Delta = \Delta_A \Delta_B$ gives

$$\frac{\chi'_s(G)}{\Delta_A \Delta_B} \leq \frac{1}{1-u} + 2\sqrt{\frac{1}{1-u} \cdot \frac{KL}{k\Delta}} + 2\,\frac{KL}{k\Delta}. \tag{1}$$

We estimate K and L. Since $k \leq \Delta$, we have $\ln(3k^2\Delta) \leq \ln 3 + 3\ln\Delta \leq 4\ln\Delta$. So

$$K = \left\lceil \frac{k}{u^2} \ln(3k^2\Delta) \right\rceil \leq \left\lceil \frac{4k \ln \Delta}{u^2} \right\rceil = \left\lceil 4k\Delta^{1/2} \right\rceil \leq 4k\Delta^{1/2} + 1.$$

Hence $L = \ln(9Kk\Delta) \leq \ln(9 \cdot 4.001 \cdot \Delta^{7/2}) \leq \ln 37 + 7/2 \ln \Delta$. Therefore,

$$\frac{KL}{k\Delta} \leq \frac{(4k\Delta^{1/2} + 1) \cdot (\ln 37 + 7/2 \ln \Delta)}{k\Delta} \leq 4 \cdot 3.6\, \frac{\ln \Delta}{\Delta^{1/2}} = 14.4u^2 \leq 1.44\, u.$$

Finally, $u \leq 1/10$ gives $\frac{1}{1-u} = 1 + \frac{u}{1-u} \leq 1 + \frac{10}{9}u$ and $\frac{1}{1-u} \leq \frac{10}{9}$. So the three terms on the right-hand side of (1) satisfy

$$\frac{1}{1-u} \leq 1 + \frac{10}{9}u, \qquad 2\sqrt{\frac{1}{1-u} \cdot \frac{KL}{k\Delta}} \leq 2\sqrt{\frac{10}{9} \cdot 14.4u^2} = 8\,u, \qquad 2\,\frac{KL}{k\Delta} \leq 2.88\,u.$$

Adding them,

$$\frac{\chi'_s(G)}{\Delta_A \Delta_B} \leq 1 + \Big(\frac{10}{9} + 8 + 2.88\Big)u < 1 + 12u.$$ □

Remark 4.3. The choice of the side A in the proof matters: the parameters L_0 and L grow with $\ln k$, and the error term is small because $k \leq \Delta$. The proof also shows where the error term comes from. The factor $1/(1-\delta)$ is the price for the imbalance of the counts $m_y(b)$ among K random orders, which forces K to be of order $k \log \Delta/\delta^2$; the term $2\sqrt{Kk\Delta L/(1-\delta)}$ is the price for the fluctuations of the loads $X_{x,j}$, whose expected values λ are of order $k\Delta/K$. Balancing the two gives the relative error of order $(\log \Delta)^{1/2}\Delta^{-1/4}$. We could improve 12 in Theorem 1.2 to a smaller constant if we are more careful in calculation.

AI usage declaration

Claude Opus 5.5 was used to assist in developing the proofs and in preparing the text of this article. The main ideas are due to the authors, who have verified all mathematical content and take full responsibility for its correctness.

References

[1] N. Alon and J. H. Spencer, *The Probabilistic Method*, 4th ed., Wiley, Hoboken, NJ, 2016.

[2] M. Bonamy, T. Perrett and L. Postle, Colouring graphs with sparse neighbourhoods: bounds and applications, *J. Combin. Theory Ser. B* **155** (2022), 278–317.

[3] R. A. Brualdi and J. J. Quinn Massey, Incidence and strong edge colorings of graphs, *Discrete Math.* **122** (1993), 51–58.

[4] H. Bruhn and F. Joos, A stronger bound for the strong chromatic index, *Combin. Probab. Comput.* **27** (2018), 21–43.

[5] W. Cames van Batenburg, R. J. Kang and F. Pirot, Strong cliques and forbidden cycles, *Indag. Math.* **31** (2020), 64–82.

[6] E. Davey, E. Hurley, R. de Joannis de Verclos, R. J. Kang and J. Volec, Strong edge-colouring via local flag algebras, arXiv:2607.17421, 2026.

[7] M. Dębski, Fractional strong chromatic index of bipartite graphs, *Discrete Math.* (2017), doi:10.1016/j.disc.2017.02.006.

[8] P. Erdős and J. Nešetřil, Problem, in: G. Halász and V. T. Sós (eds.), *Irregularities of Partitions*, Springer, Berlin, 1989, 162–163.

[9] R. J. Faudree, A. Gyárfás, R. H. Schelp and Zs. Tuza, The strong chromatic index of graphs, *Ars Combin.* **29B** (1990), 205–211.

[10] R. J. Faudree, A. Gyárfás, R. H. Schelp and Zs. Tuza, Induced matchings in bipartite graphs, *Discrete Math.* **78** (1989), 83–87.

[11] J.-L. Fouquet and J.-L. Jolivet, Strong edge-colorings of graphs and applications to multi-k-gons, *Ars Combin.* **16A** (1983), 141–150.

[12] Y. Hao, T. Yang and X. Yu, Strong chromatic index of bipartite graphs, arXiv:2606.23824, 2026.

[13] M. Huang, G. Yu and X. Zhou, The strong chromatic index of $(3, \Delta)$-bipartite graphs, *Discrete Math.* **340** (2017), 1143–1149.

[14] E. Hurley, R. de Joannis de Verclos and R. J. Kang, An improved procedure for colouring graphs of bounded local density, *Adv. Comb.* (2022), Paper No. 7.

[15] S. Janson, T. Łuczak and A. Ruciński, *Random Graphs*, Wiley, New York, 2000.

[16] M. Molloy and B. Reed, A bound on the strong chromatic index of a graph, *J. Combin. Theory Ser. B* **69** (1997), 103–109.

[17] K. Nakprasit, A note on the strong chromatic index of bipartite graphs, *Discrete Math.* **308** (2008), 3726–3728.

[18] M. Śleszyńska-Nowak, Clique number of the square of a line graph, *Discrete Math.* **339** (2016), 1551–1556.

[19] A. Steger and M.-L. Yu, On induced matchings, *Discrete Math.* **120** (1993), 291–295.