

Functional and differential equations

– 2

Functional-differential equations

Homework

13 сентября 2026 года

Theoretical minimum. These equations connect f and f' at different points, and the first move is always to gain regularity: an integral over a moving window gives one derivative, and iterating gives all of them. A mean value equation is solved by fixing one argument and differentiating in the other, which converts it into an ordinary differential equation; the answer depends on the mean, and for the arithmetic mean it is a quadratic polynomial. A delay equation is solved by $e^{\lambda x}$, but the characteristic equation is transcendental and has infinitely many roots, so “all solutions” is a statement about a series, not a finite basis. Equations with a composition are rigid: one compares monotonicity and growth of the two sides and usually gets a contradiction. Finally, a differentiable solution of a functional equation may be found by differentiating in each variable separately and eliminating.

1. (RMO) Find all differentiable functions $f: [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$ and $f'(x^2) = f(x)$ for all $x \geq 0$.
2. (RMO 2015) Find all differentiable $f: \mathbb{R} \rightarrow \mathbb{R}$ such that $f'(a) = 0$ for every integer a and $f(x) = 0$ implies $f'(x) = 0$.
3. Prove that there is no differentiable $f: \mathbb{R} \rightarrow \mathbb{R}$ with $f'(x) = f(f(x))$ and $f(0) > 0$, and compare with the IMC problem on $f'(t) > f(f(t))$ in the homework of Differentiable functions – 1.
4. Find all differentiable f with $f'(x) = f(x+1)$ for all x ; describe the solutions of exponential type and show that the characteristic equation $\lambda = e^\lambda$ has no real roots.

5. Find all continuous $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $f(x) = \frac{1}{2h} \int_{x-h}^{x+h} f$ for a fixed $h > 0$ and all x , and compare the answer with the mean value property of harmonic functions (Complex analysis – 8).
6. Find all differentiable $f: \mathbb{R} \rightarrow \mathbb{R}$ with $f(x+y) = f(x) + f(y) + f'(x)f'(y)$ for all x, y .
7. Let f be differentiable with $f(a) - f(b) = (a-b)f'(\frac{2ab}{a+b})$ for all positive a, b (the harmonic mean). Find all such f , and explain the general principle: for which means μ does the equation have non-polynomial solutions?
8. Find all C^1 functions with $f'(x) = f(1-x)$ for all $x \in \mathbb{R}$, and all C^2 functions with $f''(x) = f(-x)$.
9. Let f be continuous on $\mathbb{R}$ and satisfy $f(x) = f(2x)$ for all x . Prove that f is constant; then find all differentiable f with $f'(x) = f(2x)$ and $f(0) = 1$, as a power series, and determine its radius of convergence.
10. Prove that a solution of $f'(x) = f(x-1)$ which is bounded on $\mathbb{R}$ and tends to zero at $-\infty$ is identically zero, and explain which property of the characteristic roots is responsible.