arXiv:2608.14998v1 [math.CA] 15 Aug 2026

ARITHMETIC-PROGRESSION GAP SETS IN CANTOR SETS

S. SANDBERG-CLARK, K. TAYLOR, AND A. YAVICOLI

Abstract. We address the question of which common differences can arise in arithmetic progressions contained in fractal sets. For a compact set $C \subset \mathbb{R}$, we investigate not only whether arithmetic progressions occur in C, but the full collection of their common differences. More generally, for a finite pattern P, we study the set of scales at which affine copies of P appear in C.

For affine self-similar sets satisfying strong separation, we obtain explicit restrictions on admissible common differences. Specializing to middle-ε Cantor sets, we prove that the longest arithmetic progression has length four whenever $3-2\sqrt{2} < \varepsilon \leq 1/3$, showing that the maximal progression length drops immediately from six at the critical parameter $\varepsilon = 3-2\sqrt{2}$. We further develop recursive bounds for the sets of admissible common differences and derive explicit blackout intervals, namely ranges of scales for which arithmetic progressions cannot occur.

On the positive side, sufficiently thick Cantor sets exhibit the opposite behavior. Combining a refinement of the Hunt-Kan-Yorke construction with the Newhouse Gap Lemma, we prove that every sufficiently small common difference occurs in a three-term arithmetic progression. In particular, if the largest bounded gap of a Cantor set is at most $0.067\,\mathrm{diam}(C)$ and its thickness is at least $6.96268\ldots$, then every common difference in $(0, 0.435\,\mathrm{diam}(C)]$ occurs in a three-term arithmetic progression contained in C. Analogous interval results are obtained for four-term arithmetic progressions and asymmetric three-point patterns.

The occurrence of finite configurations in large or structured sets is a central theme in additive combinatorics and fractal geometry. In the integer setting, largeness is often expressed through density. For compact subsets of Euclidean space, Hausdorff dimension, Fourier decay, and Newhouse thickness provide different quantitative notions of size. The present paper asks not only whether a compact set contains a prescribed configuration, but also at which scales that configuration occurs.

For a compact set $C \subset \mathbb{R}$ and an integer $k \geq 2$, define the *arithmetic-progression gap set*

$$G_{AP}^k(C) := \big\{t > 0 : \ \exists x \in \mathbb{R} \text{ with } x, x+t, \ldots, x+(k-1)t \in C\big\}, \tag{0.1}$$

where we refer to $t \in G_{AP}^k(C)$ as a *common difference*. More generally, let

$$P = \{0 = p_1 < p_2 < \cdots < p_k = 1\} \subset [0,1]$$

be a normalized finite pattern. Its *scale set* in C is

$$G_P(C) := \{s > 0 : \ \exists x \in \mathbb{R} \text{ such that } x + sP \subset C\}. \tag{0.2}$$

Here s is the diameter of the embedded copy. We write

$$\delta_j(P) = p_j - p_{j-1}, \qquad \delta_{\min}(P) = \min_{2\leq j\leq k} \delta_j(P), \qquad \delta_{\max}(P) = \max_{2\leq j\leq k} \delta_j(P).$$

2020 *Mathematics Subject Classification.* Primary 28A80; Secondary 11B25, 28A78.

Key words and phrases. Cantor sets, arithmetic progressions, Newhouse thickness, pattern scales, Hunt–Kan–Yorke theorem.

For

$$\mathcal{A}_k = \left\{0, \frac{1}{k-1}, \dots, 1\right\},$$

the two conventions satisfy

$$G_{\mathcal{A}_k}(C) = (k-1)G_{AP}^k(C). \tag{0.3}$$

For asymmetric three-point patterns we use

$$P_\theta = \{0, \theta, 1\}, \qquad 0 < \theta < 1.$$

The scale-set problem is an affine-configuration analogue of the distance-set problem. Falconer's conjecture asks whether $\dim_H E > d/2$ forces the distance set $\Delta(E) = \{|x-y| : x, y \in E\}$ to have positive Lebesgue measure [1]; Mattila and Sjölin proved nonempty interior under the stronger condition $\dim_H E > (d+1)/2$ [2]. Arithmetic progressions impose a simultaneous multipoint relation, and dimensional information alone does not determine their existence. Keleti constructed full-dimensional compact subsets of $\mathbb{R}$ avoiding prescribed three-point similarity patterns [3]. Yavicoli strengthened this to positive measure for gauges arbitrarily close to the ambient-dimensional gauge while avoiding countably many linear patterns [4]. Shmerkin constructed full-dimensional Salem sets without nontrivial three-term arithmetic progressions [7]. Conversely, sufficiently strong dimensional and Fourier-decay estimates do force three-term progressions [5]. More recently, Carnovale and Senger studied Lebesgue- and Hausdorff-size conclusions for the set of three-term progression step sizes under Fourier-analytic hypotheses [6]. Our emphasis is complementary: we seek exact or recursive information for specific Cantor sets, explicit blackout intervals, and guaranteed intervals of scales arising from self-similarity and thickness. These results motivate the use of geometric structure, and especially thickness, rather than dimension alone.

Arithmetic progressions in middle Cantor sets and in homogeneous self-similar sets have also been studied directly. Chaika proved that the middle-Nth Cantor set contains arithmetic progressions of length at least a constant multiple of $N/\log_2 N$ [9]. Broderick, Fishman, and Simmons obtained, for sufficiently small ε, quantitative bounds of the form

$$\frac{1/\varepsilon}{\log(1/\varepsilon)} \lesssim L_{AP}(C_\varepsilon) \leq \frac{1}{\varepsilon} + 1,$$

where $L_{AP}(E)$ denotes the maximal length of a nontrivial arithmetic progression in E; they also recorded the Newhouse-Gap-Lemma consequence $L_{AP}(C_\varepsilon) \geq 4$ for $0 < \varepsilon \leq 1/3$ [8]. Xi, Jiang, and Pei investigated existence and maximal-length questions for homogeneous self-similar sets using multiple β-expansions and asked whether $L_{AP}(C_\varepsilon)$ is an increasing staircase function of the contraction ratio [10]. Tian, Lou, and Shang answered that question negatively; in particular, they proved the exact endpoint value

$$L_{AP}(C_{3-2\sqrt{2}}) = 6. \tag{0.4}$$

Thus the strict lower endpoint in our maximal-length-four theorem is essential [11]. In contrast with this earlier maximal-length literature, the main objects here are the entire gap and scale sets $G_{AP}^k(C)$ and $G_P(C)$, including their recursive structure, blackout intervals, and intervals of admissible scales.

The middle-ε Cantor set C_ε is our principal model. It is generated by the iterated function system

$$f_0(x) = \lambda x, \qquad f_1(x) = 1 - \lambda + \lambda x,$$

where $\lambda = (1-\varepsilon)/2$. In the case where $\varepsilon = 1/3$, we get the middle-third Cantor set which illustrates the distinction between distances and progression gaps: $\Delta(C_{1/3}) = [0,1]$, whereas

$$G^3_{AP}(C_{1/3}) = G^4_{AP}(C_{1/3}) = \{3^{-n} : n \in \mathbb{N}\}.$$

Thickness, higher-order intersections, and the interior problem. The interior question can be formulated as a uniform higher-order intersection problem. Indeed,

$$t \in G^k_{AP}(C) \quad \Longleftrightarrow \quad \bigcap_{j=0}^{k-1}(C - jt) \neq \varnothing, \tag{0.5}$$

and, for a normalized pattern $P = \{p_1, \ldots, p_k\}$,

$$s \in G_P(C) \quad \Longleftrightarrow \quad \bigcap_{j=1}^{k}(C - sp_j) \neq \varnothing. \tag{0.6}$$

Thus proving that a scale set has interior requires a single geometric hypothesis to force these intersections for every parameter in an interval, not merely for one exceptional scale.

For two-set intersections, Newhouse thickness provides precisely such a robust mechanism. The Newhouse Gap Lemma [13] implies, for example, that if $\tau(C) \geq 1$, then $C \cap (C+t) \neq \varnothing$ for every $0 < t < \operatorname{diam}(C)$; equivalently, the two-point scale set contains an interval. The same lemma can be used more indirectly to produce an individual three-term arithmetic progression in every compact set of thickness at least one: one separates suitable pieces of the set and intersects the set with an appropriate midpoint set. See [17, Proposition 20] and [14] for this idea and its extensions to general three-point configurations. Stronger thickness hypotheses also force larger prescribed finite patterns [16]. These results, however, are primarily existence statements and do not by themselves give a uniform interval of admissible scales.

The theorem of Hunt, Kan, and Yorke gives a route from two-set to higher-order intersections: under suitable thickness assumptions, the intersection of two interleaved Cantor sets contains a compact subset of positive thickness [12]. For three-term progressions, the natural strategy is to construct

$$K_s \subseteq (C - s) \cap (C + s)$$

with quantitative lower bounds for both $\tau(K_s)$ and $\operatorname{diam}(K_s)$, and then apply the Gap Lemma once more to K_s and C. The qualitative HKY statement is not enough for an explicit interval of parameters, because the second application requires bounds that are uniform in s. In Section 4 we extract such bounds directly from the initial stage of the HKY construction. This yields the explicit three-term interval in Theorem 4.1; a second Gap Lemma argument, applied to the HKY output and a short translate of it, gives the four-term interval in Theorem 4.3.

Main results. The paper develops complementary exclusion and interior results.

(1) (Maximal AP length For $3 - 2\sqrt{2} < \varepsilon \leq 1/3$, the longest arithmetic progression in C_ε has exactly four terms (Theorem 2.4). Together with (0.4), this shows that the maximal length drops from six at the endpoint to four immediately to its right.
(2) (Blackout intervals) For affine self-similar sets satisfying strong separation, a general first-splitting theorem confines $G_P(C)$ to an explicit union of rescaled intervals (Theorem 3.1).

A level-two refinement for C_ε lowers the threshold obtained from the first-level bound and proves that $G^3_{AP}(C_\varepsilon)$ contains no interval $(0, r)$ whenever

$$\frac{4-\sqrt{13}}{3} < \varepsilon \leq \frac{1}{3}.$$

The argument also gives explicit blackout intervals in the additional parameter range (see Theorem 3.9).

(3) (Interior of the 3-AP gap set) An explicit initial-stage lower bound extracted from the Hunt–Kan–Yorke construction implies that, if C has diameter d, largest bounded gap at most $0.067d$, and thickness at least $(1-0.067)/(2\cdot 0.067)$, then

$$(0, 0.435d] \subseteq G^3_{AP}(C)$$

(see Theorem 4.1).

(4) (Asymmetric three-point patterns) Using the same assumptions above, for every $0 < \theta < 1$ we prove both blackout criteria and positive intervals for $G_{P_\theta}(C)$. In particular, if $0 < \varepsilon \leq 0.067$, then

$$(0, 0.87] \subseteq G_{P_\theta}(C)$$

(see Corollary 4.2).

(5) (Interior of the 4-AP gap set) Combining the same two-set construction with a second application of the Newhouse Gap Lemma gives

$$(0, 0.30d] \subseteq G^4_{AP}(C)$$

when $\tau(C) \geq 32.83333\ldots$ (see Theorem 4.3).

The novelty and attribution of these results can be summarized as follows. Prior work on middle and homogeneous self-similar Cantor sets concerns existence and maximal AP length [9, 8, 10, 11]. Our first two contributions instead determine complete gap sets in the middle-ε case and develop recursive outer descriptions of the full step-size set, including new explicit blackout intervals. The exact value four on $3-2\sqrt{2} < \varepsilon \leq 1/3$ complements the endpoint value six in [11]. In the positive-scale results, the underlying two-set construction, its qualitative positive-thickness conclusion, the gap-size estimate used in the thickness calculation, and the three-set mechanism are due to Hunt–Kan–Yorke [12]. What is derived here is an explicit initial-stage lower bound within that construction, together with the numerical three-term, four-term, and asymmetric-pattern consequences. We do not claim to introduce the general problem of step-size sets; the Fourier-analytic literature includes, in particular, [6].

Figure 1 records the main parameter ranges for middle-ε Cantor sets. Section 1 reviews thickness and proves the qualitative four-point existence result used later. Sections 2 and 3 contain the exclusion results. Section 4 develops the quantitative Hunt–Kan–Yorke input and its applications: the three- and four-term interval theorems and the three-point asymmetric patterns.

1. Preliminaries

We review thickness, iterated function systems, and related results, as these notions will be used throughout the paper. For a nonempty bounded set $E \subset \mathbb{R}$, write $|E| = \operatorname{diam}(E)$. A common way to formulate thickness is:



FIGURE 1. Principal conclusions for the middle-ε family. A bar whose left endpoint is 0 records a conclusion valid for every smaller positive ε. The endpoints shown in the positive-interval results are conservative rounded constants; strict and non-strict parameter endpoints are as stated in the theorems.

Definition 1.1. (bridge thickness) Let $C \subset \mathbb{R}$ be a compact set, and let (G_n) be the bounded connected components of the complement, $\mathbb{R} \setminus C$, ordered by nonincreasing length. For $n \in \mathbb{N}$, each bounded gap G_n is removed from a closed parent interval I_n, leaving behind two closed children intervals L_n and R_n called bridges, the left and right parts of $I_n \setminus G_n$, respectively. The *Newhouse thickness* of C is defined by

$$\tau(C) := \inf_{n \in \mathbb{N}} \frac{\min\{|L_n|, |R_n|\}}{|G_n|}.$$

As an alternative to bridges and gaps, Hunt–Kan–Yorke defined chunks; a set P is a *chunk* of C, written $P \propto C$, if $P = C \cap I$ is a nonempty proper set where I is a closed interval and $d(P, C \setminus P) > 0$.

Definition 1.2. (chunk thickness) Given a compact set $C \subset \mathbb{R}$, we define the thickness of C to be

$$\tau(C) = \inf_{P \propto C} \frac{|P|}{d(P, C \setminus P)}$$

provided C has a chunk. Otherwise, let $\tau(C) = 0$ if C is empty or consists of a single point, and $\tau(C) = \infty$ if C is an interval with positive length.

Note that these definitions of thickness are equivalent; see [12] for details.

We often want to know when two sets intersect; a precursor to which is being interleaved. Two compact sets are *interleaved* if each meets the interior of the convex hull of the other. Knowing two sets are interleaved and have sufficiently large thickness is enough to guarantee they intersect:

Proposition 1.3 (Newhouse Gap Lemma [13])**.** *If interleaved compact sets $K, L \subset \mathbb{R}$ satisfy $\tau(K)\tau(L) \geq 1$, then $K \cap L \neq \varnothing$.*

Lemma 1.4 (Short translates are interleaved)**.** *Let $K \subset \mathbb{R}$ be a nonempty compact set. If $0 < u < \operatorname{diam}(K)$, then K and $K + u$ are interleaved.*

Proof. Write $\operatorname{conv}(K) = [a, b]$. Since $a, b \in K$ and $b - a > u$,

$$b \in K \cap (a + u, b + u) \quad \text{and} \quad a + u \in (K + u) \cap (a, b).$$

Thus each set meets the interior of the convex hull of the other. □

Consequently, if C is compact, $d = \operatorname{diam}(C) > 0$, and $\tau(C) \geq 1$, then Proposition 1.3 applied to C and $C + t$ gives $d \subseteq G^2_{AP}(C)$. A related Gap Lemma argument shows that every compact set of thickness at least one contains every prescribed nondegenerate three-point configuration on the line; see [17, Proposition 20] for arithmetic progressions and [14] for the general three-point statement. Quantitative existence results for longer finite patterns under stronger thickness hypotheses are developed in [16]. These are qualitative or existence-oriented results; our focus is the geometry of the full scale set.

The symmetry of C_ε gives a useful four-point existence result.

Proposition 1.5 (Symmetric four-point patterns)**.** *Let $C \subset \mathbb{R}$ be a Cantor set with $\operatorname{conv}(C) = [0, 1]$. Assume that C is symmetric about $1/2$, that $1/2 \notin C$, and that $\tau(C) \geq 1$. If g is the length of the largest bounded gap of C, then for every $u \in [g, 1)$ the set C contains an affine copy of*

$$\{-1, -u, u, 1\}.$$

Equivalently, for

$$Q_u = \left\{0, \frac{1-u}{2}, \frac{1+u}{2}, 1\right\},$$

one has $G_{Q_u}(C) \neq \varnothing$.

Proof. Put

$$K = C - \frac{1}{2}, \qquad L = \frac{1}{u}\left(C - \frac{1}{2}\right).$$

Both sets have thickness $\tau(C)$, and

$$\operatorname{conv}(K) = \left[-\frac{1}{2}, \frac{1}{2}\right] \subseteq \left[-\frac{1}{2u}, \frac{1}{2u}\right] = \operatorname{conv}(L).$$

The set L cannot lie in a bounded gap of K because $|L| = 1/u > 1 = |K|$. If $u > g$, every bounded gap of L has length at most $g/u < 1 = |K|$, so K cannot lie in a bounded gap of L either. Thus K and L are interleaved, and Proposition 1.3 gives $K \cap L \neq \varnothing$. For $z \in K \cap L$, the points $\pm z, \pm uz$ lie in K; since $0 \notin K$, they form the required nondegenerate pattern after translating by $1/2$.

It remains to consider $u = g$. The same argument applies unless K is contained in the closure of a bounded gap of L. In that exceptional case, the gap has length at most $g/u = 1$, while $|K| = 1$, so its closure must equal $[-1/2, 1/2]$. Scaling back shows that the central gap of C has length u. Its two endpoints, together with 0 and 1, form exactly the normalized pattern Q_u. □

Corollary 1.6. *For $0 < \varepsilon < 1$, the set C_ε contains a four-term arithmetic progression if and only if $\varepsilon \leq 1/3$.*

Proof. If $\varepsilon \leq 1/3$, then $\tau(C_\varepsilon) \geq 1$ and the largest gap has length ε. Proposition 1.5, with $u = 1/3$, produces the normalized four-term pattern $\mathcal{A}_4$.

Conversely, suppose that C_ε contains a three-term progression. Pull it back from the smallest construction interval containing it. At the first level, two consecutive points lie in one interval of length λ, so the common difference is at most λ; another consecutive pair crosses the central gap, so it is at least ε. Hence $\varepsilon \leq \lambda$, equivalently $\varepsilon \leq 1/3$. A four-term progression contains a three-term one, which proves the converse. □

These results are due to the strong self-similarity properties of the middle-ε Cantor sets which leads us to consider a more general family of self-similar compact sets constructed using iterated function systems.

Definition 1.7 (IFS)**.** Fix $I \subset \mathbb{R}$. Define the *iterated function system* $\{f_i\}_1^N$ to be a finite family of contractive maps $f_i : I \to I$ with contraction factor $0 < r_i < 1$; i.e., $|f_i(x) - f_i(y)| \le r_i|x - y|$ for all $x, y \in I$, $1 \le i \le N$. The *attractor* of $\{f_i\}$ is the unique nonempty compact set $C \subset I$ satisfying

$$C = \bigcup_{i=1}^{N} f_i(C).$$

For a *word* $w = (w_1, \dots, w_n) \in \{0, 1, \dots, N\}^k$, write $f_w = f_{w_1} \circ \cdots \circ f_{w_k}$, and let $f_\emptyset$ be the identity.

We say an IFS has *strong separation* if the first-generation pieces $f_i(C)$ are pairwise disjoint.

Remark 1.8. In the special case of the middle-ε Cantor set, the set C_ε is the attractor of the functions

$$f_0(x) = \lambda x \quad \text{and} \quad f_1(x) = 1 - \lambda + \lambda x,$$

with domain $[0, 1]$ and $\lambda = (1 - \varepsilon)/2$. Its level-$n$ construction consists of 2^n intervals of length λ^n, and

$$\tau(C_\varepsilon) = \frac{\lambda}{\varepsilon} = \frac{1 - \varepsilon}{2\varepsilon}.$$

The above definition preserves the notion of parent and children intervals mentioned above; for example, I is the parent interval to the children intervals $f_1(I), f_2(I), \dots, f_N(I)$.

2. Longest arithmetic progressions in middle-ε Cantor sets

In this section we prove that, for $\varepsilon \in (3 - 2\sqrt{2}, 1/3]$, the longest arithmetic progressions in C_ε have length exactly four. The key point is a precise description of the possible midpoints of three-term progressions.

2.1. The middle-third Cantor set.

Proposition 2.1. *Let $C_{1/3}$ be the middle-third Cantor set. Then*

$$G^4_{AP}(C_{1/3}) = G^3_{AP}(C_{1/3}) = \left\{ \frac{1}{3^n} : n \in \mathbb{N} \right\},$$

and $G^k_{AP}(C_{1/3}) = \varnothing$ for every $k \ge 5$. Moreover, $G^3_{AP}(C_\varepsilon) = \varnothing$ whenever $\varepsilon > 1/3$.

Proof. For $C_{1/3}$ we have $\lambda = \varepsilon = 1/3$, and

$$\lambda^{n-1} \left\{ 0, \frac{1}{3}, \frac{2}{3}, 1 \right\} \subset C_{1/3} \qquad (n \in \mathbb{N}).$$

Thus $\{3^{-n} : n \in \mathbb{N}\}$ is contained in both $G^3_{AP}(C_{1/3})$ and $G^4_{AP}(C_{1/3})$.

Conversely, let $a < b < c$ be a three-term arithmetic progression in $C_{1/3}$, and let I be the smallest construction interval containing all three points. Write I_0, I_1 for the two children of I. Their lengths and the length of the gap between them are all $|I|/3$. If $a, b \in I_0$ and $c \in I_1$, then

$$b - a \le \frac{|I|}{3} \le c - b.$$

Since the two distances are equal, equality holds throughout, and the common difference is $|I|/3$. The case $a \in I_0$ and $b, c \in I_1$ is symmetric. Thus every common difference is 3^{-n} for some $n \in \mathbb{N}$.

The four endpoints of I_0 and I_1 form a four-term progression with common difference $|I|/3$. A five-term progression cannot occur: after rescaling its smallest containing construction interval to $[0, 1]$, a first-level split would force its common difference to be at least $1/3$, whereas its total span forces it to be at most $1/4$, a contradiction.

Finally, if $\varepsilon > 1/3$, then the first-level gap has length ε and each first-level interval has length $\lambda < \varepsilon$. In any first-level splitting three-term progression, two of the points lie in the same first-level interval, forcing the common difference to be at most λ, while a consecutive pair crossing the central gap forces it to be at least ε. This contradiction proves $G^3_{AP}(C_\varepsilon) = \varnothing$. □

2.2. **Midpoints of three-term progressions.** Define the midpoint set

$$M_{AP}(C) := \{x \in C : \ \exists t > 0 \text{ with } x - t, x, x + t \in C\}.$$

Theorem 2.2 (Midpoints of three-term progressions)**.** *Let $0 < \varepsilon \leq 1/3$. Then*

$$M_{AP}(C_\varepsilon) = \bigcup_{n \in \mathbb{N}_0} \bigcup_{w \in \{0,1\}^n} f_w \left(C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \frac{1+\lambda}{2} \right] \right),$$

where the f_w are as in Remark 1.8. More precisely, the midpoints of the three-term progressions that split between the two first-level intervals are exactly

$$C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \frac{1+\lambda}{2} \right].$$

Proof. Suppose first that $\{x - t, x, x + t\} \subset C_\varepsilon$ splits at the first level. By symmetry, assume that $x \in [0, \lambda]$. Then $x - t \in [0, \lambda]$ and $x + t \in [1 - \lambda, 1]$. Hence

$$1 - \lambda - x \leq t \leq x,$$

so $x \geq (1 - \lambda)/2$. Therefore

$$x \in C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \lambda \right].$$

Reflecting about $1/2$ gives the corresponding right-hand interval, proving necessity.

Conversely, let

$$x \in C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \lambda \right].$$

If $x = (1 - \lambda)/2$, take $t = x$. Then $x - t = 0$ and $x + t = 1 - \lambda$, so the required progression is immediate. We may therefore assume $x > (1 - \lambda)/2$. When $\varepsilon = 1/3$, the interval under consideration contains only $x = 1/3$, already covered by the boundary case; hence we may also assume $\varepsilon < 1/3$, or equivalently $\varepsilon < \lambda$.

Define

$$A_x := x - \big(C_\varepsilon \cap [0, \lambda^2]\big), \qquad B_x := \big(C_\varepsilon \cap [1 - \lambda, 1]\big) - x.$$

These are affine copies of C_ε, with

$$\tau(A_x) = \tau(B_x) = \frac{\lambda}{\varepsilon} > 1,$$

and

$$\operatorname{conv}(A_x) = [x - \lambda^2, x], \qquad \operatorname{conv}(B_x) = [1 - \lambda - x, 1 - x].$$

The strict inequality $x > (1-\lambda)/2$ implies that the two convex hulls overlap nontrivially. The largest bounded gap of A_x has length $\lambda^2\varepsilon < |B_x| = \lambda$, while the largest bounded gap of B_x has length $\lambda\varepsilon < |A_x| = \lambda^2$. Consequently neither set can lie in a bounded gap of the other; the hull overlap excludes the unbounded gaps. Thus A_x and B_x are interleaved, and applying the Gap Lemma gives $A_x \cap B_x \neq \varnothing$.

For $t \in A_x \cap B_x$ we have

$$x - t \in C_\varepsilon \cap [0, \lambda^2], \qquad x + t \in C_\varepsilon \cap [1-\lambda, 1].$$

The two points lie in different first-generation intervals, so $t > 0$. Thus x is the midpoint of a first-level splitting progression. The right-hand half follows by symmetry.

Finally, every three-term progression is contained in a smallest construction interval $f_w([0,1])$, determined by a word w. Pulling back by the corresponding inverse branch f_w^{-1} produces a first-level splitting progression in $[0,1]$, and applying the branch f_w again recovers the original interval $f_w[0,1]$. This gives the stated union over words. □

Lemma 2.3 (Endpoint localization)**.** *Assume* $3 - 2\sqrt{2} < \varepsilon \leq 1/3$. *Suppose*

$$\{x - t, x, x + t\} \subset C_\varepsilon$$

splits at the first level and

$$x \in C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \lambda\right].$$

Then $x - t \in [0, \lambda^2]$.

Proof. The point x belongs to the right child of the left first-level interval, since

$$\frac{1-\lambda}{2} \geq \lambda - \lambda^2.$$

If $x - t$ also belonged to that child, then

$$x - t \in [\lambda - \lambda^2, x),$$

so $t \leq \lambda^2$. Since the progression splits at the first level, $x + t \in [1-\lambda, 1]$, and $x \leq \lambda$ gives $t \geq 1 - 2\lambda = \varepsilon$. Thus $\varepsilon \leq \lambda^2$, contrary to

$$\varepsilon > 3 - 2\sqrt{2} \quad \Longleftrightarrow \quad \varepsilon > \lambda^2.$$

Hence $x - t$ lies in the other second-level child, namely $[0, \lambda^2]$. □

Theorem 2.4 (Sharp four-term threshold)**.** *Let* $\varepsilon \in (3 - 2\sqrt{2}, 1/3]$. *Then the longest arithmetic progression contained in* C_ε *has length exactly four.*

Proof. Corollary 1.6 gives a four-term progression, so it remains to exclude five-term progressions.

Suppose, to the contrary, that

$$a, a + t, a + 2t, a + 3t, a + 4t \in C_\varepsilon$$

for some $t > 0$. By passing to the smallest construction interval containing the progression and applying its inverse branch, we may assume that the progression splits at the first level. Set $x = a + 2t$. The triple

$$\{x - 2t, x, x + 2t\}$$

then splits at the first level. By symmetry and Theorem 2.2, we may assume

$$x \in C_\varepsilon \cap \left[\frac{1-\lambda}{2}, \lambda\right].$$

Lemma 2.3, applied with common difference $2t$, gives

$$x - 2t \in [0, \lambda^2].$$

Also $x + 2t$ lies in the right first-level interval.

There are two cases. If $x + t$ lies in the right first-level interval, then $\{x - t, x, x + t\}$ splits at the first level, so Lemma 2.3 gives $x - t \in [0, \lambda^2]$. Since we already know $x - 2t \in [0, \lambda^2]$, the two consecutive points $x - 2t$ and $x - t$ lie in an interval of length λ^2; hence

$$t = (x - t) - (x - 2t) \leq \lambda^2.$$

On the other hand, $x + t \geq 1 - \lambda$ and $x \leq \lambda$, so $t \geq \varepsilon$, contradicting $\varepsilon > \lambda^2$.

If $x + t$ lies in the left first-level interval, then $\{x, x + t, x + 2t\}$ splits at the first level. Its midpoint $x + t$ lies in the left midpoint interval, and the localization lemma gives $x \in [0, \lambda^2]$. This contradicts

$$x \geq \frac{1-\lambda}{2} > \lambda^2,$$

where the strict inequality holds for every $\lambda < 1/2$.

Thus no five-term progression exists. By the nesting $G_{AP}^{k+1}(C) \subseteq G_{AP}^{k}(C)$, no longer progression exists either, and the maximal length is four. □

3. Blackout intervals for pattern scale sets

This section considers when attractors of iterated function systems contain scale sets

$$G_P(C) = \{s > 0 : \exists x \in \mathbb{R} \text{ such that } x + sP \subset C\},$$

and arithmetic-progression gap sets

$$G_{AP}^{k}(C) = \{t > 0 : \exists x \in \mathbb{R} \text{ with } x, x + t, \ldots, x + (k-1)t \in C\},$$

as in equation (0.2) and (0.1), respectively.

A copy $x + sP$ first splits at level $n + 1$ when it lies completely in one level-n interval but intersects at least two children of that interval.

An open interval disjoint from $G_P(C)$ will be called a *blackout interval*. The preceding notation permits a single first-splitting argument for arithmetic and asymmetric patterns.

Theorem 3.1 (General pattern blackout bound)**.** *Let C be the attractor of an affine IFS with $M \geq 2$,*

$$f_i(x) = r_i x + a_i, \qquad 1 \leq i \leq M, \qquad 0 < r_i < 1,$$

with strong separation, $\operatorname{conv}(C) = [0, 1]$*, and*

$$f_i([0, 1]) = [a_i, b_i], \qquad r_i = b_i - a_i,$$

where

$$0 = a_1 < b_1 < a_2 < b_2 < \cdots < a_M < b_M = 1.$$

Set

$$g_{\min} := \min_{2 \leq j \leq M} (a_j - b_{j-1}).$$

Let $P = \{0 = p_1 < \cdots < p_k = 1\}$ be a normalized k-point pattern, and write $\delta_{\min} = \delta_{\min}(P)$ and $\delta_{\max} = \delta_{\max}(P)$. Then

$$G_P(C) \subseteq \bigcup_{n\in\mathbb{N}_0} \bigcup_{(i_1,\dots,i_n)\in\{1,\dots,M\}^n} (r_{i_1}\cdots r_{i_n})[L_P, U_P],$$

where

$$L_P := \frac{g_{\min}}{\delta_{\max}}$$

and

$$U_P := \begin{cases} \min\left\{1, \dfrac{\sum_{i=1}^M r_i}{\delta_{\min}(k-M)}, \dfrac{\max_{1\le i\le M} r_i}{\delta_{\min}(\lceil k/M\rceil - 1)}\right\}, & k > M, \\ 1, & 2 \le k \le M. \end{cases}$$

Proof. Let D_P be the set of scales $s \in G_P(C)$ for which some copy $x + sP \subset C$ is split at the first level. Every copy either is already split or lies in a single first-level interval. In the latter case, applying the inverse branch produces another copy of P in C. This iteration must terminate after finitely many steps. Indeed, let $r_{\max} = \max_i r_i < 1$. If the original copy of scale $s > 0$ remained in a single construction interval for the first n levels, then that interval would have diameter at most $r_{\max}^n$, and therefore $s \le r_{\max}^n$. This is impossible for every n, since $r_{\max}^n \to 0$. Thus a first split occurs after finitely many inverse branches, and

$$G_P(C) \subseteq \bigcup_{n\in\mathbb{N}_0} \bigcup_{(i_1,\dots,i_n)\in\{1,\dots,M\}^n} (r_{i_1}\cdots r_{i_n})D_P.$$

Take $s \in D_P$. At least one pair of consecutive points of the embedded pattern lies in different first-level intervals. For some$j \in \{2,\dots,k\}$,

$$\delta_j(P)s \ge g_{\min},$$

which implies

$$s \ge \frac{g_{\min}}{\delta_j(P)} \ge \frac{g_{\min}}{\delta_{\max}} = L_P.$$

Since $x + sP \subset [0,1]$ and $\operatorname{diam}(P) = 1$, we also have $s \le 1$.

Assume $k > M$. If $A_i = (x + sP) \cap [a_i, b_i]$, then points of A_i are separated by at least $\delta_{\min} s$, and hence

$$\#A_i \le \left\lfloor \frac{r_i}{\delta_{\min} s} \right\rfloor + 1.$$

Therefore

$$k \le \sum_{i=1}^M \#A_i \le \frac{\sum_{i=1}^M r_i}{\delta_{\min} s} + M,$$

which yields

$$s \le \frac{\sum_{i=1}^M r_i}{\delta_{\min}(k-M)}.$$

Moreover, some first-level interval contains at least $\lceil k/M\rceil$ points. Because an interval contains a consecutive block of the ordered pattern, those points span at least $\lceil k/M\rceil - 1$ consecutive pattern gaps. Thus

$$(\lceil k/M\rceil - 1)\delta_{\min} s \le \max_{1\le i\le M} r_i,$$

which gives the third upper bound. Hence $D_P \subseteq [L_P, U_P]$, and the first-splitting decomposition proves the theorem. □

Corollary 3.2 (Homogeneous IFS)**.** *Under the hypotheses of Theorem 3.1, suppose $r_i = r$ for every i. Then*

$$G_P(C) \subseteq \bigcup_{n\in\mathbb{N}_0} r^n[L_P, U_P].$$

If $L_P > U_P$, then $G_P(C) = \varnothing$. If $rU_P < L_P$, then the intervals $r^n[L_P, U_P]$ are pairwise disjoint and $G_P(C)$ contains no interval of the form $(0, \eta)$.

Proof. Every word of length n has contraction ratio r^n, so the asserted containment is the specialization of Theorem 3.1. If $L_P > U_P$, the first-split scale set is empty and hence so is $G_P(C)$. If $rU_P < L_P$, then for every $n \geq 0$,

$$r^{n+1}U_P < r^n L_P,$$

so consecutive containing intervals are separated by the nonempty open gap $(r^{n+1}U_P, r^n L_P)$. These gaps accumulate at zero, which rules out an interval $(0, \eta)$ in $G_P(C)$. □

Corollary 3.3 (Arithmetic progressions)**.** *Under the IFS hypotheses of Theorem 3.1, for $k \geq 3$,*

$$G^k_{AP}(C) \subseteq \bigcup_{n\in\mathbb{N}_0} \bigcup_{(i_1,\ldots,i_n)\in\{1,\ldots,M\}^n} (r_{i_1}\cdots r_{i_n})[g_{\min}, c_k],$$

where

$$c_k := \begin{cases} \min\left\{\dfrac{1}{k-1}, \dfrac{\sum_{i=1}^M r_i}{k-M}, \dfrac{\max_{1\leq i\leq M} r_i}{\lceil k/M\rceil - 1}\right\}, & k > M, \\ \dfrac{1}{k-1}, & 2 \leq k \leq M. \end{cases}$$

Proof. Apply Theorem 3.1 to $P = \mathcal{A}_k$. Here $\delta_{\min} = \delta_{\max} = 1/(k-1)$ and (0.3) gives $G_{\mathcal{A}_k}(C) = (k-1)G^k_{AP}(C)$. Dividing the resulting interval bounds by $k-1$ proves the claim. □

Corollary 3.4 (Small-scale exclusion criterion)**.** *Let $0 < \varepsilon \leq 1/3$, $\lambda = (1-\varepsilon)/2$, and $k \geq 3$. Then*

$$G^k_{AP}(C_\varepsilon) \subseteq \bigcup_{n\in\mathbb{N}_0} \lambda^n[\varepsilon, c_k(\varepsilon)],$$

where

$$c_k(\varepsilon) := \min\left\{\frac{1}{k-1}, \frac{2\lambda}{k-2}, \frac{\lambda}{\lceil k/2\rceil - 1}\right\}.$$

If $\varepsilon > c_k(\varepsilon)$, then $G^k_{AP}(C_\varepsilon) = \varnothing$. If $\lambda c_k(\varepsilon) < \varepsilon$, then $G^k_{AP}(C_\varepsilon)$ contains no interval of the form $(0, r)$. In particular, for $k = 3$ we have $c_3(\varepsilon) = \lambda$, so this conclusion holds whenever

$$\varepsilon > \lambda^2 \quad \Longleftrightarrow \quad \varepsilon > 3 - 2\sqrt{2}.$$

Proof. The middle-ε IFS has $M = 2$, common contraction ratio λ, and first-level gap $g_{\min} = \varepsilon$. Substituting these data into Corollary 3.3 gives the stated formula for $c_k(\varepsilon)$ and the scale containment. The two conclusions follow from Corollary 3.2. For $k = 3$, $\lambda \leq 1/2$ gives

$$c_3(\varepsilon) = \min\left\{\frac{1}{2}, 2\lambda, \lambda\right\} = \lambda.$$

Finally,

$$\varepsilon > \lambda^2 \quad \Longleftrightarrow \quad 4\varepsilon > (1-\varepsilon)^2 \quad \Longleftrightarrow \quad \varepsilon^2 - 6\varepsilon + 1 < 0,$$

which, in $0 < \varepsilon \le 1/3$, is equivalent to $\varepsilon > 3 - 2\sqrt{2}$. □

Corollary 3.5 (Asymmetric three-point patterns)**.** *Let* $0 < \varepsilon \le 1/3$, $\lambda = (1-\varepsilon)/2$, *and* $P_\theta = \{0, \theta, 1\}$ *with* $1/2 \le \theta < 1$. *Then*

$$G_{P_\theta}(C_\varepsilon) \subseteq \bigcup_{n\in\mathbb{N}_0} \lambda^n \left[\frac{\varepsilon}{\theta}, U_\theta(\varepsilon)\right], \qquad U_\theta(\varepsilon) := \min\left\{1, \frac{\lambda}{1-\theta}\right\}.$$

If

$$\lambda U_\theta(\varepsilon) < \frac{\varepsilon}{\theta},$$

then $G_{P_\theta}(C_\varepsilon)$ *contains no interval of the form* $(0, r)$. *The case* $0 < \theta < 1/2$ *follows by reflecting the pattern and replacing* θ *by* $1-\theta$.

Proof. For $1/2 \le \theta < 1$,

$$\delta_{\min}(P_\theta) = 1 - \theta, \qquad \delta_{\max}(P_\theta) = \theta.$$

Theorem 3.1, with $M = 2$, $r_1 = r_2 = \lambda$, and $g_{\min} = \varepsilon$, therefore gives

$$L_{P_\theta} = \frac{\varepsilon}{\theta}$$

and

$$U_{P_\theta} = \min\left\{1, \frac{2\lambda}{1-\theta}, \frac{\lambda}{1-\theta}\right\} = U_\theta(\varepsilon).$$

The blackout criterion is now Corollary 3.2. If $0 < \theta < 1/2$, the reflection $x \mapsto 1 - x$ carries P_θ onto $P_{1-\theta}$. Since C_ε is symmetric about $1/2$, reflection preserves its pattern scale sets, and the result follows from the case $1 - \theta > 1/2$. □

3.1. **A geometric refinement for three-term progressions.** We now specialize to the middle-ε Cantor set from the local setup above. The first-splitting argument records only the diameter and the central gap of the two first-generation intervals. For three-term progressions one can retain more of the product geometry which yields both an exact renormalization formula and a strictly sharper blackout criterion.

Write $C_0 = f_0(C_\varepsilon)$ and $C_1 = f_1(C_\varepsilon)$ for the two first-generation pieces.

Definition 3.6 (First-split spectrum)**.** Define

$$B_\varepsilon := \left\{ t > 0 : \begin{array}{l} \text{there exists } x \text{ with } x, x+t, x+2t \in C_\varepsilon, \\ \text{with at least one of these points in each } C_0, C_1 \end{array} \right\}.$$

Thus B_ε consists of the progression gaps whose first separation occurs at the first construction level.

Lemma 3.7 (Exact first-split decomposition)**.** *For every* $0 < \varepsilon \le 1/3$,

$$\boxed{G^3_{AP}(C_\varepsilon) = \bigcup_{n=0}^{\infty} \lambda^n B_\varepsilon.}$$

Proof. This follows by self-similarity of the middle-ε Cantor set. □

We next compute a level-two outer approximation to B_ε. Let $C_\varepsilon^{(2)}$ be the union of the four level-two construction intervals, which we denote by

$$\begin{aligned} I_{00} &= [0, \lambda^2], & I_{01} &= [\lambda - \lambda^2, \lambda], \\ I_{10} &= [1-\lambda, 1-\lambda+\lambda^2], & I_{11} &= [1-\lambda^2, 1], \end{aligned}$$

where

$$I_0 = [0, \lambda] \quad \text{and} \quad I_1 = [1-\lambda, 1],$$

are the two first-generation construction intervals. Define

$$B_\varepsilon^{(2)} := \left\{ t > 0 : \begin{array}{l} \text{there exists } x \text{ with } x, x+t, x+2t \in C_\varepsilon^{(2)}, \\ \text{and these three points meet both } I_0 \text{ and } I_1 \end{array} \right\}.$$

Since $C_\varepsilon \subseteq C_\varepsilon^{(2)}$ and $C_i \subseteq I_i$ for $i = 0, 1$, we have

$$B_\varepsilon \subseteq B_\varepsilon^{(2)}.$$

For fixed t, let

$$\ell_t^{(1)} = \{(u, v) : v = u + t\}, \qquad \ell_t^{(2)} = \{(u, v) : v = u + 2t\}.$$

A progression $x, x+t, x+2t \in C_\varepsilon^{(2)}$ exists exactly when the first-coordinate projections of

$$\ell_t^{(1)} \cap (C_\varepsilon^{(2)} \times C_\varepsilon^{(2)}) \quad \text{and} \quad \ell_t^{(2)} \cap (C_\varepsilon^{(2)} \times C_\varepsilon^{(2)})$$

intersect. The following proposition computes the permitted parameters.

Proposition 3.8 (Level-two outer approximation)**.** *Set*

$$\rho := \frac{3 - \sqrt{5}}{2}, \qquad \eta := \sqrt{2} - 1, \qquad \kappa := \frac{\sqrt{13} - 1}{6},$$

and

$$a_\lambda := \frac{1 - \lambda - \lambda^2}{2}, \qquad b_\lambda := \frac{1 - \lambda + \lambda^2}{2}, \qquad c_\lambda := 1 - \lambda - \lambda^2.$$

For $1/3 \le \lambda < 1/2$*,*

$$B_\varepsilon^{(2)} = \begin{cases} [\varepsilon, \lambda], & \frac{1}{3} \le \lambda \le \rho, \\ [a_\lambda, b_\lambda], & \rho < \lambda < \eta, \\ [\varepsilon, \lambda^2] \cup [a_\lambda, b_\lambda] \cup [c_\lambda, \lambda], & \eta \le \lambda < \kappa, \\ [\varepsilon, \lambda], & \kappa \le \lambda < \frac{1}{2}. \end{cases}$$

At $\lambda = \eta$*, the first and third intervals in the third line are singletons.*

Proof. We first record a general interval calculation. Let

$$I = [p, p + L], \qquad J = [q, q + L], \qquad K = [r, r + L].$$

The conditions $x \in I$, $x + t \in J$, and $x + 2t \in K$ are equivalent to

$$[p, p + L] \cap [q - t, q + L - t] \cap [r - 2t, r + L - 2t] \neq \varnothing.$$

Comparing every lower endpoint with every upper endpoint shows that the admissible $t \geq 0$ form

$$\Big[\max\left\{q-p-L, \frac{r-p-L}{2}, r-q-L, 0\right\}, \min\left\{q-p+L, \frac{r-p+L}{2}, r-q+L\right\}\Big]. \tag{3.1}$$

provided that the left endpoint does not exceed the right endpoint.

Because C_ε is symmetric about $1/2$, it is enough to analyze only the cases where $x \in I_{00}$ or I_{01}. There are six nondecreasing level-two intervals. Substituting their endpoints into (3.1), with $L = \lambda^2$, gives

$(x, x+t, x+2t)$	candidate interval		nonempty condition
(I_{00}, I_{00}, I_{10})	$[c_\lambda, \lambda^2]$		$\lambda \geq 1/2$
(I_{00}, I_{00}, I_{11})	$[1-2\lambda^2, \lambda^2]$		$\lambda \geq 1/\sqrt{3}$
(I_{00}, I_{01}, I_{10})	$[\varepsilon, \lambda]$	if $\lambda \leq \rho$	
	$[a_\lambda, b_\lambda]$	if $\lambda \geq \rho$	
(I_{00}, I_{01}, I_{11})	$[c_\lambda, \lambda]$		$\lambda \geq \eta$
(I_{01}, I_{01}, I_{10})	$[\varepsilon, \lambda^2]$		$\lambda \geq \eta$
(I_{01}, I_{01}, I_{11})	$[c_\lambda, \lambda^2]$		$\lambda \geq 1/2$.

As $1/3 \leq \lambda < 1/2$, only the middle three rows can contribute. For the central row, the switch occurs when

$$a_\lambda = \varepsilon \quad \text{and} \quad b_\lambda = \lambda,$$

which is equivalent to $\lambda = \rho$. The two side intervals first appear at $\lambda = \eta$, because

$$\varepsilon = \lambda^2 \quad \text{and} \quad c_\lambda = \lambda \quad \Longleftrightarrow \quad \lambda = \eta.$$

Finally,

$$\lambda^2 = a_\lambda \quad \Longleftrightarrow \quad b_\lambda = c_\lambda \quad \Longleftrightarrow \quad 3\lambda^2 + \lambda - 1 = 0,$$

whose positive root is κ. At and beyond this value the three intervals overlap and their union is $[\varepsilon, \lambda]$. This proves the four cases in the statement. For ease of independent verification, Appendix A records all six endpoint comparisons before simplification. □

The level-two geometry improves the previous threshold. Define

$$\varepsilon_* := \frac{4-\sqrt{13}}{3} = 0.1314829081\ldots < 3-2\sqrt{2} = 0.1715728753\ldots.$$

Theorem 3.9 (Improved blackout threshold)**.** *If*

$$\varepsilon_* < \varepsilon \leq \frac{1}{3},$$

then $G^3_{AP}(C_\varepsilon)$ contains no interval of the form $(0, r)$. Moreover, in the additional parameter range

$$\varepsilon_* < \varepsilon \leq 3-2\sqrt{2} \qquad \big(\eta \leq \lambda < \kappa\big),$$

one has the explicit blackout intervals

$$\boxed{\left(\lambda^{n+2}, \frac{1-\lambda-\lambda^2}{2}\,\lambda^n\right) \cap G^3_{AP}(C_\varepsilon) = \varnothing \qquad (n \geq 0).}$$

Proof. When $\varepsilon > 3 - 2\sqrt{2}$, the conclusion follows already from Corollary 3.4. It remains to consider $\varepsilon_* < \varepsilon \leq 3 - 2\sqrt{2}$, equivalently $\eta \leq \lambda < \kappa$. Proposition 3.8 gives

$$B_\varepsilon \subseteq [\varepsilon, \lambda^2] \cup [a_\lambda, b_\lambda] \cup [c_\lambda, \lambda], \qquad \lambda^2 < a_\lambda. \tag{3.2}$$

We also need to rule out interference from larger scales. On $[\eta, \kappa]$,

$$2(\varepsilon - \lambda a_\lambda) = \lambda^3 + \lambda^2 - 5\lambda + 2 > 0. \tag{3.3}$$

Indeed, the polynomial on the right is decreasing on $[0, 1/2]$, and at $\lambda = \kappa$ it equals $(82 - 22\sqrt{13})/27 > 0$.

Fix $n \geq 0$ and let

$$J_n = (\lambda^{n+2}, a_\lambda \lambda^n).$$

By Lemma 3.7, it is enough to show that J_n misses $\lambda^m B_\varepsilon^{(2)}$ for every $m \geq 0$. If $m \geq n+1$, then

$$\max(\lambda^m B_\varepsilon^{(2)}) \leq \lambda^{m+1} \leq \lambda^{n+2}.$$

If $m = n$, the exclusion follows directly from (3.2). Finally, if $m \leq n-1$, then by (3.3),

$$\min(\lambda^m B_\varepsilon^{(2)}) = \varepsilon \lambda^m \geq \varepsilon \lambda^{n-1} > a_\lambda \lambda^n.$$

Thus $J_n \cap G_{AP}^3(C_\varepsilon) = \varnothing$ for every n. Since the intervals J_n accumulate at zero, $G_{AP}^3(C_\varepsilon)$ cannot contain any interval $(0, r)$. □

Remark 3.10. The threshold ε_* is the strongest one obtainable by combining the exact first-split decomposition with this complete level-two outer containment alone. Indeed, when $\lambda \geq \kappa$ (equivalently $\varepsilon \leq \varepsilon_*$), Proposition 3.8 gives $B_\varepsilon^{(2)} = [\varepsilon, \lambda]$, exactly the original first-level bound. A further improvement would therefore require level three or deeper symbolic information.

4. The Hunt-Kan-Yorke Method and Intervals of Pattern Scales

In this section, we build on a result of Hunt-Kan-Yorke [12] to investigate the gap set

$$G_{AP}^k(C) := \{t > 0 : \exists\, x \text{ so that } x, x+t, \ldots, x+(k-1)t \in C\}$$

to quantify the range of common differences that can occur in 3- and 4-term arithmetic progressions contained in an admissible compact set $C \subset \mathbb{R}$; in particular, we provide explicit conditions for the gap sets $G_{AP}^3(C)$ and $G_{AP}^4(C)$ to contain intervals about 0.

Theorem 4.1 (Interior of the 3-Gap Set)**.** *Let $C \subset \mathbb{R}$ be a Cantor set with $d = \operatorname{diam}(C) > 0$ and largest bounded gap G satisfying $|G| \leq 0.067d$ and thickness*

$$\tau(C) \geq \frac{1 - 0.067}{2(0.067)} = 6.96268\ldots$$

Then $G_{AP}^3(C)$ contains the interval $(0, 0.435d]$.

As we will see in the subsections that follow, the key mechanism used to prove Theorem 4.1 is applying Proposition 4.12 to the sets $C_1 = C - s$, $C_2 = C + s$, and $C_3 = C$, where $s > 0$, from which it follows $s \in G_{AP}^3(C)$. Note, however, that Proposition 4.12 is incredibly flexible in the choice of C_i, and one could also choose $C_1 = C + s\theta$, $C_2 = C - s(1-\theta)$, and $C_3 = C$, where $0 < \theta < 1$. Thus it follows as a corollary that $C_{P_\theta}(C)$ also contains an interval.

Corollary 4.2 (Interior of the Asymmetric 3-Point Patterns)**.** *Let $C \subset \mathbb{R}$ be a Cantor set with $d = \operatorname{diam}(C) > 0$ and largest bounded gap G satisfying $|G| \leq 0.067d$ and thickness*

$$\tau(C) \geq \frac{1 - 0.067}{2(0.067)} = 6.96268\ldots$$

Then $G_{P_\theta}(C)$ contains the interval $(0, 0.87d]$.

Theorem 4.3 (Interior of the 4-Gap Set)**.** *Let $C \subset \mathbb{R}$ be a Cantor set with $d = \operatorname{diam}(C) > 0$ and thickness*

$$\tau(C) \geq \frac{1 - 0.015}{2(0.015)} = 32.83333\ldots$$

Then $G^4_{AP}(C)$ contains the interval $(0, 0.3d]$.

This process is split into two parts: §4.1 which provides a self-contained argument building on the work of Hunt-Kan-Yorke [12] to provide explicit and computable conditions for three and four Cantor sets to intersect and §4.2 which applies the results of the previous subsection to prove Theorems 4.1 and 4.3.

4.1. **Intersection of Three Cantor Sets.** In this section, we improve a result of [12]:

Proposition 4.4 (Theorem 2, [12])**.** *There is a function $\varphi(\tau_1, \tau_2)$ which is positive in regions III and IV of Figure 2 such that for all interleaved compact sets $C_1, C_2 \subset \mathbb{R}$ with $\tau(C_1) \geq \tau_1$ and $\tau(C_2) \geq \tau_2$, there is a set $K \subset C_1 \cap C_2$ with thickness at least $\varphi(\tau_1, \tau_2)$.*

In practice, this proposition only establishes that $\tau(C) > 0$. To better explain why this is and how we will improve this proposition, we provide a brief overview of its proof. We postpone introducing some notation until after.

Outline of Proof. Let C_1 and C_2 be sets of sufficiently large thickness; i.e., $\tau_1 := \tau(C_1)$ and $\tau_2 := \tau(C_2)$ lie in regions III or IV of Figure 2.

(i.) Construct a new gap set (K_n) which contains the gaps of C_1 and C_2. Consequently, $K := (\cup_n K_n)^{\complement}$ is contained in $C_1 \cap C_2$.

(ii.) Argue the thickness of K can be determined using the gaps (K_n); i.e.,

$$\tau(K) \geq \frac{d(K_n, K_m)}{|K_m|}. \tag{4.1}$$

In fact, taking the minimum over all such n, m there will exist some function $\varphi(\tau_1, \tau_2)$ such that

$$\tau(K) \geq \varphi(\tau_1, \tau_2).$$

(iii.) Bound the denominator of equation (4.1) using Lemma 4.8.

(iv.) Bound the numerator of equation (4.1) using ([12], Lemma 5).

(v.) Combine the above upper and lower bounds to get

$$\tau(K) \geq \varphi(\tau_1, \tau_2) := \frac{t_* \left((\tau_1 - t_*)(\tau_2 - t_*) - (1 + t_*)^2\right)}{(\tau_1 - t_*)(\tau_2 - t_*) + (1 + t_*)^2 \left(2\max(\tau_1, \tau_2) + 1\right)} \psi_{t_*}(\tau_1, \tau_2)$$

where $0 < t_* < \frac{\tau_1\tau_2 - 1}{\tau_1 + \tau_2 + 2}$ is chosen to maximize $\varphi(\tau_1, \tau_2)$ and

$$\psi_t(\tau_1, \tau_2) := (1 - \lambda_1 - \lambda_2)^N \min\left\{1 - \sigma_1\lambda_1 - \sigma_2 b_*, 1 - \sigma_2\lambda_2 - (1 - \lambda_2)\frac{\sigma_1\lambda_1}{1 + \lambda_1}\right\}$$

where for $i = 1, 2$

$$\lambda_i := \frac{1+t}{\tau_i + 1 + t} \quad \text{and} \quad \sigma_i := \frac{(\tau_i - t)(\tau_{3-i} + 1)}{(\tau_i - t)(\tau_{3-i} - t) - (1+t)^2}.$$

□

Let us elaborate on the function $\psi_t(\tau_1, \tau_2)$ from (v.). In regions III or IV, ([12], Lemma 5) establishes the lower bound on the denominator from (4.1); i.e.,

$$d(K_m, K_n) \geq (1 - \lambda_1 - \lambda_2)^N \min\left\{1 - \sigma_1\lambda_1 - \sigma_2 b_*, 1 - \sigma_2\lambda_2 - (1-\lambda_2)\frac{\sigma_1\lambda_1}{1+\lambda_1}\right\} =: \psi_t(\tau_1, \tau_2),$$

for some interdependent large $N \in \mathbb{N}$ and $b_* \in \mathbb{R}_+$. Observe that as N increases, $\psi_t(\tau_1, \tau_2)$ decreases exponentially to zero and, consequently, the function $\varphi(\tau_1, \tau_2)$—which provides a lower bound for the thickness of C—decreases to zero even faster. Thus, not only is $\psi_t(\tau_1, \tau_2)$ small, it is also incredibly difficult to calculate explicitly. Even if one calculates $\psi_t(\tau_1, \tau_2)$ explicitly, they must then substitute it into $\varphi(\tau_1, \tau_2)$ and optimize for t_*. It is probable that $N = 1$ is sufficient for thickness pairs in region III, though it would take further investigation to verify this claim.

To overcome these difficulties, we ask what constraints would allow us to consider the simpler $N = 0$ case. This leads to the discovery of region IV of Figure 2, and we prove in Lemma 4.9 that

$$\tilde{\psi}_t(\tau_1, \tau_2) := 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2$$

is sufficient as a lower bound. We then substitute this new function into Proposition 4.4 and drop the assumption that $\varphi(\tau_1, \tau_2)$ has been optimized by t_*; consequently, $\tilde{\varphi}_t(\tau_1, \tau_2)$ is calculable and there is a range of valid t that makes $\tilde{\varphi}_t(\tau_1, \tau_2)$ positive and large.

Proposition 4.5. *There is a function*

$$\tilde{\varphi}_t(\tau_1, \tau_2) := \frac{t\left((\tau_1 - t)(\tau_2 - t) - (1+t)^2\right)}{(\tau_1 - t)(\tau_2 - t) + (1+t)^2\left(2\max(\tau_1, \tau_2) + 1\right)}\tilde{\psi}_t(\tau_1, \tau_2), \tag{4.2}$$

where

$$\tilde{\psi}_t(\tau_1, \tau_2) := 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2,$$

which is positive in region IV for infinitely many $0 < t < (\tau_1\tau_2 - 1)/(\tau_1 + \tau_2 + 2)$ such that for all interleaved compact sets $C_1, C_2 \subset \mathbb{R}$ with $\tau(C_1) \geq \tau_1$ and $\tau(C_2) \geq \tau_2$, there is a set $K \subset C_1 \cap C_2$ with thickness at least $\tilde{\varphi}_t(\tau_1, \tau_2)$.

Remark 4.6. One can expand $\tilde{\psi}_t(\tau_1, \tau_2)$ to find

$$\begin{aligned}\tilde{\psi}_t(\tau_1, \tau_2) &= 1 - \frac{(\tau_1 - t)(\tau_2 + 1)}{(\tau_1 - t)(\tau_2 - t) - (1+t)^2} \cdot \frac{1+t}{\tau_1 + 1 + t} - \frac{(\tau_2 - t)(\tau_1 + 1)}{(\tau_1 - t)(\tau_2 - t) - (1+t)^2} \cdot \frac{1+t}{\tau_2 + 1 + t} \\ &= \frac{at^2 + bt + c}{\left[(\tau_1 - t)(\tau_2 - t) - (1+t)^2\right](\tau_1 + 1 + t)(\tau_2 + 1 + t)}\end{aligned}$$

where

$$\begin{aligned}A &= -\left(3\tau_1\tau_2 + 2\tau_1 + 2\tau_2 + 1\right) \\ B &= -\left(\tau_1^2\tau_2 + \tau_1\tau_2^2 + 8\tau_1\tau_2 + 4\tau_1 + 4\tau_2 + 2\right) \\ C &= \tau_1^2\tau_2^2 - 4\tau_1\tau_2 - 2\tau_1 - 2\tau_2 - 1.\end{aligned}$$

Thus we let $t_{\max} = \min\left\{\frac{\tau_1\tau_2-1}{\tau_1+\tau_2+2}, \frac{-B+\sqrt{B^2-4AC}}{2A}\right\}$, and it follows that $\tilde{\psi}_t(\tau_1,\tau_2)$ is positive whenever $0 < t < t_{\max}$.

Remark 4.7. It is known that for $t > 0$ sufficiently close to zero, $\tilde{\varphi}_t(\tau_1,\tau_2)$ is positive; however, the exact value of t that maximizes $\tilde{\varphi}_t(\tau_1,\tau_2)$ increases as τ_1, τ_2 increase.

In our applications (§4.2), we fix $\tau_1 = \tau_2$ and choose a particular t to demonstrate large thickness $\tilde{\varphi}_t(\tau_1,\tau_1)$. We then extend this lower bound to all sets of thickness $\tau \geq \tau_1$.

Thus in what follows we construct the function $\tilde{\psi}_t(\tau_1,\tau_2)$, and the result that

$$\tau(K) \geq \tilde{\varphi}_t(\tau_1,\tau_2)$$

follows for free. Our outline is as follows.

In §4.1.1, we introduce our revised thickness conditions for (τ_1,τ_2); this is region IV of Figure 2.

In §4.1.2, we review how to construct the set K in $C_1 \cap C_2$. In [12], this is accomplished by first taking the largest gap G of C_1, C_2 and then expanding it to absorb nearby gaps that are within distance $t|G|$ for some arbitrary $0 < t < \frac{\tau_1\tau_2-1}{\tau_1+\tau_2+2}$. They then repeat this process until no gaps remain. This creates a new system of gaps (K_n), which are dependent on the number t; note that smaller values of t make for smaller gaps K_n, while larger values of t make for larger gaps K_n. Please note that we let the choice of t be arbitrary; we can choose a particular value later. K is then the complement of the system of gaps K_n.

In §4.1.3, we restate Lemma 4.8. This lemma bounds the denominator and aids in bounding the numerator of (4.1) by establishing how far K_n can extend past the gaps of C_1, C_2 that it contains. As we use this lemma without modification, we leave it as a black box.

In §4.1.4, we prove our key Lemma 4.9 which bounds the numerator of (4.1); i.e.,

$$d(K_n, K_m) \geq \tilde{\psi}_t(\tau_1,\tau_2)\, d(G,H),$$

where

$$\tilde{\psi}_t(\tau_1,\tau_2) := 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2$$

and for $i = 1, 2$

$$\lambda_i := \frac{1+t}{\tau_i+1+t} \quad \text{and} \quad \sigma_i := \frac{(\tau_i-t)(\tau_{3-i}+1)}{(\tau_i-t)(\tau_{3-i}-t)-(1+t)^2}.$$

All of the framework included is to make this paper self-contained; we encourage interested readers to review [12] for further details.

4.1.1. *Regions.* In Figure 2, the curve dividing regions I and II is given by $\tau_1\tau_2 = 1$; it is a consequence of the Newhouse Gap Lemma that sets C_1 and C_2 with thicknesses above the line $\tau_1\tau_2 \geq 1$ have at least one point of intersection; additionally, this result is sharp.

The curves dividing regions II and III are given by the inequalities

$$\tau_1 \geq \tau_2, \quad \tau_1 > \frac{\tau_2^2+3\tau_2+1}{\tau_2^2}, \quad \text{and} \quad \tau_2 > \frac{(2\tau_1+1)^2}{\tau_1^3},$$

or

$$\tau_2 \geq \tau_1, \quad \tau_2 > \frac{\tau_1^2+3\tau_1+1}{\tau_1^2}, \quad \text{and} \quad \tau_1 > \frac{(2\tau_2+1)^2}{\tau_2^3}.$$



FIGURE 2. The intersection of two interleaved compact sets with thicknesses τ_1 and τ_2: can be empty for $(\tau_1, \tau_2) \in I$; is nonempty for $(\tau_1, \tau_2) \in II$; contains a set of positive thickness for $(\tau_1, \tau_2) \in III$; and contains a set with an explicit minimum thickness in IV.

These inequalities come from the original argument in Hunt-Kan-Yorke [12], and all thickness pairs (τ_1, τ_2) in regions III or IV guarantee the intersection of C_1 and C_2 contain a set of positive thickness. Moreover, this dividing line is sharp; see [15].

The red line dividing regions III and IV is our new contribution and the focus of this section, §4; it is given by

$$\tau_1^2\tau_2^2 - 4\tau_1\tau_2 - 2\tau_1 - 2\tau_2 - 1 = 0,$$

or equivalently

$$\tau_i = \frac{1 + 2\tau_j + \sqrt{2\tau_j^3 + 5\tau_j^2 + 4\tau_j + 1}}{\tau_j^2},$$

for $i \neq j$. **We show that all thickness pairs (τ_1, τ_2) in region IV not only contain a set of positive thickness, but we obtain an explicit lower bound for that thickness.**

4.1.2. *Construction of a new system of gaps K_n.* We revisit how to construct the sets (K_n). Recall that we want the gaps (K_n) to contain all the gaps of C_1 and C_2 from which it follows that $K \subset C_1 \cap C_2$ where $K := (\cup_n K_n)^{\complement}$.

Let C_1 and C_2 be interleaved compact sets with $\tau(C_1) \geq \tau_1$ and $\tau(C_2) \geq \tau_2$ for some (τ_1, τ_2) in region IV of Figure 2.

Let the gaps of C_1 be $I_0, I_1, I_2, \ldots$, with I_0 and I_1 unbounded, I_0 to the left of I_1, and $|I_2| \geq |I_3| \geq \cdots$. For C_2 we define $J_0, J_1, J_2, \ldots$ similarly. We refer to the intervals I_n and J_n collectively as the "original gaps." Our goal is to construct the complement of K as a union of disjoint open intervals $K_0, K_1, K_2, \ldots$ with K_0 and K_1 unbounded, and with every original gap contained in some K_m whence $K \subset C_1 \cap C_2$. To get a lower bound on the thickness of K, observe that every chunk P of K is bordered on each side by a gap of K, with at least



FIGURE 3. Cases in the construction of I_* and J_*.

one of the bordering gaps being bounded. Pick a chunk P, and say P is bordered by K_m and K_n with $m > n$ and $m \geq 2$. Then

$$\frac{|P|}{d(P, K \setminus P)} = \frac{d(K_m, K_n)}{\min(|K_m|, |K_n|)} \geq \frac{d(K_m, K_n)}{|K_m|}.$$

Recalling that thickness is defined as

$$\tau(K) = \inf_{P \propto K} \frac{|P|}{d(P, K \setminus P)},$$

we see that finding a universal lower bound of $\frac{d(K_m,K_n)}{|K_m|}$ would ensure that the thickness is also bounded. Hence, Hunt-Kan-Yorke show that there exists some $\tilde{\varphi}_t(\tau_1, \tau_2) > 0$ such that whenever $m > n$ and $m \geq 2$,

$$\frac{d(K_m, K_n)}{|K_m|} \geq \tilde{\varphi}_t(\tau_1, \tau_2). \tag{4.3}$$

In this paper, we focus on improving the bound of the numerator (i.e., bounding $d(K_m, K_n)$) through improving the bound on $\tilde{\psi}_t(\tau_1, \tau_2)$.

We return to the construction of the (K_n) by finding a pair of original gaps I_* and J_* between which K will lie; that is I_* and J_* will be contained in K_0 and K_1. The properties we desire of I_* and J_* are that they are a positive distance apart, that all gaps of C_1 with an endpoint between the closures of I_* and J_* are bounded and no larger than I_* and likewise (in comparison to J_*) for gaps of C_2 between I_* and J_*. We will show later that once I_* and J_* have been determined, the diameter of K can be bounded below by a constant depending on τ_1 and τ_2 times the distance between I_* and J_*.

Assume without loss of generality that $J_0 \subset I_0$. If $I_1 \subset J_1$, Case 1 of Figure 3, then $I_* = I_0$ and $J_* = J_1$ have the above properties; they must be separated by a positive distance since C_1 and C_2 are interleaved. If $J_1 \subset I_1$, Case 2 of Figure 3, let J_* be the largest gap of C_2 with an endpoint between I_0 and I_1, and let I_* be whichever of I_0 and I_1 is farthest from J_*. At least one of I_0 and I_1 must be a positive distance from J_* since C_1 and C_2 are interleaved.

Next, let t be a positive constant whose precise value will be chosen later; for now we assume that $0 < t < (\tau_1\tau_2 - 1)/(\tau_1 + \tau_2 + 2) < \min(\tau_1, \tau_2)$. Assume without loss of generality that I_* lies to the left of J_*. We begin constructing K_0 by requiring that it contain I_*. We then require that K_0 contain the rightmost bounded J_n with $d(I_*, J_n) \leq t|J_n|$ (we leave the verification that there is a rightmost gap satisfying this condition to the original authors). If there does not exist such a J_n that is not already contained in I_*, we stop the construction and let $K_0 = I_*$. Otherwise, we further require that K_0 contain the rightmost bounded I_m that is within t times its length of the previously added J_n. Again, if this requirement does



FIGURE 4. The construction of K_0.

not extend K_0 any farther rightward, we stop the construction. If not, we then add to K_0 the rightmost J_l which is within t times its length of I_m and is at most as large as J_n (see Figure 4). If a next step is necessary, we consider gaps of C_1 which are no larger than I_m, and so forth. We may have to continue this process infinitely often, but if so we must converge to a right endpoint for K_0, since there is no way this construction can extend past the rightmost point in $C_1 \cup C_2$.

We define K_1 similarly, starting with the requirement that K_1 contain J_* and extending K_1 to the left if necessary in the same way we constructed K_0.

Next, to construct K_2 we first require that it contain the largest original gap (choose any one in the case of a tie) not contained in $K_0 \cup K_1$ (if no such gap exists, we leave K_2 undefined and let S be the complement of $K_0 \cup K_1$). Then we extend it on both the left and right in the same manner as before, but considering only gaps that are at most as large as the one we started with, to obtain the endpoints of K_2. We next start with the largest original gap not contained in $K_0 \cup K_1 \cup K_2$, proceeding similarly to define K_3, and so forth. Any given original gap must eventually be contained in some K_n because there can only be finitely many original gaps that are as large or larger than the given one. We do not yet know that the K_n are disjoint from each other; this will follow when we prove (4.3), though.

Let us now examine our construction more closely. Define $l(I)$ and $r(I)$ to be the left and right endpoints of an interval I, respectively. For a given K_n, let G_0 be the gap we started with in its construction, which for $n \geq 2$ must be the largest original gap it contains, or at least tied for the largest. For simplicity and without loss of generality we assume that G_0 is a gap of C_1. Consider the collection E of all J_n gaps of C_2 with $|J_n| \leq |G_0|$, $r(J_n) > r(G_0)$, and $d(G_0, J_n) \leq t|J_n|$. We claim—see [12] for details—that the members of E are increasing in size from left to right.

We likewise define G_2 to be the rightmost gap of C_1 which is at most as large as G_0 and lies within t times its length of G_1; again, if no such gap exists with $r(G_2) > r(G_1)$ we say that $|G_2| = 0$ and $r(G_2) = r(G_1)$. Next, to define G_3 we consider only gaps of S_2 which are at most as large as G_1, for G_4 we look only at gaps of C_1 no larger than G_2, and so forth. Define $G_{-1}, G_{-2}, \cdots$ similarly to be the leftmost (and largest) gaps added to K_n at each stage of the process extending K_n leftward. Then we may think of the open interval K_n as being defined by

$$l(K_n) = \lim_{m \to -\infty} l(G_m), \quad r(K_n) = \lim_{m \to \infty} r(G_m).$$

Each limit exists because it is the limit of a bounded monotonic sequence.

In the above construction, the even-numbered G_m are gaps of C_1 and the odd-numbered ones are gaps of C_2, but if G_0 had been a gap of C_2 it would be the other way around. In any case, G_0 is the largest even-numbered G_m and either G_1 or G_{-1} is the largest odd-numbered one. Also, the even-numbered G_m decrease monotonically in size as one moves either rightward or leftward from the largest, and the same statement holds for the odd-numbered G_m. We

call a given G_m either a "1-gap" or "2-gap" of K_n according to whether it is a gap of C_1 or C_2. Notice that not all original gaps contained in K_n are 1-gaps or 2-gaps, only those that have been given a label G_m in the construction of K_n. When we refer henceforth to left-to-right ordering or adjacency among the 1-gaps and 2-gaps of a given K_n, it is with respect to the ordering $\ldots, G_{-2}, G_{-1}, G_0, G_1, G_2, \ldots$. Thus, 1-gaps can only be adjacent to 2-gaps and vice versa.

This concludes our review of the construction of the gaps (K_n). The above notation will be used in the following lemmas, and further information can be found in [12].

4.1.3. *Black Box Lemma.* The following lemma is used in bounding both the numerator and denominator of the left side of inequality (4.3). For all $m \geq 0$, it establishes a bound on how far K_n can extend to the right of G_m in terms of how far G_{m+1} extends past G_m, and similarly for $m \leq 0$ on the left.

Lemma 4.8 (Lemma 4, [12])**.** *Assume* $0 < t < (\tau_1\tau_2 - 1)/(\tau_1 + \tau_2 + 2)$. *Let*

$$\sigma_1 = \frac{(\tau_1 - t)(\tau_2 + 1)}{(\tau_1 - t)(\tau_2 - t) - (1 + t)^2}$$

and

$$\sigma_2 = \frac{(\tau_2 - t)(\tau_1 + 1)}{(\tau_1 - t)(\tau_2 - t) - (1 + t)^2}.$$

Let G be a 1-gap of K_n which is at least as large as all 1-gaps of K_n to its right. Let H be the next 2-gap of K_n to the right of G. Then

$$r(K_n) - r(G) \leq \sigma_2(r(H) - r(G)).$$

The same statement with "1" and "2" interchanged holds, as do the corresponding results for left endpoints.

Lemma 4.8 is a key tool in proving both the original Lemma 5, [12] and our modification (Lemma 4.9) below; hence, we will treat this lemma as a black box in what follows.

4.1.4. *Explicit formula for $\tilde{\psi}_t(\tau_1, \tau_2)$.* As established above, we want to show that the left side of (4.3) is positive for all $m > n$ and $m \geq 2$. Lemma 4.9 provides an improved bound for the denominator $d(K_n, K_m)$ in region IV:

Lemma 4.9. *There exists a function*

$$\tilde{\psi}_t(\tau_1, \tau_2) := 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2$$

that is positive whenever $(\tau_1, \tau_2) \in IV$, for which the following statement holds. For $m \neq n$, let G be a 1-gap of K_m and H be a 2-gap of K_n. If all 1-gaps of K_m or K_n with at least one endpoint between the closures of G and H are bounded and at most as large as G, and all similarly situated 2-gaps are bounded and at most as large as H, then

$$d(K_n, K_m) \geq \tilde{\psi}_t(\tau_1, \tau_2)d(G, H).$$

Proof of Lemma 4.9. Let G be a 1-gap of K_m and H be a 2-gap of K_n satisfying the hypotheses. We assume without loss of generality that $(\tau_1, \tau_2) \in IV$, so

$$\tau_i > \frac{1 + 2\tau_j + \sqrt{2\tau_j^3 + 5\tau_j^2 + 4\tau_j + 1}}{\tau_j^2},$$



FIGURE 5. The gaps G_i and H_i.

for $i \neq j$. If $d(G, H) = 0$, the inequality to be proven is trivial. Otherwise, let us normalize $d(G, H)$ to one, and assume G lies to the left of H. Let $G_0 = G$ and $H_0 = H$. Let G_1 be the 1-gap of K_n adjacent to H_0 on its left, and let H_1 be the 2-gap of K_m adjacent to G_0 on its right. Let G_2 be the adjacent 1-gap of K_m rightward from H_1, and likewise define $H_2, G_3, H_3, \ldots$. For $i \geq 0$ let

$$x_i = \begin{cases} l(H_i) - l(G_{i+1}), & i \text{ even}, \\ r(G_{i+1}) - r(H_i), & i \text{ odd}, \end{cases}$$

and

$$y_i = \begin{cases} r(H_{i+1}) - r(G_i), & i \text{ even}, \\ l(G_i) - l(H_{i+1}), & i \text{ odd}. \end{cases}$$

Let $R_i = d(G_i, H_i)$; then $R_0 = 1$ and $R_{i+1} = \max(R_i - x_i - y_i, 0)$ for $I \geq 0$. Let R_∞ be the limit as i goes to infinity of R_i. Then $d(K_m, K_n) = R_\infty$, so we wish to show that there is a positive lower bound on R_∞ which depends only on τ and t.

We first find bounds for x_i and y_i in terms of each other. For all even $i \geq 0$, G_i are 1-gaps of K_n, H_{i+1} are 2-gaps of K_m, and $|G_i| \geq |G_{i+2}|$; see Figure 5. Then

$$\tau_1 |G_{i+2}| \leq d(G_i, G_{i+2}) \leq d(H_{i+1}, G_{i+2}) + r(H_{i+1}) - r(G_i) \leq t|G_{i+2}| + r(H_{i+1}) - r(G_i),$$

where the first inequality follows from the definition of thickness and the last inequality follows from construction of the 1- and 2-gaps; i.e., G_{i+2} is the largest gap of S_1 with size at most $|G_i|$ within distance $t|G_{i+2}|$ of H_{i+1}. Because $t \leq \tau_1$, we re-arrange as

$$|G_{i+2}| \leq \frac{r(H_{i+1}) - r(G_i)}{\tau_1 - t}.$$

Hence,

$$\begin{aligned} x_{i+1} = r(G_{i+2}) - r(H_{i+1}) &\leq |G_{i+2}| + d(H_{i+1}, G_{i+2}) \\ &\leq (1+t)|G_{i+2}| \leq \frac{1+t}{\tau_1 - t} \left(r(H_{i+1}) - r(G_i) \right) = \frac{1+t}{\tau_1 - t} y_i. \end{aligned}$$

A similar argument works for odd i; consequently,

$$x_{i+1} \leq \frac{1+t}{\tau_1 - t} y_i. \tag{4.4}$$

Similarly, for y_i

$$y_{i+1} \leq \frac{1+t}{\tau_2 - t} x_i. \tag{4.5}$$

Furthermore, by Lemma 4 [12] we have

$$y_i + x_{i+1} + y_{i+2} + \cdots \leq \sigma_1 y_i$$

and

$$x_i + y_{i+1} + x_{i+2} + \cdots \leq \sigma_2 x_i.$$

Thus, for each i,

$$d(K_m, K_n) = R_\infty \geq R_i - x_i - y_i - x_{i+1} - y_{i+1} - \cdots \geq R_i - \sigma_1 x_i - \sigma_2 y_i, \tag{4.6}$$

We will show that for some i, the right side of (4.6) is positive. In fact, in region IV we can take $i = 0$.

Let us obtain upper bounds on x_0 and y_0. We know

$$x_0 = l(H_0) - l(G_1) \leq |g_1| + d(G_1, H_0) \leq (1+t)|G_1|, \tag{4.7}$$

and by hypothesis $|G_1| \leq |G_0|$, so

$$x_0 = l(H_0) - r(G_0) - [l(G_1) - r(G_0)] = 1 - d(G_0, G_1) \leq 1 - \tau_1|G_1|. \tag{4.8}$$

Eliminating $|G_1|$ from these inequalities yields

$$x_0 \leq \frac{1+t}{\tau_1 + 1 + t} =: \lambda_1. \tag{4.9}$$

Similarly,

$$y_0 \leq \frac{1+t}{\tau_2 + 1 + t} =: \lambda_2. \tag{4.10}$$

We can obtain similar bounds on x_i and y_i for $i \geq 1$, but it is no longer necessary.

We will show that at $t = 0$,

$$1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 > 0,$$

from which it holds for t sufficiently near 0 by continuity. This in turn will imply

$$d(K_m, K_n) \geq R_0 - \sigma_1 x_0 - \sigma_2 y_0 \geq 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 > 0.$$

Observe that at $t = 0$,

$$\begin{aligned} 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 &= 1 - \frac{\tau_1(\tau_2+1)}{\tau_1\tau_2 - 1}\frac{1}{\tau_1+1} - \frac{\tau_2(\tau_1+1)}{\tau_1\tau_2-1}\frac{1}{\tau_2+1} \\ &= \frac{(\tau_1\tau_2 - 1)(\tau_1+1)(\tau_2+1) - \tau_1(\tau_2+1)^2 - \tau_2(\tau_1+1)^2}{(\tau_1\tau_2-1)(\tau_1+1)(\tau_2+1)} \\ &= \frac{\tau_1^2\tau_2^2 - 4\tau_1\tau_2 - 2\tau_1 - 2\tau_2 - 1}{(\tau_1\tau_2-1)(\tau_1+1)(\tau_2+1)}. \end{aligned}$$

Hence, $1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 > 0$ if and only if $(\tau_i^2)\tau_j^2 + (-2 - 4\tau_i)\tau_j + (-1 - 2\tau_i) > 0$ for $i \neq j$ which is true so long as

$$\tau_i > \frac{1 + 2\tau_j + \sqrt{2\tau_j^3 + 5\tau_j^2 + 4\tau_j + 1}}{\tau_j^2},$$

for $i \neq j$. Thus, $1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 > 0$ for $t = 0$ and (τ_1, τ_2) in region IV. Thus, $d(K_m, K_n) \geq 1 - \sigma_1\lambda_1 - \sigma_2\lambda_2 > 0$ for t sufficiently near 0. Recalling that $d(G, H)$ was normalized to 1, the conclusion follows. □

Remark 4.10. We emphasize a particular case of Lemma 4.9 as we use it in §4.2.

Let C_1, C_2 be two compact sets with thicknesses in region IV such that the intersection of their convex hulls, written $Q := \text{conv}(C_1) \cap \text{conv}(C_2)$, contains neither C_1 nor C_2. Then Proposition 4.5 applies and we obtain a compact set $K \subset C_1 \cap C_2$ constructed from the gaps (K_n); in particular, we can apply Lemma 4.9 to the unbounded gaps K_0 and K_1 of K from whence

$$d(K_0, K_1) \geq \tilde{\psi}_t(\tau_1, \tau_2) d(I_*, J_*).$$

Because Q contains neither C_1 nor C_2, we are in Case 1 of Figure 3; this implies

$$|\,\text{conv}(K)| \geq \tilde{\psi}_t(\tau_1, \tau_2)|Q|.$$

Remark 4.11. In the special case where $\tau := \tau_1 = \tau_2$, we have

$$\tilde{\psi}_t(\tau) = 1 - 2\frac{\tau - t}{\tau - 2t - 1}\frac{1 + t}{\tau + 1 + t} = \frac{\tau^2 - 2\tau - 1 - t(3\tau + 1)}{(\tau - 2t - 1)(\tau + 1 + t)}, \tag{4.11}$$

which is positive so long as $0 < t < \frac{\tau^2 - 2\tau - 1}{3\tau + 1}$. Consequently, referring to the definition of $\tilde{\varphi}_t(\tau_1, \tau_2)$ in (4.2), we can simplify $\tilde{\varphi}_t$ as

$$\tilde{\varphi}_t(\tau) = \frac{t\left[(\tau - t)^2 - (1 + t)^2\right]\tilde{\psi}_t(\tau)}{(\tau - t)^2 + (1 + t)^2(2\tau + 1)} = \frac{t(\tau - 2t - 1)\tilde{\psi}_t(\tau)}{\tau + 1 + 2t(t + 1)}, \tag{4.12}$$

which is positive so long as $0 < t < \frac{\tau - 1}{2}$ and $\tilde{\psi}_t(\tau) > 0$.

Putting this together, $\tilde{\varphi}_t(\tau)$ is positive so long as

$$0 < t < \frac{\tau^2 - 2\tau - 1}{3\tau + 1} < \frac{\tau - 1}{2}.$$

As a corollary, Hunt-Kan-Yorke explain when the Gap Lemma can be applied to the set $K \subset C_1 \cap C_2$ and some other set C_3. The argument follows by explaining how to apply the Gap Lemma 1.3 to the sets K and C_3.

Proposition 4.12 (Corollary 6, [12])**.** *Let C_1 and C_2 be two interleaved compact sets whose thicknesses (τ_1, τ_2) lie in region IV and for which the intersection Q of their convex hulls contains neither C_1 nor C_2. If C_3 is a compact set with largest bounded gap G such that*

(i) the convex hull of C_3 contains Q,
(ii) $|G| < \tilde{\psi}_t(\tau_1, \tau_2)|Q|$, and
(iii) $\tau(C_3)\tilde{\varphi}_t(\tau_1, \tau_2) \geq 1$,

for $0 < t < t_{max}$. Then $C_1 \cap C_2 \cap C_3$ is nonempty.

Proof. Indeed, let C_1, C_2 be two compact sets with thicknesses in region IV. Apply Proposition 4.5 to C_1 and C_2 to obtain the set $K \subset C_1 \cap C_2$ with minimum thickness $\tilde{\varphi}_{t_*}(\tau_1, \tau_2)$, for some $0 < t_* < t_{\max}$. To apply the Gap Lemma 1.3, we need to establish that neither C_3 nor K lie in a gap of the other and that the product of their thicknesses is greater than or equal to 1.

To see that neither set lies in a gap of the other, we first need information about the diameter of K. Recall that K lies between the unbounded gaps K_0 and K_1, so we use Remark 4.10 to obtain

$$|\,\text{conv}(K)| \geq \tilde{\psi}_t(\tau_1, \tau_2)|Q|,$$

where $Q := \text{conv}(C_1) \cap \text{conv}(C_2)$. Thus condition (i) guarantees

$$C \subset Q \subset \text{conv}(C_3),$$

and condition (ii) guarantees the largest gap G of C_3 is not wider than $\operatorname{conv}(K)$:

$$|G| < \tilde{\psi}_t(\tau_1, \tau_2)|Q| < |\operatorname{conv}(K)|.$$

Hence, we conclude that neither K nor C_3 lie in a gap of the other.

Lastly, condition (iii) establishes that the product of the thicknesses is greater than one; i.e.,

$$\tilde{\varphi}_t(\tau_1, \tau_2)\tau(K)\tau(C_3) \geq \tau(C_3) \geq 1. \tag{4.13}$$

Thus we conclude $C_1 \cap C_2 \cap C_3$ is nonempty. □

An analogous argument can be used to obtain quadruple intersections $C_1 \cap C_2 \cap C_3 \cap C_4 \neq \varnothing$, for compact subsets $C_i \subset \mathbb{R}$, by applying Proposition 4.12 to C_1, C_2 and C_3, C_4 to obtain new compact sets $K_1 \subset C_1 \cap C_2$ and $K_2 \subset C_3 \cap C_4$ of thickness $\tau(K_1), \tau(K_2) \geq 1$. One can then apply the Gap Lemma 1.3 to K_1 and K_2 to show $C_1 \cap C_2 \cap C_3 \cap C_4 \neq \varnothing$. However, this case requires strict conditions on the convex hulls $\operatorname{conv}(C_i)$ to ensure that the sets K_1 and K_2 are interleaved. Thus, we show only the special case which proves the existence of 4-term arithmetic progressions; i.e., $C_i := C + is$ for $0 \leq i \leq 3$.

Proposition 4.13 (four-term arithmetic progression criterion)**.** *Let $C \subset \mathbb{R}$ be compact and $0 < t < \frac{\tau^2 - 2\tau - 1}{3\tau + 1}$ be such that*

$$\tilde{\varphi}_t(\tau) \geq 1,$$

where $\tau := \tau(C)$. It necessarily follows that $\tau \gg 1 + \sqrt{2}$. Then for $d = \operatorname{diam}(C) > 0$,

$$\left[0, \frac{\tilde{\psi}_t(\tau)}{1 + 2\tilde{\psi}_t(\tau)} d\right] \subset G^4_{AP}(C).$$

Proof. We will show $C \cap (C + s) \cap (C + 2s) \cap (C + 3s) \neq \varnothing$ for $s \in \left[0, \frac{\tilde{\psi}_t(\tau)}{1+2\tilde{\psi}_t(\tau)} d\right]$ by first applying Proposition 4.5 to the intersection $C \cap (C + 2s)$. As thickness is invariant under similarities and arithmetic-progression gaps scale linearly, it is enough to treat the normalization $\operatorname{conv}(C) = [0, 1]$. Because $0 < t < \frac{\tau^2 - 2\tau - 1}{3\tau + 1}$, we know $\tilde{\psi}_t(\tau) > 0$, and by hypothesis $\tilde{\varphi}_t(\tau) > 1$. Let

$$0 \leq s \leq \frac{\tilde{\psi}_t(\tau)}{1 + 2\tilde{\psi}_t(\tau)}. \tag{4.14}$$

Since $\tilde{\psi}_t(\tau)/(1 + 2\tilde{\psi}_t(\tau)) < 1/2$, we have $s < 1/2$. Therefore the sets

$$C_0 := C, \qquad C_2 := C + 2s$$

are interleaved: the right endpoint $1 \in C_0$ lies in $(2s, 1 + 2s) = \operatorname{int}\operatorname{conv}(C_2)$, and the left endpoint $2s \in C_2$ lies in $(0, 1) = \operatorname{int}\operatorname{conv}(C_0)$. Thus we apply Proposition 4.5 to C_0 and C_2 and obtain a compact set

$$K \subseteq C \cap (C + 2s)$$

for which

$$\tau(K) \geq \tilde{\varphi}_t(\tau),$$

and utilizing Remark 4.10

$$\operatorname{diam}(K) = |\operatorname{conv}(K)| \geq \tilde{\psi}_t(\tau)(1 - 2s).$$

The choice of s is in (4.14) equivalent to

$$\tilde{\psi}_t(\tau)(1-2s) \geq s;$$

consequently, $s \leq \operatorname{diam}(K)$.

We now investigate the set $(C+s) \cap (C+3s)$ by translating our set K from above: Translating the inclusion $K \subseteq C \cap (C+2s)$ gives

$$K+s \subseteq (C+s) \cap (C+3s), \qquad \tau(K+s) = \tau(K) \geq \tilde{\varphi}_t(\tau).$$

Because $s \leq \operatorname{diam}(K)$, Lemma 1.4 shows that K and $K+s$ are interleaved. Moreover,

$$\tau(K)\tau(K+s) = \tau(K)^2 \geq \tilde{\varphi}_t(\tau)^2 \geq 1.$$

Then applying the Newhouse Gap Lemma gives a point

$$x \in K \cap (K+s).$$

Since $x \in K \subseteq C \cap (C+2s)$, we have

$$x \in C, \qquad x-2s \in C.$$

Since $x \in K+s$, the point $x-s$ belongs to K; consequently

$$x-s \in C, \qquad x-3s \in C.$$

Thus

$$x-3s, \quad x-2s, \quad x-s, \quad x$$

is a four-term arithmetic progression in C with common difference s; hence $s \in G^4_{AP}(C)$.

Rescaling restores the factor d. □

This process of combining the Gap Lemma with the HKY criterion can be repeated for intersections of five or more sets; however, there is significant attrition of necessary information (e.g., exact length and position of convex hulls) that make this style of argument deteriorate. Thus we conclude our modifications of [12] here.

4.2. **Non-Empty Gap Set.** In this section, we prove Theorem 4.1; i.e., the gap set,

$$G^3_{AP}(C) := \{t > 0 : \exists\, x \text{ so that } x, x+t, x+2t \in C\},$$

for sufficiently thick C—with conditions on the largest gap G_1—contains the interval $(0, 0.432]$. With our new Lemma 4.9 and Proposition 4.5, we now do so by utilizing Proposition 4.12:

Notice that if $(C-s) \cap C \cap (C+s) \neq \varnothing$ for a shift $s > 0$, then C contains a 3-term arithmetic progression $\{c-s, c, c+s\}$ for some $c \in C$; hence, we consider the case where $C_1 = C-s$, $C_2 = C+s$, and $C_3 = C$ for a shift $s > 0$. Because thickness is translation invariant, we know $\tau := \tau_1 = \tau_2 = \tau(C)$; hence, we use the simplified $\tilde{\varphi}_t(\tau)$ and $\tilde{\psi}_t(\tau)$ in equations (4.12) and (4.11), respectively.

We prove Theorem 4.1 for a fixed thickness $\tau_\star$ and a fixed $t_\star$. Then, we extend to sets of larger thickness $\tau \geq \tau_\star$ by claiming that

$$\tilde{\psi}_{t_\star}(\tau) \geq \tilde{\psi}_{t_\star}(\tau_\star) \quad \text{and} \quad \tilde{\varphi}_{t_\star}(\tau) \geq \tilde{\varphi}_{t_\star}(\tau_\star,)$$

for all $\tau \geq \tau_\star$.

We begin by verifying the above claims.

Claim 4.14. For fixed $0 < t < \frac{\tau-1}{2}$, $\tilde{\psi}_t(\tau)$ increases as τ increases.

Proof. Fix $0 < t < \frac{\tau-1}{2}$. Then by equation (4.11),

$$\tilde{\psi}_t(\tau) = \frac{\tau^2 - (2+3t)\tau - 1 - t}{\tau^2 - t\tau - (2t+1)(t+1)}.$$

Taking the derivative with respect to τ yields

$$\begin{aligned}\tilde{\psi}_t'(\tau) &= \frac{\begin{array}{r}[\tau^2 - t\tau - (2t+1)(t+1)]\,[2\tau - (2+3t)] \\ -\,[\tau^2 - (2+3t)\tau - 1 - t]\,[2\tau - t]\end{array}}{[\tau^2 - t\tau - (2t+1)(t+1)]^2} \\ &= \frac{2(1+t)\tau^2 - t(4t+5)\tau + 2(1+t)(3t^2+3t+1)}{[\tau^2 - t\tau - (2t+1)(t+1)]^2}.\end{aligned}$$

Thus, $\tilde{\psi}_t'(\tau)$ is positive whenever $n_t(\tau) = 2(1+t)\tau^2 - t(4t+5)\tau + 2(1+t)(3t^2+3t+1)$ is positive. Observe that $n_t(\tau)$ is an upward facing parabola with negative discriminant $D = -32t^4 - 104t^3 - 135t^2 - 90t - 16$; hence, $n_t(\tau)$ is always positive which means $\tilde{\psi}_t(\tau)$ is increasing for fixed t. □

Claim 4.15. For fixed $0 < t < \frac{\tau-1}{2}$, $\tilde{\varphi}_t(\tau)$ increases as τ increases.

Proof. Fix $0 < t < \frac{\tau-1}{2}$. Recall from equation (4.12) that

$$\begin{aligned}\tilde{\varphi}_t(\tau) &= \frac{t(\tau^2 - 2\tau - 1) - t^2(3\tau + 1)}{(\tau+1)^2 + 3(\tau+1)t + 2(\tau+2)t^2 + 2t^3} \\ &= \frac{t\tau^2 + (-3t^2 - 2t)\tau + (-t^2 - t)}{\tau^2 + (2t^2+3t+2)\tau + (2t^3+4t^2+3t+1)}.\end{aligned}$$

Taking the derivative with respect to τ,

$$\begin{aligned}\tilde{\varphi}_t'(\tau) &= \frac{\begin{array}{r}[\tau^2 + (2t^2+3t+2)\tau + (2t^3+4t^2+3t+1)]\,[2t\tau + (-3t^2-2t)] \\ -\,[t\tau^2 + (-3t^2-2t)\tau + (-t^2-t)]\,[2\tau + (2t^2+3t+2)]\end{array}}{[\tau^2 + (2t^2+3t+2)\tau + (2t^3+4t^2+3t+1)]^2} \\ &= \frac{(2t^3+6t^2+4t)\tau^2 + (4t^4+8t^3+8t^2+4t)\tau + (-6t^5-14t^4-12t^3-4t^2)}{[\tau^2 + (2t^2+3t+2)\tau + (2t^3+4t^2+3t+1)]^2}.\end{aligned}$$

Hence, $\tilde{\varphi}_t'(\tau)$ is positive when the numerator

$$n_t(\tau) = (2t^3+6t^2+4t)\tau^2 + (4t^4+8t^3+8t^2+4t)\tau + (-6t^5-14t^4-12t^3-4t^2)$$

is positive. Note that $n_t(\tau)$ is an upward-facing parabola and $n_t(0) = -6t^5-14t^4-12t^3-4t^2 < 0$ for $0 < t$. Thus, if $n_t(\rho) > 0$ for some $\rho > 0$, then $n_t(\tau) > 0$ for all $\tau \geq \rho$. Observe that $0 < t < \frac{\tau-1}{2}$ is equivalent to $1 < 2t+1 < \tau$. Then

$$\begin{aligned}n_t(\tau) &= (2t^3+6t^2+4t)\tau^2 + (4t^4+8t^3+8t^2+4t)\tau + (-6t^5-14t^4-12t^3-4t^2) \\ &> (2t^3+6t^2+4t)(2t+1)^2 + (4t^4+8t^3+8t^2+4t)(2t+1) + (-6t^5-14t^4-12t^3-4t^2) \\ &= 10t^5 + 38t^4 + 54t^3 + 34t^2 + 8t > 0.\end{aligned}$$

Thus for fixed t the function $\tilde{\varphi}_t(\tau)$ increases as τ increases. □

We now have the prerequisite tools to prove that the gap set of sufficiently thick C with sufficiently small largest gap contains an interval including 0.

Proof of Theorem 4.1. Let C be a compact set of thickness

$$\tau := \tau(C) \geq 6.96,$$

where, by similarity invariance of thickness, we normalize $\operatorname{conv}(C) = [0, 1]$. Let the largest gap G satisfy $|G| \leq 0.067$ and take $s \in (0, 0.435]$. We will apply Proposition 4.12 to the sets

$$C_1 = C - s, \quad C_2 = C + s, \quad \text{and} \quad C_3 = C$$

using the function $\tilde{\psi}_t(\tau)$ from Lemma 4.9 to obtain $C_1 \cap C_2 \cap C_3 \neq \varnothing$; in particular, that $G^3_{\mathrm{AP}}(C) \supset (0, 0.435]$. We do so by comparing against the anchor values $\tau_\star = 6.96$, $|G_\star| = 0.067$, and $t_\star = 0.55$ and using Claims 4.14 and 4.15 to extend to sets of larger thickness.

First, note that C_1 and C_2 are interleaved because $s < 1/2$ and because $Q = [s, 1 - s]$, we know Q contains neither C_1 nor C_2.

(i) Observe that $\operatorname{conv}(C_3) = [0, 1] \supset [s, 1 - s] = Q$.

(ii) Next, we want $|G| < \tilde{\psi}_{t_\star}(\tau)|Q|$ which is equivalent to

$$\tilde{\psi}_{t_\star}(\tau) > \frac{|G|}{1 - 2s},$$

which holds for our chosen $t_\star$, $|G_\star|$, and $s \in (0, 0.435]$ as

$$\tilde{\psi}_{t_\star}(\tau) \geq \tilde{\psi}_{t_\star}(\tau_\star) = 0.51954\ldots \geq 0.51538\ldots = \frac{|G_\star|}{1 - 2s} \geq \frac{|G|}{1 - 2s}. \tag{4.15}$$

(iii) Because $\tau(C_3) = \tau \geq 6.96$, the condition that $\tau(C_3)\tilde{\varphi}_{t_\star}(\tau(C_1), \tau(C_2)) \geq 1$ is equivalent to

$$\tilde{\varphi}_{t_\star}(\tau) \geq \frac{1}{\tau},$$

which is true by our choice of $t_\star$ and τ; in particular, Claim 4.15 gives

$$\tilde{\varphi}_{t_\star}(\tau) \geq \tilde{\varphi}_{t_\star}(\tau_\star) = 0.14368\ldots \geq 0.14367\ldots = \frac{1}{\tau_\star} \geq \frac{1}{\tau}.$$

Thus Proposition 4.12 gives $(C - s) \cap C \cap (C + s) \neq \varnothing$ for all $s \in (0, 0.435]$; hence,

$$G^3_{AP}(C) \supset (0, 0.435].$$

Scaling back by d proves our theorem in full generality. □

Proof of Corollary 4.2. This follows from the proof above, though this time we apply Proposition 4.12 to the sets

$$C_1 = C + s\theta, \quad C_2 = C - s(1 - \theta), \quad \text{and} \quad C_3 = C,$$

where $s \in [0, 0.87]$ and $\theta \in (0, 1)$.

The minor changes that follow are that $Q = [s\theta, 1 - s(1 - \theta)]$ from which $|Q| = 1 - s$; this changes equation (4.15) to

$$\tilde{\psi}_{t_\star}(\tau) \geq \tilde{\psi}_{t_\star}(\tau_\star) = 0.51954\ldots \geq 0.51538\ldots = \frac{|G_\star|}{1 - s} \geq \frac{|G|}{1 - s},$$

from which the doubled range of s comes from. □

Lastly, we go one step beyond Theorem 4.1 to establish and quantify the existence of quadruple arithmetic progressions.

Proof of Theorem 4.3. To apply Proposition 4.13, we need only find a set C of sufficiently large thickness $\tau := \tau(C)$ such that

$$\tilde{\varphi}_t(\tau) \geq 1.$$

We claim that any Cantor set $C \subset \mathbb{R}$ with normalized length $\operatorname{conv}(C) = [0,1]$ and thickness

$$\tau := \tau(C) \geq \frac{1 - 0.015}{2(0.015)} = 32.83333\ldots$$

is sufficient.

Consider the anchor thickness $\tau_\star := 32.8$ and t-value $t_\star := 2.5$. Observe that

$$0 < t_\star < \frac{\tau_\star^2 - 2\tau_\star - 1}{3\tau_\star + 1} \leq \frac{\tau^2 - 2\tau - 1}{3\tau + 1},$$

and

$$0.78197\ldots = \tilde{\psi}_{t_\star}(\tau_\star) \leq \tilde{\psi}_{t_\star}(\tau) \quad \text{and} \quad 1.02129\ldots = \tilde{\varphi}_{t_\star}(\tau_\star) \leq \tilde{\varphi}_{t_\star}(\tau).$$

Thus, Proposition 4.13 applies and we conclude

$$[0, 0.3] \subset G^4_{AP}(C)$$

where the upper bound of the interval comes from

$$0.30498\ldots = \frac{\tilde{\psi}_{t_\star}(\tau_\star)}{1 + 2\tilde{\psi}_{t_\star}(\tau_\star)} \leq \frac{\tilde{\psi}_{t_\star}(\tau)}{1 + 2\tilde{\psi}_{t_\star}(\tau)}.$$

□

5. Concluding remarks

These results exhibit two complementary mechanisms for progression scale sets. Self-similar first splitting creates recursively scaled blackout intervals, while thickness and thick-intersection arguments create intervals of admissible scales. The level-two computation shows that product geometry can improve a purely first-level estimate without requiring a full symbolic description of the Cantor set.

Several natural problems remain. The most immediate is to determine the first-split spectrum B_ε exactly, or to compute a tractable level-three outer approximation. This could lower the blackout threshold below $(4 - \sqrt{13})/3$. It would also be useful to optimize the numerical HKY parameters and to develop analogous recursive descriptions for longer or asymmetric patterns.

Appendix A. Endpoint certificate for the level-two calculation

This appendix gives a direct check of the six rows used in Proposition 3.8. Write

$$A = I_{00}, \qquad B = I_{01}, \qquad C = I_{10}, \qquad D = I_{11},$$

and identify each interval with its left endpoint

$$0, \qquad \lambda - \lambda^2, \qquad 1 - \lambda, \qquad 1 - \lambda^2,$$

respectively; all four intervals have length $L = \lambda^2$. For a triple of left endpoints (p, q, r), formula (3.1) gives the lower candidates

$$q - p - L, \qquad \frac{r - p - L}{2}, \qquad r - q - L, \qquad 0$$

and the upper candidates

$$q - p + L, \qquad \frac{r - p + L}{2}, \qquad r - q + L.$$

The following comparisons hold throughout $1/3 \leq \lambda < 1/2$.

(i) For (A, A, C), the maximum lower candidate is $c_\lambda = 1 - \lambda - \lambda^2$ and the minimum upper candidate is λ^2. The interval is nonempty exactly when $c_\lambda \leq \lambda^2$, equivalently $\lambda \geq 1/2$.
(ii) For (A, A, D), the endpoints are $1-2\lambda^2$ and λ^2. Nonemptiness is equivalent to $\lambda \geq 1/\sqrt{3}$, so this row is empty in the range under study.
(iii) For (A, B, C), the lower endpoint is $\max\{\varepsilon, a_\lambda\}$ and the upper endpoint is $\min\{\lambda, b_\lambda\}$. The simultaneous equalities $a_\lambda = \varepsilon$ and $b_\lambda = \lambda$ reduce to $\lambda^2 - 3\lambda + 1 = 0$, whose root in $(0, 1/2)$ is $\rho = (3 - \sqrt{5})/2$. Hence this row contributes $[\varepsilon, \lambda]$ for $\lambda \leq \rho$ and $[a_\lambda, b_\lambda]$ for $\lambda \geq \rho$.
(iv) For (A, B, D), the maximum lower candidate is c_λ and the minimum upper candidate is λ. Thus the contribution is $[c_\lambda, \lambda]$, nonempty exactly when $1 - 2\lambda - \lambda^2 \leq 0$, or $\lambda \geq \eta = \sqrt{2} - 1$.
(v) For (B, B, C), the maximum lower candidate is $\varepsilon = 1 - 2\lambda$ and the minimum upper candidate is λ^2. Hence the contribution is $[\varepsilon, \lambda^2]$, again nonempty exactly when $\lambda \geq \eta$.
(vi) For (B, B, D), the calculation is the same as in (i), and there is no contribution for $\lambda < 1/2$.

It remains only to merge the three possible nonempty intervals. For $\rho < \lambda < \eta$, only $[a_\lambda, b_\lambda]$ remains. For $\eta \leq \lambda < \kappa$, their order is

$$\varepsilon \leq \lambda^2 < a_\lambda \leq b_\lambda < c_\lambda \leq \lambda.$$

The two strict gaps close simultaneously when

$$\lambda^2 = a_\lambda \quad \Longleftrightarrow \quad b_\lambda = c_\lambda \quad \Longleftrightarrow \quad 3\lambda^2 + \lambda - 1 = 0,$$

namely at $\kappa = (\sqrt{13} - 1)/6$. For $\lambda \geq \kappa$ the three intervals overlap and fill $[\varepsilon, \lambda]$. This independently reproduces all four cases of Proposition 3.8.

Acknowledgements

A. Y. is supported in part by the Natural Sciences and Engineering Research Council of Canada, NSERC (GR030571 and GR030540). S.S. is supported in part by NSF DMS-2231565 K.T. is supported in part by the Simons Foundation Grant GR137264.

References

[1] K. J. Falconer, *On the Hausdorff dimensions of distance sets*, Mathematika **32** (1985), no. 2, 206–212.
[2] P. Mattila and P. Sjölin, *Regularity of distance measures and sets*, Math. Nachr. **204** (1999), 157–162.
[3] T. Keleti, *Construction of one-dimensional subsets of the reals not containing similar copies of given patterns*, Anal. PDE **1** (2008), no. 1, 29–33.
[4] A. Yavicoli, *Large sets avoiding linear patterns*, Proc. Amer. Math. Soc. **149** (2021), no. 10, 4057–4066.
[5] I. Łaba and M. Pramanik, *Arithmetic progressions in sets of fractional dimension*, Geom. Funct. Anal. **19** (2009), no. 2, 429–456.
[6] M. Carnovale and S. Senger, *On the number of 3APs in fractal sets*, preprint, arXiv:2602.03029 (2026).
[7] P. Shmerkin, *Salem sets with no arithmetic progressions*, Int. Math. Res. Not. IMRN (2017), no. 7, 1929–1941.
[8] R. Broderick, L. Fishman, and D. Simmons, *Quantitative results using variants of Schmidt's game: dimension bounds, arithmetic progressions, and more*, Acta Arith. **188** (2019), no. 3, 289–316.

[9] J. Chaika, *Arithmetic progressions in middle Nth Cantor sets*, arXiv:1703.08998 (2017).
[10] L. Xi, K. Jiang, and Q. Pei, *Arithmetic progressions in self-similar sets*, Front. Math. China **14** (2019), no. 5, 957–966.
[11] Z. Tian, M. Lou, and L. Shang, *Arithmetic progressions and thickness properties in self-similar Cantor sets*, Discrete Contin. Dyn. Syst. Ser. S **25** (2026), 160–175.
[12] B. R. Hunt, I. Kan, and J. A. Yorke, *When Cantor sets intersect thickly*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 869–888.
[13] S. E. Newhouse, *Nondensity of axiom $A(a)$ on S^2*, in Global Analysis (Proc. Sympos. Pure Math., Vol. XIV), Amer. Math. Soc., Providence, RI, 1970, pp. 191–202.
[14] K. Taylor and S. Sandberg-Clark, *Triangles in the plane and arithmetic progressions in thick compact subsets of $\mathbb{R}^d$*, Canad. J. Math. **78** (2026), no. 3, 946–986.
[15] R. F. Williams, *How big is the intersection of two thick Cantor sets?*, in Continuum Theory and Dynamical Systems (Arcata, CA, 1989), Contemp. Math. **117**, Amer. Math. Soc., Providence, RI, 1991, pp. 163–175.
[16] A. Yavicoli, *Patterns in thick compact sets*, Israel J. Math. **244** (2021), no. 1, 95–126.
[17] A. Yavicoli, *A survey on Newhouse thickness, fractal intersections and patterns*, Math. Comput. Appl. **27** (2022), no. 6, Art. 111.

Department of Mathematical Sciences, United States Military Academy at West Point
Email address: samantha.sandbergclark@westpoint.edu

Department of Mathematics, The Ohio State University
Email address: taylor.2952@osu.edu

Department of Mathematics, The University of British Columbia
Email address: yavicoli@math.ubc.ca