# Cingulate-Precuneus Interactions: A New Locus of Dysfunction in Adult Attention-Deficit/ Hyperactivity Disorder

F. Xavier Castellanos, Daniel S. Margulies, Clare Kelly, Lucina Q. Uddin, Manely Ghaffari, Andrew Kirsch, David Shaw, Zarrar Shehzad, Adriana Di Martino, Bharat Biswal,

Edmund J. S. Sonuga-Barke, John Rotrosen, Lenard A. Adler, and Michael P. Milham

**Background:** Pathophysiologic models of attention-deficit/hyperactivity disorder (ADHD) have focused on frontal-striatal circuitry with alternative hypotheses relatively unexplored. On the basis of evidence that negative interactions between frontal foci involved in cognitive control and the non-goal-directed "default-mode" network prevent attentional lapses, we hypothesized abnormalities in functional connectivity of these circuits in ADHD.

**Methods:** Resting-state blood oxygen level– dependent functional magnetic resonance imaging (fMRI) scans were obtained at 3.0-Tesla in 20 adults with ADHD and 20 age- and sex-matched healthy volunteers.

**Results:** Examination of healthy control subjects verified presence of an antiphasic or negative relationship between activity in dorsal anterior cingulate cortex (centered at x = 8, y = 7, z = 38) and in default-mode network components. Group analyses revealed ADHD-related compromises in this relationship, with decreases in the functional connectivity between the anterior cingulate and precuneus/posterior cingulate cortex regions (p < .0004, corrected). Secondary analyses revealed an extensive pattern of ADHD-related decreases in connectivity between precuneus and other default-mode network components, including ventromedial prefrontal cortex ( $p < 3 \times 10^{-11}$ , corrected) and portions of posterior cingulate (p < .02, corrected).

**Conclusions:** Together with prior unbiased anatomic evidence of posterior volumetric abnormalities, our findings suggest that the long-range connections linking dorsal anterior cingulate to posterior cingulate and precuneus should be considered as a candidate locus of dysfunction in ADHD.

**Key Words:** ADHD, anterior cingulate cortex, default-mode network, functional magnetic resonance imaging, precuneus, posterior cingulate cortex

athophysiologic models of attention-deficit/hyperactivity disorder (ADHD) have focused on prefrontal-striatal and mesolimbic circuits (1) on the basis of findings of executive and motivational dysfunction (2,3). However, as pointed out in a recent quantitative meta-analysis of functional imaging studies of ADHD (4), the focus on executive dysfunction and the frontal lobes has left alternative hypotheses unexplored. Additionally, standard task-based functional imaging studies depend on the precise specification of underlying neuropsychological deficits. Although the hypothesis that ADHD results from a primary inhibitory deficit was proposed a decade

From the Phyllis Green and Randolph Cöwen Institute for Pediatric Neuroscience (FXC, DSM, CK, LQU, MG, ZS, ADM, EJSS-B, MPM), New York University Child Study Center; Nathan Kline Institute for Psychiatric Research (FXC), Orangeburg, New York; Department of Psychiatry (AK, DS, JR, LAA), New York University School of Medicine, New York; Department of Radiology (BB), University of Medicine and Dentistry of New Jersey, Newark; Institute for Disorders of Impulse and Attention (EJSS-B), School of Psychology, University of Southampton, Southampton, and Social, Genetic, Developmental Psychiatry Research Centre (EJSS-B), Institute of Psychiatry, Kings College London, London, United Kingdom.

Address reprint requests to F. Xavier Castellanos, M.D., Phyllis Green and Randolph Cowen Institute for Pediatric Neuroscience, NYU Child Study Center, 215 Lexington Avenue, 14th Floor, New York, NY 10016; E-mail: castef01@nyumc.org.

Received April 4, 2007; revised May 31, 2007; accepted June 29, 2007.

ago (5), the accumulated evidence has been equivocal at best (1).

A complementary approach is to examine the neural substrates of ADHD-relevant behaviors, such as attentional lapses (6,7), and assess whether the underlying circuits are implicated in ADHD through analysis of the temporal correlations among distributed brain regions. This method of functional connectivity (8), first applied to mapping the motor cortical system (9), provides remarkably detailed spatial maps of putatively functionally related regions (10).

Momentary lapses in attention have been associated with failure to suppress activity in the default-mode network (11). First identified by Raichle *et al.* (12), the default-mode network is a large and robustly replicable network (13–15) that comprises ventral medial prefrontal cortex (VMPFC) and posterior cingulate cortex (PCC) and precuneus in the midline and that has been linked to non-goal-directed processes (16). By definition, default-mode network activity can be detected during resting state scans, and abnormalities in this network have been reported in ADHD (17,18), Alzheimer's disease (19,20), autism (21), schizophrenia (22), and depression (23).

The starting point for this study was the finding that decreased cue-related activation in three prefrontal regions predicted slower response times and decreased default-mode network suppression in healthy adult volunteers (11), suggesting potential control loci for default-mode regulation. We hypothesized that these fronto-default mode interactions may represent loci of dysfunction in ADHD. Given reports that fronto-default-mode interactions are intrinsically represented in spontaneous activity at rest (14,24), we tested for the presence of ADHD-related

Table 1. Demographics

	ADHD Subjects	Healthy Comparison Subjects
Total Subjects	20	20
Mean ± SD Age (years)	$34.9 \pm 9.9$	$31.2 \pm 9.0$
Number (%) Male	16 (80%)	14 (70%)
Years of Education Completed	$17.2 \pm 1.9$	$17.6 \pm 1.6$
Inattentive Symptoms <sup>a</sup>	$7.4 \pm 1.6$	$1.2 \pm 0.4$
Hyperactive/Impulsive		
Symptoms <sup>a</sup>	$5.9 \pm 2.6$	$1.5 \pm 0.6$
Number (%) Treated	9 (45%)	0

ADHD, attention-deficit/hyperactivity disorder.

<sup>a</sup>Current DSM-IV inattention and hyperactivity/impulsivity symptoms (lifetime symptoms for healthy comparisons) were assessed using the Adult ADHD Clinical Diagnostic Scale 2.1. Symptom data on this scale were unavailable for two subjects diagnosed with combined-type ADHD by clinical interview. Age, sex distributions, and educational attainment did not differ significantly (p > .20).

differences in functional connectivity between each of the three previously identified frontal foci (dorsal anterior cingulate cortex [dACC]; right inferior frontal gyrus [rIFG]; right middle frontal gyrus [rMFG]) and the default-mode network.

### **Methods and Materials**

### **Participants**

Twenty adult participants with ADHD were recruited from the New York University (NYU) School of Medicine Adult ADHD Program (directed by author LA). Twenty age-matched comparisons were recruited through local media advertisements. All prospective participants were screened with the Symptom Checklist-90-Revised (SCL-90-R) to exclude a broad range of psychiatric psychopathology (25). Exclusion criteria for both groups included the following:1) lifetime history of psychotic, bipolar, or substance use disorders; 2) current history of mood, psychotic, anxiety, or substance use disorders; 3) lifetime history of treatment with psychotropics other than stimulants (for ADHD group only); or 4) history of neurological or chronic medical illness. All participants were evaluated with the Adult ADHD Clinical Diagnostic Scale (ACDS) Version1.2 (26), a semistructured interview that probes for the presence, severity, and impairment associated with ADHD symptoms in childhood and adulthood and which has been validated with informant-only reports (27). All participants with ADHD met lifetime criteria for Combined Type ADHD; they were also administered the Structured Clinical Interview for DSM-IV (SCID) to rule out other Axis I comorbid diagnoses. Best estimate diagnoses for probands were established by LA. A psychiatrist (AK or MG) administered a semistructured clinical interview to healthy comparison subjects to rule out all Axis I psychiatric disorders, including presence of ADHD or learning disorders in childhood or adulthood. Educational attainment (level attained, e.g., high school graduate = 12; college graduate = 16) and strong right-handedness were obtained from self-report. Demographic characteristics are shown in Table 1. All participants reported graduating from high school and attending at least 1 year of college. Nine patients were currently being treated with stimulants, which were discontinued for at least 1 day before scanning. All participants provided signed informed consent as approved by the institutional review boards of NYU and the NYU School of Medicine. Participants were compensated.

Each participant underwent one resting state scan consisting of 197 contiguous whole-brain functional volumes using echo planar imaging on a Siemens 3.0-Tesla Allegra (repetition time = 2000 msec; echo time = 25 msec; flip angle = 90, 39 slices,  $matrix = 64 \times 64$ ; field of view = 192 mm; acquisition voxel size =  $3 \times 3 \times 3$  mm; 6.5 min). Participants were verbally instructed to relax and remain still with eyes open while the word "Relax" was centrally displayed. A high-resolution T1-weighted magnetization prepared gradient echo sequence was also obtained.

### Preprocessing


Initial image preprocessing, including motion correction, despiking, and slice time correction, was done using AFNI software (http://www.afni.nimh.nih.gov/afni). Subsequent preprocessing, including spatial filtering (full width at half maximum [FWHM] = 6 mm), temporal bandpass filtering, and spatial normalization, was performed using FSL software (http://www.fmrib.ox.ac. uk/fsl).

### **Study Specific Template Generation**

To optimize automated spatial normalization and avoid possible group differences related to registration errors (28), we created a study-specific template based on the approach employed in optimized voxel-based morphometry protocols (28). To create the template, all participants' brain images were 1) spatially normalized to Montréal Neurological Institute (MNI) space using a 12-parameter affine transformation in the FLIRT program (http://www.fmrib.ox.ac.uk/fsl), 2) spatially filtered (FWHM = 8 mm), and 3) averaged. Using this template, we recalculated the spatial transformation for each participant using a 12-parameter affine implemented in FLIRT. Repeating our analyses using the standard adult 152 brain MNI template rather than the study specific template yielded nearly identical results.

# **Functional Connectivity Analyses**

To obtain time series for each seed in each participant, we 1) transform the subject's time series into MNI space using a 12 degrees of freedom linear affine implemented in FLIRT (voxel size =  $1 \times 1 \times 1$  mm) and 2) calculate the mean time series for each of the three mask spheres centered around the three spherical regions of interest (ROIs; 123 voxels, radius = 3.5 mm) centered at the coordinates reported by Weissman et al. (2006) converted to MNI space (dACC: x = 8, y = 7, z = 38; rIFG: x =34, y = 45, z = 23; rMFG: x = 49, y = 19, z = 0). For each ROI, individual participant analyses were carried out using the GLM implemented in FEAT, using a seed-based regression approach employed in our prior work (10), with the time series for the ROI, as well as for the nuisance covariates (time series regressors for global signal intensity, white matter, cerebrospinal fluid, and six motion parameters) as predictors. (See Figure 1 for schematic outline). We produced individual subject-level maps of all positively and negatively predicted voxels for each regressor, correcting for multiple comparisons at the cluster level using Gaussian random field theory (min Z > 2.3; cluster significance: p < .05, corrected). Group-level analyses (within and between) were conducted using a mixed-effects model (FLAME) implemented in FSL, which produced thresholded Z score maps of activity associated with each ROI. On the basis of the assumption that cortical connectivity should be reciprocal (29), we used the time series corresponding to the region that differed significantly between groups in our primary analyses and derived the corre-



**Figure 1.** Data analytical path for preprocessing, extraction of region of interest and nuisance covariate time series, individual subject multiple regression analyses, and mixed effects analyses of group results.

sponding positively and negatively predicted networks and their significant group differences.

Exploratory correlations between inattention and hyperactivity/impulsivity symptom severity reports were performed with

**Table 2.** ADHD-Related Decreases in Functional Connectivity with ACC (MNI coordinates x=8, y=7, z=38)

Region <sup>a</sup>	ВА	х	у	z	Z Score
Posterior Cingulate Gyrus	31	15	-56	28	4.42
Precuneus	7	-8	-63	38	4.14
Precuneus	7	22	-59	28	4.05
Precuneus	7	1	-80	41	3.83
Precuneus	7	4	-66	37	3.5
Precuneus	7	22	-60	37	3.37

ACC, anterior cingulate cortex; ADHD, attention-deficit/hyperactivity disorder; BA, Brodmann's area.


<sup>a</sup>Peak activations reported for cluster identified in precuneus (x = 4, y = -65, z = 34;  $p = 3.7 \times 10^{-4}$ , corrected; cluster size = 3661). All table coordinates in Talairach space.

the regression parameter estimates for the significant between group difference at a two-sided alpha of .05.

### **Results**

Although both rIFG and rMFG ROIs were significantly negatively related to precuneus and PCC in both groups (see Supplement 1), these relationships did not differentiate the groups significantly. By contrast, functional connectivity analyses of the dACC ROI demonstrated significantly less negatively correlated activity (p < .0004, corrected; 3989 cubic voxels 1.5 mm per side = 13,436 mm³) in precuneus/PCC in subjects with ADHD (see Figure 2 and Table 2).

As shown in Figure 3, the precuneus/PCC mask obtained from the dACC ROI-based group difference robustly predicted positively and negatively correlated activity in healthy volunteers. The positively correlated regions coincided with the default-mode network in control subjects, but the ADHD group lacked the anterior component of that network. As the right panel of Figure 3 shows, the groups differed significantly in medial prefrontal cortex and



**Figure 2. (A)** Functional connectivity analyses were carried out using a spherical region of interest (ROI) located in anterior cingulate cortex (ACC; diameter = 7 mm, number of voxels = 123; MNI coordinates x = 8, y = 7, z = 38), based on the findings of Weissman *et al.* (2006). **(B)** Voxels positively (red) and negatively (blue) predicted by the time series for the ACC ROI (sagittal slice: x = 6; coronal slice: y = -62). A robust negative or antiphasic relationship was noted between the ACC seed region and default-mode network components (i.e., an increase in ACC activity predicts a decrease in default-mode activity). Attention-deficit/hyperactivity disorder-related decreases in functional connectivity were noted between the ACC and precuneus.




Figure 3. (A) Secondary functional connectivity analyses were carried out using the posterior cingulate/precuneus cluster identified in our primary analyses as the seed region. (B) For each group (attention-deficit/hyperactivity disorder [ADHD] and control subjects), voxels positively (red) and negatively (blue) predicted by the time series for the seed region are indicated. These analyses provided further support for ADHD-related decreases in precuneus/anterior cingulate cortex (ACC) connectivity. Furthermore, they identified areas of ADHD-related decreases in connectivity among precuneus and other default-mode network components, including ventromedial prefrontal cortex and anterior portions of posterior cingulate cortex.

superior frontal gyrus ( $p < 3 \times 10^{-11}$ , corrected) and also in PCC/precuneus (p < .02, corrected; see Table 3). The ADHD group also showed significantly less negatively correlated functional activity in ACC (Brodmann's area [BA] 24, 31, 32) and medial frontal gyrus (BA 6) (p < .00002, corrected) and in superior temporal gyrus (BA 22, 41; p < .04, corrected), including claustrum and precentral gyrus (BA 43, 44). Figure 4 shows the distribution of regression parameter estimates for the negatively correlated relationship between dACC ROI and PCC/precuneus in the two groups (means  $\pm$ 


SD  $-.23 \pm 0.12$  and  $-.01 \pm .15$  for healthy control and ADHD groups, respectively; t(38) = 5.10,  $p = 9.6 \times 10^{-6}$ ; d = 1.61, 95% confidence interval .79-2.43). The regression parameter estimates were significantly negatively correlated with ADHD symptoms across the entire sample (Pearson Correlation Coefficients ranged between -.61 and -.54 (n = 38, p < .001) for the childhood or adulthood symptoms of hyperactivity/impulsivity or inattention. However, despite our prediction, we did not detect significant relationships between the parameter estimates and severity of

Table 3. ADHD-Related Decreases in Negatively and Positively Correlated Functional Connectivity with Precuneus/Posterior Cingulate Regions

Region	BA	Х	у	Z	Cluster Size (No. Voxels)	Cluster Significance	Peak Z Scores
			Negatively	Correlate	ed Functional Connectivity		
Middle Frontal Gyrus <sup>a</sup>	6	14	-6	49	5201	$p = 1.5 \times 10^{-5}$ , corrected	_
Cingulate Gyrus	24	9	3	37	_	· —	4.41
Cingulate Gyrus	24/32	9	6	37	_	_	4.17
Middle Frontal Gyrus	6	6	0	55	_	_	4.09
Cingulate Gyrus	32	9	10	37	<del>_</del>	_	4.08
Cingulate Gyrus	24	13	4	44	<del>_</del>	_	4.02
Cingulate Gyrus	31	19	-36	43	<del>_</del>	_	3.82
Superior Temporal Gyrus <sup>a</sup>	22	-47	-10	5	1704	p = .038, corrected	_
Superior Temporal Gyrus	41	-47	-23	10	_	<u> </u>	4.17
Claustrum	_	-32	-12	11	_	<del>_</del>	4.08
Claustrum	_	-38	-12	5	_	<del>_</del>	3.66
Superior Temporal Gyrus	22	-53	3	1	_	_	3.44
Precentral Gyrus	43	-59	-8	10	_	<del>_</del>	3.38
Precentral Gyrus	44	-49	3	7	_	_	3.36
			Positively	Correlate	d Functional Connectivity		
Middle Frontal Gyrus <sup>a</sup>	9	-4	45	25	13,479	$p = 3 \times 10^{-11}$ , corrected	_
Superior Frontal Gyrus	9	-7	55	26	_	<del>_</del>	4.47
Superior Frontal Gyrus	10	-19	63	22	_	<del>_</del>	4.36
Superior Frontal Gyrus	10	-19	66	22	_	_	4.24
Middle Frontal Gyrus	9	-2	48	22	_	<del>_</del>	4.1
Middle Frontal Gyrus	9	-2	51	20	<del>_</del>	_	4.05
Middle Frontal Gyrus	6	31	18	53	_	<del>_</del>	4
Cingulate Gyrus <sup>a</sup>	31	0	-49	28	1971	p = .018, corrected	_
Posterior Cingulate	31	-4	-54	26	_	<del>_</del>	3.66
Precuneus	31	-7	-60	26	_	<del>_</del>	3.59
Cingulate Gyrus	31	6	-39	29	<del>_</del>	_	3.48
Posterior Cingulate	23	-5	-59	19	<del>_</del>	<del>_</del>	3.34
Cuneus	7	3	-63	31	<del>_</del>	_	3.33
Cingulate Gyrus	31	1	-29	37	<del>_</del>	_	3.18

ADHD, attention-deficit/hyperactivity disorder; BA, Brodmann's area.

<sup>&</sup>lt;sup>a</sup>Peak activations reported for cluster.



**Figure 4.** Decreased anterior cingulate cortex (ACC)/precuneus connectivity in attention-deficit/hyperactivity disorder (ADHD). The scatterplot depicts the mean parameter estimates for dorsal ACC connectivity (seed region: x=8, y=7, z=38) in the precuneus/posterior cingulate cortex (PCC) region found to exhibit ADHD-related decrease in antirelationship to the ACC. As depicted in the plot, spontaneous activity in ACC negatively predicted activity in precuneus/PCC for control subjects, but no such relationship was found in ADHD ( $p<9\times10^{-6}$ ).

inattentive symptoms in the ADHD group (or the control group), presumably reflecting the highly restricted ranges of symptom counts within each group.

## Discussion

By analyzing the intrinsic functional connectivity of putative frontal control loci in unbiased whole brain comparisons, we confirmed that the negative relationship posited by Weissman et al. (11) between control regions and the default-mode network is intrinsically represented in brain. Consistent with our hypothesis that ADHD is associated with abnormalities in fronto-defaultmode interactions implicated in preventing attentional lapses, we found ADHD-related decreases in functional connectivity between a dorsal ACC seed and posterior components of the default-mode network (i.e., precuneus and PCC). Iteratively using the precuneus/PCC region as a "seed" for an additional regression analysis revealed ADHD-related decreases in connectivity among components of the default-mode network, notably between precuneus/PCC and ventromedial prefrontal cortex. We did not find a specific relationship between functional connectivity measures and inattention symptoms within the ADHD group, perhaps because inattentive and hyperactive symptoms, both of which were significantly negatively correlated with the regression parameter estimates, were highly correlated in our combined-type ADHD sample (r = .68). Of note, these novel results fit with the three unbiased voxel-based morphometric findings of diminished volume (30,31) or decreased cortical thickness (32) in precuneus and PCC in ADHD.

Although not the primary focus of this study, our finding of ADHD-related decreases in connectivity between anterior and posterior default mode components may suggest a novel locus of dysfunction for working-memory deficits commonly observed in ADHD (33). Better performance on a working-memory task has been reported to be positively related to the strength of functional connectivity between anterior and posterior default-mode components (VMPFC and PCC) (34). In this light, our findings of ADHD-related decreases in functional connectivity among default-mode components and abnormal connectivity with control regions suggest a potential neural basis for a linkage between

working memory deficits and attentional fluctuations that will be addressed in future studies. Similarly, future studies combining pharmacologic manipulations and electrophysiologic methods or magnetoencephalography are required to move beyond functional connectivity to effective connectivity, the influence exerted by one neural system on another (8).

Our results do not confirm a prior report of greater functional connectivity in adolescents with ADHD between dACC and widespread regions including thalamus, cerebellum, insula, and pons (17). There are several possible explanations for our divergent findings. First, the ACC is functionally and structurally complex (35,10); we "seeded" an ACC subregion empirically linked to default-mode interactions, while the seed ROI used by Tian et al. (17) comprised the entire dorsal ACC. Second, Tian et al. did not differentiate the antiphase relationships ("anticorrelations") (14,24) that accounted for our findings. It is possible that undiagnosed learning disorders in the ADHD participants may have contributed to our finding, although the reasonably high level of educational attainment in all participants tends to diminish this concern. Finally, the studies differed with respect to age group (child, adult). In future studies, we plan to examine age effects and directly assess cognitive measures such as attentional lapses (6,7), working memory (33), response variability (36), and medication response.

In summary, we found strong evidence of disconnection between an anterior cingulate control region implicated in ADHD (32,37) and posterior components of the default-mode network (i.e., precuneus and PCC). In the context of increasing awareness of the complex role of precuneus and PCC in "highlevel integration between posterior association processes and anterior executive functions" (38, p. 578), our findings suggest that structural and functional circuits linking the dACC to precuneus and PCC may represent "small-world network" long-range connections (39) that should be considered as a candidate locus of dysfunction in ADHD.

This work was supported in part by grants provided to FXC by the Stavros S. Niarchos Foundation, National Institute of Mental Health (NIMH; Grant Nos. 5R21MH066393 and 5T32MH067763), the Leon Lowenstein Foundation, NARSAD (The Mental Health Research Association), and gifts from Linda and Richard Schaps, Jill and Bob Smith, and the Taubman Foundation.

Drs. Castellanos, Kelly, Uddin, Ghaffari, Kirsch, Di Martino, Biswal, Milham and Messrs. Margulies, Shaw, and Shehzad reported no biomedical financial interests or potential conflicts of interest.

Dr. Sonuga-Barke has potential conflicts of interest in relation to UCB Pharmaceuticals (consultancy, advisory board, and speaker honoraria), Janssen Cilag (consultancy, research grant, speaker honoraria), Shire Pharmaceuticals (advisory board), and Medice (speaker honoraria).

Dr. Rotrosen reports consulting income from Axonyx, Inc., and United BioSource Corp.; research support from Alkermes, Inc., National Institute on Drug Abuse, and NIMH; equity interests in Alcon, American Oriental Bioengineering, Amgen, Atrion, Avon Products, Biogen Idec, Biosite, Colgate-Palmolive, Cooper Companies, Cyberoptics, Dentsply International, Dov Pharmaceutical, Dow Chemical, Elan, Genentech, Gillette, ICOS, Invitrogen, Ivax, Kinetic Concepts, Lyondell Chemical, Mikron Infrared, Mylan Labs, Polymedica, Procter and Gamble, PDL Biopharma Inc., Respironics, St. Jude Medical, and Tyco; and no potential conflicts of interest.

Dr. Adler receives grant/research support from Abbot Labora-

tories, Cortex Pharmaceuticals, Bristol-Myers Squibb, Merck & Co, Novartis Pharmaceuticals, Pfizer, Shire, Eli Lilly, Ortho McNeil/Johnson and Johnson, New River Pharmaceuticals, Cephalon, Neurosearch. He is a member of speakers bureaus for Eli Lilly and Shire and serves on the advisory boards and as a consultant for Abbot Laboratories, Cortex Pharmaceuticals, Novartis Pharmaceuticals, Pfizer, Shire, Eli Lilly, Ortho McNeil// Johnson and Johnson, New River Pharmaceuticals, Cephalon, Merck, and Neurosearch.

Supplementary material cited in this article is available online.

- 1. Castellanos FX, Sonuga-Barke EJS, Milham MP, Tannock R (2006): Characterizing cognition in ADHD: Beyond executive dysfunction. Trends Cogn Sci 10:117-123.
- 2. Willcutt EG, Doyle AE, Nigg JT, Faraone SV, Pennington BF (2005): Validity of the executive function theory of attention-deficit/hyperactivity disorder: A meta-analytic review. Biol Psychiatry 57:1336-1346.
- 3. Sonuga-Barke EJS (2005): Causal models of ADHD: from common simple deficits to multiple developmental pathways. Biol Psychiatry 57:1231-
- 4. Dickstein SG, Bannon K, Castellanos FX, Milham MP (2006): The neural correlates of attention deficit hyperactivity disorder: An ALE meta-analysis. J Child Psychol Psychiatry 47:1051-1062.
- 5. Barkley RA (1997): Behavioral inhibition, sustained attention, and executive functions: Constructing a unifying theory of ADHD. Psychol Bull 121:65-94.
- 6. Castellanos FX, Sonuga-Barke EJS, Scheres A, Di Martino A, Hyde C, Walters JR (2005): Varieties of attention-deficit/hyperactivity disorderrelated intra-individual variability. Biol Psychiatry 57:1416-1423.
- 7. Sonuga-Barke EJ, Castellanos FX (2007): Spontaneous attentional fluctuations in impaired states and pathological conditions: A neurobiological hypothesis. Neurosci Biobehav Rev [Epub ahead of print March 12, 20071.
- 8. Friston KJ (1994): Functional and effective connectivity in neuroimaging: A synthesis, Hum Brain Mapp 2:56-78.
- 9. Biswal B, Yetkin FZ, Haughton VM, Hyde JS (1995): Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magn Reson Med 34:537-541.
- 10. Margulies DS, Kelly AMC, Uddin LQ, Biswal BB, Castellanos FX, Milham MP (2007): Mapping the functional connectivity of anterior cingulate cortex. Neuroimage [Epub ahead of print May 24, 2007].
- 11. Weissman DH, Roberts KC, Visscher KM, Woldorff MG (2006): The neural bases of momentary lapses in attention. Nat Neurosci 9:971-978.
- 12. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL (2001): A default mode of brain function. Proc Natl Acad Sci U S A 98:676-682.
- 13. Greicius MD, Menon V (2004): Default-mode activity during a passive sensory task: Uncoupled from deactivation but impacting activation. J Cogn Neurosci 16:1484 –1492.
- 14. Fransson P (2005): Spontaneous low-frequency BOLD signal fluctuations: An fMRI investigation of the resting-state default mode of brain function hypothesis. Hum Brain Mapp 26:15-29.
- 15. Damoiseaux JS, Rombouts SA, Barkhof F, Scheltens P, Stam CJ, Smith SM, et al. (2006): Consistent resting-state networks across healthy subjects. Proc Natl Acad Sci U S A 103:13848-13854.
- 16. Raichle ME, Snyder AZ (2007): A default mode of brain function: A brief history of an evolving idea. Neuroimage [Epub ahead of print March 3, 2007].
- 17. Tian L, Jiang T, Wang Y, Zang Y, He Y, Liang M, et al. (2006): Altered resting-state functional connectivity patterns of anterior cingulate cortex in adolescents with attention deficit hyperactivity disorder. Neurosci Lett 400:39-43.
- 18. Zang YF, Yong H, Chao-Zhe Z, Qing-Jiu C, Man-Qiu S, Meng L, et al. (2007): Altered baseline brain activity in children with ADHD revealed by resting-state functional MRI. Brain Dev 29:83-91.

- 19. Greicius MD, Srivastava G, Reiss AL, Menon V (2004): Default-mode network activity distinguishes Alzheimer's disease from healthy aging: Evidence from functional MRI. Proc Natl Acad Sci U S A 101:4637–4642.
- 20. Buckner RL, Snyder AZ, Shannon BJ, LaRossa G, Sachs R, Fotenos AF, et al. (2005): Molecular, structural, and functional characterization of Alzheimer's disease: Evidence for a relationship between default activity, amyloid, and memory. J Neurosci 25:7709-7717.
- 21. Kennedy DP, Redcay E, Courchesne E (2006): Failing to deactivate: Resting functional abnormalities in autism. Proc Natl Acad Sci U S A 103: 8275-8280.
- 22. Liang M, Zhou Y, Jiang T, Liu Z, Tian L, Liu H, et al. (2006): Widespread functional disconnectivity in schizophrenia with resting-state functional magnetic resonance imaging. NeuroReport 17:209-213.
- 23. Greicius MD, Flores BH, Menon V, Glover GH, Solvason HB, Kenna H, et al. (2007): Resting-state functional connectivity in major depression: Abnormally increased contributions from subgenual cingulate cortex and thalamus. Biol Psychiatry [Epub ahead of print January 8, 2007].
- 24. Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van E, Raichle ME (2005): The human brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc Natl Acad Sci USA 102:9673-9678.
- 25. Deragotis LR (1986): Symptom Checklist-90-R: Administration, Scoring, and Procedures Manual, 3rd ed. Minneapolis: National Computer Systems, Inc.
- 26. Adler LA, Cohen J (2004): Diagnosis and evaluation of adult ADHD. Psychiatr Clin North Am 27:187-201.
- 27. Kessler RC, Adler L, Barkley R, Biederman J, Conners CK, Demler O, et al. (2006): The prevalence and correlates of adult ADHD in the United States: Results from the National Comorbidity Survey Replication. Am J Psychiatry 163:716-723.
- 28. Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS (2001): A voxel-based morphometric study of ageing in 465 normal adult human brains. Neuroimage 14:21-36.
- 29. Sporns O, Tononi G, Edelman GM (2000): Connectivity and complexity: The relationship between neuroanatomy and brain dynamics. Neural Netw 13:909-922.
- 30. Overmeyer S, Bullmore ET, Suckling J, Simmons A, Williams SCR, Santosh PJ, et al. (2001): Distributed grey and white matter deficits in hyperkinetic disorder: MRI evidence for anatomical abnormality in an attentional network. Psychol Med 31:1425-1435.
- 31. Carmona S, Vilarroya O, Bielsa A, Tremols V, Soliva JC, Rovira M, et al. (2005): Global and regional gray matter reductions in ADHD: A voxelbased morphometric study. Neurosci Lett 389:88-93.
- 32. Makris N, Biederman J, Valera EM, Bush G, Kaiser J, Kennedy DN, et al. (2007): Cortical thinning of the attention and executive function networks in adults with attention-deficit/hyperactivity disorder. Cereb Cortex 17:1364-1375.
- 33. Martinussen R, Hayden J, Hogg-Johnson S, Tannock R (2005): A metaanalysis of working memory impairments in children with attentiondeficit/hyperactivity disorder. J Am Acad Child Adolesc Psychiatry 44: 377-384.
- 34. Hampson M, Driesen NR, Skudlarski P, Gore JC, Constable RT (2006): Brain connectivity related to working memory performance. J Neurosci 26:13338-13343.
- 35. Bush G, Luu P, Posner MI (2000): Cognitive and emotional influences in anterior cingulate cortex. Trends Cogn Sci 4:215-222.
- 36. Simmonds DJ, Fotedar SG, Suskauer SJ, Pekar JJ, Denckla MB, Mostofsky SH (2007): Functional brain correlates of response time variability in children. Neuropsychologia [Epub ahead of print January 21, 2007].
- 37. Shaw P, Lerch J, Greenstein D, Sharp W, Clasen L, Evans A, et al. (2006): Longitudinal mapping of cortical thickness and clinical outcome in children and adolescents with attention-deficit/hyperactivity disorder. Arch Gen Psychiatry 63:540-549.
- 38. Cavanna AE, Trimble MR (2006): The precuneus: A review of its functional anatomy and behavioural correlates. Brain 129:564-583.
- 39. Bassett DS, Bullmore E (2006): Small-world brain networks. Neuroscientist 12:512-523.