

Abstract Writing for Medical Research Papers

医学论著英语摘要写作

Qi Hui, Chen Feina, Guo Haiyan

Fujian Medical University

Department of Arts and Sciences

译者序

本书是由福建医科大学文理艺术学院的齐晖、陈菲娜、郭海燕老师为主编，陈晶为学术秘书，交由复旦大学出版社出版，专供福医大学生使用的英语摘要写作教科书。我本人也是三位老师的学生，纵然课堂生动有趣、干货满满，但苦于同校前辈制作的扫描件观感不佳，笔记整理不便，译者决心要进行文字重排处理。其中自觉原书排版不善之处，皆进行重新编排，以符合译者审美。

原书通本以英文编写，编者似乎意图借此提升我等英语阅读水平，奈何文本中穿插语言学专有名词，初学时疲于翻译、苦不堪言。此外，期末复习期间，全英文本并不利于提升复习效率，故译者对主要文本进行翻译，对照复习。本套重置本将基于该译本进行整理，包含三种排版样式——原文重排版、双语对照版、译文版。以上三个版本请学弟学妹们按需取用，20届学长祝各位期末考试顺利。

本书为个人翻译作品，若发现纰漏，请在GitHub上提交issue。在此声明，此书仅供学习交流使用，请勿用于商业用途或其他领域。

neurocane

2023/7/7

编译环境

- 操作系统: Windows
- 语言: \LaTeX
- 编译环境: \XeLaTeX
- **TeX Live**版本: TeX Live 2022

目录

第一章 Overview of Abstracts	1
1.1 Definition of an Abstract	1
1.2 Importance and Functions of an Abstract	1
1.3 Types of Abstracts	2
1.4 Types of Informative Abstracts	2
1.5 Glossary	7
chapter Exercice	8
第二章	11
第三章	13
第四章	15
第五章	17
第六章	19

第一章 Overview of Abstracts

1.1 Definition of an Abstract

The American Psychological Association (APA) Style (2010) states that an abstract is a brief, comprehensive summary of the content of an article. According to the American National Standards Institute (1979), an abstract is an abbreviated accurate representation of the content of a document, preferably prepared by its author(s) for publication with it. In general, an abstract is a concise, accurate and comprehensive statement of the content of an article. It is original rather than excerpted.

1.2 Importance and Functions of an Abstract

An abstract is a distinct genre, and to some extent plays a pivotal role in academic reading and writing. In the era of information explosion, an enormous number of new publications are produced in the academic community each day. There is no practical way for every reader to get access to every new article, or to read every new publication even if it is accessible. The abstracts published online, which are concise and comprehensive, can be obtained easily and quickly. Abstract reading, then, may be a useful starting point of any academic reading and writing. In this sense, an abstract is the most read part of an article.¹

An abstract has at least three functions (Huckin, 2001). First, it serves as a stand-alone mini text, giving readers a quick summary of a study's objectives, methodology, findings and conclusions, which are the major components of abstracts. Second, it serves as a screening device, and gives readers an adequate view on whether the full-length article is of great value to their needs and worth further reading. A good abstract, to some extent, increases the chance of being cited or referenced. Third, for those readers who do opt to read the article as a whole, the abstract serves as a preview, creating an interpretive frame that can guide reading.

¹ 劳伦衣普桑，认为将指点效
则机，最你更枝。想极整月正
进好志次回总般，段然取向使
张规军证回，世市总李率英茹
持伴。

1.3 Types of Abstracts

Generally, abstracts fall into two categories, indicative and informative, depending on the type of information they convey. A typical distinction between them is that the indicative abstract, viewed as the outline of the paper, is usually shorter and simpler, while the informative abstract, viewed as the summary of the paper is usually longer and more thorough.

These two types of abstracts also differ in the components they contain. Indicative abstracts often include the purpose, scope, and methods of the report or study, but seldom include the results or conclusions. Reading indicative abstracts could not substitute reading the paper, because not all the crucial components are covered. It is more widely used in social science papers. On the other hand, informative abstracts usually include all the crucial components of the study, such as the background, purpose, methods, results, and conclusions. It is the type of abstracts widely used in medical field. In this book, we focus on the writing of informative abstracts. The abstracts referred to in the following chapters are informative abstracts.

1.4 Types of Informative Abstracts

There are two types of informative abstracts, structured abstracts and unstructured abstracts.

Where a heading or label is used at the beginning of the text in each section, it is a structured abstract. Each section is usually written in a separate paragraph, but sometimes sections are written in a sole or continuous paragraph. Headings might be background, objectives, methods, results, conclusions, and so on. They vary according to the criteria set by different journals. Structured abstracts appear to be favored by medically-relevant publications.

Where no heading or label is used to indicate different parts of an abstract, it is an unstructured abstract. It is always a sole paragraph. The major difference between the two types of abstracts lies in whether there are headings or not. In an unstructured abstract the content and sequence of the items are written as it is in the structured one.

Journals mandate which style should be used, so check the author guidelines if you're not sure. If it is not mentioned, keep an eye out for the type of abstracts preferable in the journals where you are willing to have your paper submitted and published. Write your abstracts in the style which dominates.

Sample 1.4.1:**BACKGROUND**

In patients with acute heart failure, early intervention with an intravenous vasodilator has been proposed as a therapeutic goal to reduce cardiac-wall stress and, potentially, myocardial injury, thereby favorably affecting patients' long-term prognosis.

METHODS

In this double-blind trial, we randomly assigned 2,157 patients with acute heart failure to receive a continuous intravenous infusion of either ularitide at a dose of 15 ng per kilogram of body weight per minute or matching placebo for 48 hours, in addition to accepted therapy. Treatment was initiated a median of 6 hours after the initial clinical evaluation. The coprimary outcomes were death from cardiovascular causes during a median follow-up of 15 months and a hierarchical composite end point that evaluated the initial 48-hour clinical course.

RESULTS

Death from cardiovascular causes occurred in 236 patients in the ularitide group and 225 patients in the placebo group (21.7% vs. 21.0%; hazard ratio, 1.03; 96% confidence interval, 0.85 to 1.25; $P=0.75$). In the intention-to-treat analysis, there was no significant between-group difference with respect to the hierarchical composite outcome. The ularitide group had greater reductions in systolic blood pressure and in levels of N-terminal pro-brain natriuretic peptide than the placebo group. However, changes in cardiac troponin T levels during the infusion did not differ between the two groups in the 55% of patients with paired data.

CONCLUSIONS

In patients with acute heart failure, ularitide exerted favorable physiological effects (without affecting cardiac troponin levels), but short-term treatment did not affect a clinical composite end point or reduce long-term cardiovascular mortality.

—Effect of Ularitide on Cardiovascular Mortality in Acute Heart Failure.

New England Journal of Medicine (2017)

Sample 1.4.2:**OBJECTIVE**

To evaluate the association between the parameters of 24-hour multi-channel intraluminal impedance (MII)-pH monitoring and the symptoms or quality of life (QoL) in laryngopharyngeal reflux (LPR) patients.

DESIGN

Prospective cohort study without controls.

SETTING

University teaching hospital.

METHODS

Forty-five LPR patients were selected from subjects who underwent 24-hour MII-pH monitoring and were diagnosed with LPR from September 2014 to May 2015. Reflux Symptom Index (RSI), Health-related Quality of Life (HRQoL), Short Form 12 (SF-12) Survey questionnaires were surveyed. Spearman's correlation was used to analyse the association between the symptoms or QoL and 24-hour MII-pH monitoring.

RESULTS

Most parameters in 24-hour MII-pH monitoring showed weak or no correlation with RSI, HRQoL and SF-12. Only number of non-acid reflux events that reached the larynx and pharynx (LPR-non-acid) and number of total reflux events that reached the larynx and pharynx (LPR-total) parameters showed strong correlation with heartburn in RSI ($R=0.520$, $P < 0.001$, $R=0.478$, $P = 0.001$, respectively). Multiple regression analysis showed that there was only one significant regression coefficient between LPR-non-acid and voice/hoarseness portion of HRQoL ($b=1.719$, $P = 0.022$).

CONCLUSION

Most parameters of 24-hour MII-pH monitoring did not reflect subjective symptoms or QoL in patients with LPR.

—Association between 24-hour combined multichannel intraluminal impedance-pH monitoring and symptoms or quality of life in patients with laryngopharyngeal reflux.

Clinical Otolaryngology (2017)

Sample 1.4.3:

Due to the high incidence of recurrent squamous cell carcinoma of the head and neck and the toxicity profile of current salvage regimens, there is a need for tolerable and effective treatment options. We performed a retrospective matched case series to report our experience with recurrent high-risk patients who received capecitabine (CAP) therapy in the adjuvant setting after salvage therapy. The 5-year recurrence-free survival rates for the CAP and control cohorts were 54% (95% CI, 0.27%–0.75%) and 27% (95% CI, 0.09%–0.50%), respectively. Multivariable Cox modeling showed a significant improvement in recurrence-free survival in the CAP cohort (hazard ratio, 0.19; 95% CI, 0.04–0.92; $P = .0392$). While this was a retrospective analysis that could not control for all variables, these exploratory findings offer insights that may inform a prospective study to determine CAP efficacy.

—Capecitabine after Surgical Salvage in Recurrent Squamous Cell Carcinoma of Head and Neck.
Otolaryngology—Head & Neck Surgery (2017)

Note 1.4.1: Corpus used for this book

The data used and analyzed in this book are from a custom-built corpus with 1.15 million tokens of medical research article (RA) abstracts. The discipline of medicine is divided into 18 sub-disciplines, and RA abstracts from 2 to 3 leading journals are randomly retrieved in each sub-discipline with relatively similar number of texts for each sub-discipline (Table 1.1). The journals selected are all with relatively high impact factors.

表 1.1: Sub-disciplines and journals in each sub-discipline

Sub-discipline	Journal
Anesthesiology	<i>British Journal of Anaesthesia</i>
	<i>Anesthesiology</i>
	<i>Anesthesia and Analgesia</i>
Dermatology	<i>Journal of American Academy of Dermatology</i>
	<i>Giornale Italiano di Dermatologia e Venereologia</i>

Continued on next page

表 1.1: Sub-disciplines and journals in each sub-discipline (Continued)

Sub-discipline	Journal
Emergency Medicine	<i>Annals of Emergency Medicine</i> <i>Internal and Emergency Medicine</i> <i>Academic Emergency Medicine</i>
Geriatrics	<i>Neurobiology of Aging</i> <i>Aging Cell</i> <i>Age and Ageing</i>
Internal Medicine	<i>The New England Journal of Medicine</i> <i>The Lancet</i> <i>JAMA-Journal of the American Medical Association</i>
Medical Imaging	<i>The Journal of Nuclear Medicine</i> <i>Investigative Radiology</i> <i>Radiology</i>
Medical Laboratory	<i>Clinical Chemistry and Laboratory Medicine</i> <i>Clinical Biochemistry</i>
Neurology	<i>The Lancet Neurology</i> <i>Annals of Neurology</i>
Obstetrics and Gynecology	<i>Obstetrics & Gynecology</i> <i>American Journal of Obstetrics & Gynecology</i> <i>An International Journal of Obstetrics & Gynecology</i>
Oncology	<i>Journal of Clinical Oncology</i> <i>The lancet Oncology</i>
Ophthalmology	<i>Ophthalmology</i> <i>American Journal of Ophthalmology</i> <i>Archives of Ophthalmology</i>
Otolaryngology (ENT)	<i>Head & Neck</i> <i>Clinical Otolaryngology</i> <i>Otolaryngology—Head & Neck Surgery</i>
Pain Medicine	<i>The Clinical Journal of Pain</i> <i>Pain Medicine</i> <i>Regional Anesthesia and Pain Medicine</i>
Pediatrics	<i>Journal of the American academy of child & Adolescent psychiatry</i> <i>Pediatrics</i> <i>JAMA pediatrics</i>

Continued on next page

表 1.1: Sub-disciplines and journals in each sub-discipline (Continued)

Sub-discipline	Journal
Physical medicine and rehabilitation	<i>Neurorehabilitation and neural repair</i> <i>Journal of fluency disorders</i>
Psychiatry	<i>Molecular psychiatry</i> <i>The American journal of psychiatry</i> <i>JAMA psychiatry</i>
Sports medicine	<i>Medicine and Science in Sports and Exercise</i> <i>Sports Medicine</i> <i>The American Journal of Sports Medicine</i>
Surgery	<i>Annals of Surgery</i> <i>American Journal of Transplantation</i> <i>Journal of Neurology, Neurosurgery & Psychiatry</i>

Task 1.4.1: Corpus-based task

Can you build your own corpus with at least 100, 000 tokens?

Note 1.4.2: Corpus used for this book

In this book, "corpus-based tasks" are designed to enhance your ability to explore language realizations of medical RA abstracts with corpus approach. Most of these tasks might require the use of software such as AntConc, WordSmith, and so on.

1.5 Glossary

表 1.2: Glossary of Chapter 1

WORDS	MEANING	MEANING OR EXAMPLE
excerpt /'eksɜ:pt/	<i>v.</i> 摘录; 引用	If a long piece of writing or music is excerpted, short pieces from it are printed or played on their own.
genre /'ʒɑ:nrə/	<i>n.</i> 体裁	a particular type of art, writing, music etc, which has certain features that all examples of this type share.

Continued on next page

表 1.2: Glossary of Chapter 1 (Continued)

WORDS	MEANING	MEANING OR EXAMPLE
mandate /'mændert/	<i>v.</i> 授权; 强制执行; 委托办理	to tell someone that they must do a particular thing.
methodology /,meθə'dɒləʒi/	<i>n.</i> 方法学	a set of methods and principles used to perform a particular activity.
opt /ɒpt/	<i>v.</i> 选择; 挑选	to choose one thing or do one thing instead of another
pivotal /'pɪvətl/	<i>adj.</i> 关键性的; 核心的	more important than anything else in a situation or system.

Chapter Exercise

1. Identify whether the following abstracts are structured or unstructured and tell the reasons.

Abstract 1

Objectives: The aim of this study was to analyze changes in health care utilization and cost among a sample of highly impaired children and adolescents who sought a 3-week intensive interdisciplinary pain treatment (IPT).

Materials and Methods: Claims data from 7 statutory health insurance companies were analyzed for 65 children and adolescents who sought IIP at the German Paediatric Pain Centre. The annual health care utilization and cost were determined for the following 4 areas: outpatient care, inpatient care, medications, and remedies and aids. We analyzed the changes in resource utilization in the year before (pre_1 y) IPT and in the subsequent year (post_1 y).

Results: Within the first year after IPT, overall health care costs did not decrease significantly. However, the pattern of health care utilization changed. First, significantly more children and adolescents started outpatient psychotherapy ($P = 0.001$). Second, the number of hospitalized children decreased significantly from 1-year pre to 1-year post ($P = 0.001$). Accordingly, there were significantly fewer hospitalizations for primary chronic pain disorders at 1-year post ($P < 0.001$). The prescription of nonopioids, co-analgesics and opioids was significantly reduced from 1-year pre to 1-year post (all $P < 0.013$).

Discussion: The present results indicate that the health care costs of children and adolescents with severe chronic pain disorders do not significantly decrease 1 year after IPT; however, the treatment becomes more goal-focused. Differential diagnosis measures and nonindicated therapeutic interventions decreased, and more indicated interventions, such as psychotherapy, were used. Future research is needed to investigate the economic long-term changes after IPT.

—Health Care Utilization and Cost in Children and Adolescents with Chronic Pain: Analysis of Health Care Claims Data 1 Year Before and After Intensive Interdisciplinary Pain Treatment.

The Clinical Journal of Pain (2017)

Abstract 2

Previous studies of brain structure in Tourette syndrome (TS) have produced mixed results, and most had modest sample sizes. In the present multicenter study, we used structural magnetic resonance imaging (MRI) to compare 103 children and adolescents with TS to a well-matched group of 103 children without tics. We applied voxel-based morphometry methods to test gray matter (GM) and white matter (WM) volume differences between diagnostic groups, accounting for MRI scanner and sequence, age, sex and total GM+WM volume. The TS group demonstrated lower WM volume bilaterally in orbital and medial prefrontal cortex, and greater GM volume in posterior thalamus, hypothalamus and midbrain. These results demonstrate evidence for abnormal brain structure in children and youth with TS, consistent with and extending previous findings, and they point to new target regions and avenues of study in TS. For example, as orbital cortex is reciprocally connected with hypothalamus, structural abnormalities in these regions may relate to abnormal decision making, reinforcement learning or somatic processing in TS.

—Brain structure in pediatric Tourette syndrome.

Molecular Psychiatry (2017)

Abstract 3

Objective: To assess the feasibility of detecting signature volatile organic compounds in the breath of patients with oral squamous cell carcinoma.

Study Design: Prospective cohort pilot study.

Setting: University hospital.

Subjects and Methods: Using gas chromatography and mass spectrometry, emitted volatile organic compounds in the breath of patients before and after curative surgery (n=10) were compared with those of healthy subjects (n=4). It was hypothesized that certain volatile organic compounds disappear after surgical therapy. A characteristic signature of these compounds for diseased patients was compiled and validated.

Results: Breath analyses revealed 125 volatile organic compounds in patients with oral cancer. A signature of 8 compounds that were characteristic for patients with oral cancer could be detected: 3 from this group presented were absent after surgery.

Conclusion: The presented results confirmed the hypothesis of an absence of cancer-associated volatile organic compounds in the breath after therapy. In this pilot study, we proved the feasibility of this test approach. Further studies should be initiated to establish protocols for usage in a clinical setting.

—Volatile Organic Compounds in the Breath of Oral Squamous Cell
Carcinoma Patients: A Pilot Study.

Otolaryngology Head and Neck Surgery (2017)

第二章

第三章

第四章

第五章

第六章

