



Why Are Neutrons Important?

To turn neutrino physics into a precision science, we need to understand complex neutrino-nucleus interactions (specifically on Ar):

- Neutrons carry away a **large fraction**¹ of energy.
- Neutron final states are **model dependent**.
- Neutrons are **difficult to detect** in LAr.




Contributions from neutrons vary widely from event to event!


Figure from A. Friedland and W. Li (2019).

How Does Missing Energy Affect DUNE?


- Measurements of the CP phase³ are shifted by a large amount if neutrons are ignored!



- The fraction of total energy given to hadrons (when at least one hadron is a neutron) is shown as a function of incident neutrino energy in a CC-interaction.
- GENIE varies in the lower energy regime by a factor of 2!




Blip Classification





BLIP Classification will consist of at least the following two models:

- PointNet++** - The first network will take in an entire detector readout for an event and learn to semantically label blips (e.g. PointNet++⁵).
- BlipGraph** - The second network will learn to classify clusters at different scales (e.g. PointNet⁴, DynamicEdgeConv⁶).



Below are examples of different neutron capture gammas simulated in LArSoft.

- Each cluster is normalized so that it is centered at zero and extends from [-1,1] in each variable.
- ADC is normalized over all events.



Contrastive Learning


Symmetries -> Augmentations

There are several symmetries in our dataset due to physics:

- Translation** - Translations in (channel, tdc) are handled by normalizing each point cloud to the origin.
- Rotation** - Rotations about (channel, tdc) are implemented as part of the augmentations.
- Fluctuations** - Fluctuations can occur from many sources including the electron drift.
- Parity** - Flipping of the momentum of the particle along the (channel, tdc) directions.



Rotations/translations about ADC are not symmetries!




Using the *Normalized Temperature-scaled Cross Entropy Loss (NTXent)*

$$\ell_{i,j} = -\log \frac{\exp(\text{sim}(z_i, z_j)/\tau)}{\sum_{k=1}^{2N} \mathbb{1}_{[k \neq i]} \exp(\text{sim}(z_i, z_k)/\tau)}, \quad (1)$$



we push "like" clusters closer together in the embedding space and push all others away.

Training Confusion Matrix Class particle



BlipGraph Results

This early model rejects 99.7 % of all other blip types when imposing a 90% cut on the probability to being a 4.745 MeV gamma.



Software

All of the related software can be [found on github](#), which includes the following repositories:

- Arrakis/ArrakisND** - a set of packages for constructing training datasets from LArSoft and larnd-sim simulation.
- Blip** - machine learning and clustering tools including the SparseUNet and BlipGraph..

Acknowledgements

This material is based upon work supported by the Department of Energy National Nuclear Security Administration through the Nuclear Science and Security Consortium under Award Number(s) DE-NA0003996.

- [1] O. Ronneberger et. al., *U-Net: Convolutional Networks for Biomedical Image Segmentation*, arXiv:1505.04597, 2015.
- [2] C. Choy et. al., *4D Spatio-Temporal ConvNets: Minkowski Convolutional Neural Networks*, arXiv:1904.08755, 2019.
- [3] M. Ester et. al., *A density-based algorithm for discovering clusters in large spatial databases with noise*, 1996.
- [4] Eric Church, *LArSoft: A software package for liquid argon time projection drift chambers*, arXiv:1311.6774, 2013.
- [5] DUNE Collaboration, *Status of single phase and dual phase DUNE prototype detectors at CERN*, XXXIX International Conference on High Energy Physics, 2018.