

# Exploring Quasi-crystals and algebraic structures: linking crystal bases to semigroups and beyond

António Malheiro

(joint work with Alan J. Cain, Inês Rodrigues and Fátima Rodrigues)

Center for Mathematics and Applications (NOVA Math), NOVA University of Lisbon

NBSAN, University of Manchester

June 20, 2024

NOVAMATH  
CENTER FOR MATHEMATICS  
+ APPLICATIONS

fct

Fundação  
para a Ciência  
e a Tecnologia



REPÚBLICA PORTUGUESA  
CIÊNCIA, TECNOLOGIA  
E ENSINO SUPERIOR

This work is funded by national funds through the FCT - Fundação para a Ciência e a Tecnologia, I.P., under the scope of the projects UIDB/00297/2020 and UIDP/00297/2020 (Center for Mathematics and Applications).

# Motivation

## Plactic monoid

[Lascoux, Schützenberger '81]

- ▶ Young tableaux, Schensted insertion




- ▶ Knuth relations

$$acb \equiv cab, a \leq b < c$$

$$bac \equiv bca, a < b \leq c$$


- ▶ Crystals



## Hypoplactic monoid

[Krob, Thibon '97], [Novelli '00]

- ▶ Quasi-ribbon tableaux, Krob–Thibon insertion




- ▶ Knuth + quartic relations

$$cadb \equiv acbd, a \leq b < c \leq d$$

$$bdac \equiv dbca, a < b \leq c < d$$

- ▶ Quasi-crystals



# Crystals

## Definition

A **crystal** of type  $A_{n-1}$  is a non-empty set  $\mathcal{C}$  together with maps

$$\tilde{e}_i, \tilde{f}_i : \mathcal{C} \longrightarrow \mathcal{C} \sqcup \{\perp\} \quad (\text{Kashiwara operators})$$

$$\tilde{\varepsilon}_i, \tilde{\varphi}_i : \mathcal{C} \longrightarrow \mathbb{Z} \sqcup \{-\infty\} \quad (\text{length functions})$$

$$wt : \mathcal{C} \longrightarrow \mathbb{Z}^n \quad (\text{weight function})$$

for  $i \in I := \{1, \dots, n-1\}$ , satisfying the following:

**C1.** For any  $x, y \in \mathcal{C}$ ,  $\tilde{e}_i(x) = y$  iff  $x = \tilde{f}_i(y)$ , and in that case

$$wt(y) = wt(x) + \alpha_i, \quad \tilde{\varepsilon}_i(y) = \tilde{\varepsilon}_i(x) - 1, \quad \tilde{\varphi}_i(y) = \tilde{\varphi}_i(x) + 1$$

**C2.**  $\tilde{\varphi}_i(x) = \tilde{\varepsilon}_i(x) + \langle wt(x), \alpha_i \rangle$


**C3.**  $\tilde{\varepsilon}_i(x) = -\infty \Rightarrow \tilde{e}_i(x) = \tilde{f}_i(x) = \perp$ .

where  $\alpha_i = (0, \dots, 0, 1, -1, 0, \dots, 0)$ .

(This definition is generalized for other Cartan types)

# Crystals

- The **crystal graph** associated to a crystal  $\mathcal{C}$  is the directed weighted graph where  $y \xrightarrow{i} x$  iff  $\tilde{e}_i(x) = y$  iff  $\tilde{f}_i(y) = x$ .




# Crystals

- ▶ A crystal is **seminormal** if

$$\tilde{e}_i(x) = \max\{k : \tilde{e}_i(x)^k \neq \perp\}, \quad \tilde{\varphi}_i(x) = \max\{k : \tilde{f}_i(x)^k \neq \perp\},$$

for all  $i \in I$  and  $x \in \mathcal{C}$ . In particular,  $\tilde{e}_i(x), \tilde{\varphi}_i(x) \geq 0$ .



- ▶ To compute  $\tilde{f}_i(w)$  and  $\tilde{e}_i(w)$  on a word  $w \in \{1 < \dots < n\}^*$ :
  - ▶ consider the subword with only symbols  $i$  and  $i+1$ , and cancel all pairs  $(i+1)i$  ( $i$ -inversions), until there are no pairs left.
  - ▶  $\tilde{e}_i$  changes the *leftmost*  $i+1$  to  $i$ , if possible; if not, it is  $\perp$ .
  - ▶  $\tilde{f}_i$  changes the *rightmost*  $i$  to  $i+1$ , if possible; if not, it is  $\perp$ .

1221112


12**1**112

1**2****2**112

# Stembridge crystals

- ▶ A **Stembridge crystal** is a seminormal crystal of simply-laced type that satisfies some local axioms [Stembridge '03]. These are the crystal graphs that correspond to representations of Lie algebras.
- ▶ The connected components have nice properties:
  - ▶ Unique highest weight element (source vertex), from which all vertices can be reached.
  - ▶ All components whose highest weight elements have the same weight are isomorphic.
  - ▶ In type  $A$ , the character of a connected component is a Schur function  $s_\lambda$ .


# Stembridge crystals




# Stembridge crystals

## Local axioms

**S1.** If  $\tilde{e}_i(x) = y$ , then  $\tilde{e}_j(y)$  is equal to  $\tilde{e}_j(x)$  or  $\tilde{e}_j(x) + 1$  (the second case is possible only if  $|i - j| = 1$ ).



for  $|i - j| = 1$



for  $|i - j| > 1$






# Stembridge crystals

## Local axioms

- S2.** If  $\tilde{e}_i(x) = y$  and  $\tilde{\epsilon}_j(y) = \tilde{\epsilon}_j(x) > 0$  then

$$\tilde{e}_i \tilde{e}_j(x) = \tilde{e}_j \tilde{e}_i(x) \neq \perp$$




and  $\tilde{\varphi}_i(\tilde{e}_j(x)) = \tilde{\varphi}_i(x)$ .

(and dual axioms for  $\tilde{f}_i, \tilde{f}_j$ )

- S3.** If  $\tilde{e}_i(x) = y$  and  $\tilde{e}_j(x) = z$ , and  $\tilde{\epsilon}_i(z) = \tilde{\epsilon}_i(x) + 1$  and  $\tilde{\epsilon}_j(y) = \tilde{\epsilon}_j(x) + 1$  then

$$\tilde{e}_i \tilde{e}_j^2 \tilde{e}_i(x) = \tilde{e}_j \tilde{e}_i^2 \tilde{e}_j(x) \neq \perp.$$



# Motivation

## Plactic monoid

[Lascoux, Schützenberger '81]

- ▶ Young tableaux, Schensted insertion




- ▶ Knuth relations

$$acb \equiv cab, a \leq b < c$$

$$bac \equiv bca, a < b \leq c$$


- ▶ Crystals



## Hypoplactic monoid

[Krob, Thibon '97], [Novelli '00]

- ▶ Quasi-ribbon tableaux, Krob–Thibon insertion




- ▶ Knuth + quartic relations

$$cadb \equiv acbd, a \leq b < c \leq d$$

$$bdac \equiv dbca, a < b \leq c < d$$

- ▶ Quasi-crystals



# Quasi-crystals

- ▶ First introduced by Cain and M. (2017), providing another characterization of the hypoplactic monoid of type  $A$ .
- ▶ Cain, Guilherme and M. (2023) provided a definition of abstract quasi-crystals for other Cartan types.
- ▶ For type  $A$ , each connected component has a unique highest weight element, is isomorphic to a quasi-crystal of quasi-ribbon tableaux, and its character is a fundamental quasisymmetric function  $F_\alpha$ .
- ▶ We have  $s_\lambda = \sum_{T \in \text{SYT}(\lambda)} F_{\text{DesComp}(T)}$ .
- ▶ Using this decomposition, Maas-Gariépy (2023) independently introduced quasi-crystals, as subgraphs of a connected component of a crystal graph.

# Quasi-crystals

Definition (Cain, Guilherme, M. '23)

A **quasi-crystal** of type  $A_{n-1}$  is a non-empty set  $\mathcal{Q}$  together with maps

$$\ddot{e}_i, \ddot{f}_i : \mathcal{Q} \longrightarrow \mathcal{Q} \sqcup \{\perp\} \quad (\text{quasi-Kashiwara operators})$$

$$\ddot{e}_i, \ddot{\varphi}_i : \mathcal{Q} \longrightarrow \mathbb{Z} \sqcup \{-\infty, +\infty\}$$

$$wt : \mathcal{Q} \longrightarrow \mathbb{Z}^n$$

for  $i \in \{1, \dots, n-1\}$ , satisfying the same axioms of crystals and additionally:

$$\ddot{e}_i(x) = +\infty \Rightarrow \ddot{e}_i(x) = \ddot{f}_i(x) = \perp.$$

► A quasi-crystal is **seminormal** if, for all  $i \in I$  and  $x \in \mathcal{Q}$ ,

$$\ddot{e}_i(x) = \max\{k : \ddot{e}_i(x)^k \neq \perp\}, \quad \ddot{\varphi}_i(x) = \max\{k : \ddot{f}_i(x)^k \neq \perp\}$$

whenever  $\ddot{e}_i(x) \neq +\infty$ .

## Quasi-crystals

To compute  $\ddot{f}_i(w)$  and  $\ddot{e}_i(w)$  on a word  $w \in \{1 < \dots < n\}^*$ :

- ▶ If  $w$  has an  $i$ -inversion,  $\ddot{f}_i(w) = \ddot{e}_i(w) = \perp$ .
- ▶ Otherwise,  $\ddot{f}_i(w) = \tilde{f}_i(w)$  and  $\ddot{e}_i(w) = \tilde{e}_i(w)$ .

$$\ddot{f}_1(13121) =$$

$$\ddot{f}_1(13112) =$$

$$\ddot{f}_1(1\textcolor{lightgray}{3}121) =$$

$$\ddot{f}_1(13112) =$$

$$\ddot{f}_1(1\textcolor{lightgray}{3}1\textcolor{red}{2}1) =$$

$$\ddot{f}_1(13112) =$$


$$\ddot{f}_1(13121) = \perp$$

$$\ddot{f}_1(13112) =$$

# Quasi-crystals

The **quasi-crystal graph** associated to a quasi-crystal  $\mathcal{Q}$  is the directed weighted graph where:

- ▶  $y \xrightarrow{i} x$  iff  $\ddot{e}_i(x) = y$ .
- ▶  $x$  has an  $i$ -labelled loop iff  $\ddot{e}_i(x) = +\infty$  iff  $\ddot{\varphi}_i(x) = +\infty$ .




# Local characterization of quasi-crystals

## Local quasi-crystal axioms


**LQC1.** If  $\ddot{e}_i(x) = y$ , then:

- For  $|i - j| > 1$ ,  $\ddot{e}_j(x) = \ddot{e}_j(y)$ .




- For  $j = i + 1$ ,

$$\ddot{e}_{i+1}(x) \neq \ddot{e}_{i+1}(y) \Leftrightarrow (\ddot{e}_{i+1}(x) = +\infty \wedge \ddot{e}_i(y) = 0) \Rightarrow \ddot{e}_{i+1}(y) > 0.$$



- For  $j = i - 1$ ,


$$\ddot{\varphi}_{i-1}(x) \neq \ddot{\varphi}_{i-1}(y) \Leftrightarrow (\ddot{\varphi}_{i-1}(y) = +\infty \wedge \ddot{\varphi}_i(x) = 0) \Rightarrow \ddot{\varphi}_{i-1}(x) > 0.$$



# Local characterization of quasi-crystals


## Local quasi-crystal axioms

**LQC2.**  $\ddot{e}_i(x) = 0$  iff  $\ddot{\varphi}_{i+1}(x) = 0$ , for  $i \in \{1, \dots, n-2\}$ .



**LQC3.** If both  $\ddot{e}_i(x)$  and  $\ddot{e}_j(x)$  are defined, for  $i \neq j$ , then  $\ddot{e}_i \ddot{e}_j(x) = \ddot{e}_j \ddot{e}_i(x) \neq \perp$  (and dual axiom for  $\ddot{f}_i, \ddot{f}_j$ .)

$$\ddot{e}_i \ddot{e}_j(x) = \ddot{e}_j \ddot{e}_i(x)$$





# Local characterization of quasi-crystals

Theorem (Cain, M., Rodrigues, Rodrigues '23)

*If  $\mathcal{Q}$  is a quasi-crystal of type  $A$  (not necessarily seminormal) satisfying the local axioms, and such that  $\tilde{e}_i(x) \neq +\infty$  and  $\tilde{f}_i(x) \neq +\infty$ , for all  $i \in I, x \in \mathcal{Q}$ , then  $\mathcal{Q}$  is a weak Stembridge crystal (i.e. not necessarily seminormal).*

Theorem (Cain, M., Rodrigues, Rodrigues '23)

*Let  $\mathcal{Q}$  be a connected component of a seminormal quasi-crystal graph of type  $A$ , weighted in  $\mathbb{Z}_{\geq 0}^n$ , satisfying the local axioms. Then,  $\mathcal{Q}$  has a unique highest weight element, whose weight is a composition.*

Theorem (Cain, M., Rodrigues, Rodrigues '23)

*Let  $\mathcal{Q}$  and  $\mathcal{Q}'$  be connected components of seminormal quasi-crystal graphs of type  $A$  satisfying the local axioms, with highest weight elements  $u$  and  $v$ . If  $\text{wt}(u) = \text{wt}(v)$ , then there exists a weight-preserving isomorphism between  $\mathcal{Q}$  and  $\mathcal{Q}'$ .*

# Quasi-tensor product of quasi-crystals

Cain, Guilherme, and M. (2023) introduced a notion of quasi-tensor product of seminormal quasi-crystals, denoted  $\mathcal{Q} \ddot{\otimes} \mathcal{Q}'$ , which has  $\mathcal{Q} \times \mathcal{Q}'$  as underlying set and maps:

- ▶  $wt(x \ddot{\otimes} x') = wt(x) + wt(x')$ .
- ▶ If  $\ddot{\varphi}_i(x) > 0$  and  $\ddot{\varepsilon}_i(x') > 0$ ,  $\ddot{e}_i(x \ddot{\otimes} x') = \ddot{f}_i(x \ddot{\otimes} x') = \perp$  and  $\ddot{\varepsilon}_i(x \ddot{\otimes} x') = \ddot{\varphi}_i(x \ddot{\otimes} x') = +\infty$ , otherwise,

$$\ddot{e}_i(x \ddot{\otimes} x') = \begin{cases} \ddot{e}_i(x) \ddot{\otimes} x' & \text{if } \ddot{\varphi}_i(x) \geq \ddot{\varepsilon}_i(x') \\ x \ddot{\otimes} \ddot{e}_i(x') & \text{if } \ddot{\varphi}_i(x) < \ddot{\varepsilon}_i(x') \end{cases}$$

$$\ddot{f}_i(x \ddot{\otimes} x') = \begin{cases} \ddot{f}_i(x) \ddot{\otimes} x' & \text{if } \ddot{\varphi}_i(x) > \ddot{\varepsilon}_i(x') \\ x \ddot{\otimes} \ddot{f}_i(x') & \text{if } \ddot{\varphi}_i(x) \leq \ddot{\varepsilon}_i(x') \end{cases}$$

$$\ddot{\varepsilon}_i(x) = \max\{\ddot{\varepsilon}_i(x), \ddot{\varepsilon}_i(x') - \langle wt(x), \alpha_i \rangle\}$$

$$\ddot{\varphi}_i(x) = \max\{\ddot{\varphi}_i(x) + \langle wt(x'), \alpha_i \rangle, \ddot{\varphi}_i(x')\}$$


(With this convention  $x \ddot{\otimes} y$  is identified with the word  $yx$ .)

# Quasi-tensor product of quasi-crystals


- $\mathcal{B}_n$  is the standard crystal of type  $A_{n-1}$ :

$$1 \xrightarrow{\text{red}} 2 \xrightarrow{\text{blue}} 3 \xrightarrow{\text{green}} \dots \xrightarrow{\text{teal}} n$$

- Similarly to the case of the plactic monoid, each component of the hypoplactic monoid is isomorphic to some  $\mathcal{B}_n^{\ddot{\otimes} k}$ .



A connected component of  $\mathcal{B}_3^{\ddot{\otimes} \mathcal{B}_3}$




A connected component of  $\mathcal{B}_3 \otimes \mathcal{B}_3$

# Quasi-tensor product of quasi-crystals

Theorem (Cain, M., Rodrigues, Rodrigues '23)

*Let  $Q$  and  $Q'$  be seminormal quasi-crystal graphs satisfying the local axioms. Then,  $Q \otimes Q'$  is a seminormal quasi-crystal that satisfies the same axioms.*

- ▶ The standard crystal  $\mathcal{B}_n$  satisfies the local axioms.
- ▶ A connected component of a quasi-crystal of words, being isomorphic to some  $\mathcal{B}_n^{\otimes k}$ , satisfies the local axioms.
- ▶ As a consequence, every connected component of a seminormal quasi-crystal satisfying the local axioms is isomorphic a quasi-crystal of quasi-ribbon tableaux.



# From crystals to quasi-crystals

Let  $(\mathcal{C}, \tilde{f}_i, \tilde{e}_i, \tilde{\varepsilon}_i, \tilde{\varphi}_i)$  be a connected component of a Stembridge crystal, weighted in  $\mathbb{Z}_{\geq 0}^n$ , and define  $(\mathcal{Q}, \tilde{f}_i, \tilde{e}_i, \tilde{\varepsilon}_i, \tilde{\varphi}_i)$  to have the same underlying set as  $\mathcal{C}$  and:


- Place a  $i$ -labelled loop on  $x$  if  $\tilde{\varepsilon}_i(x) < wt_{i+1}(x)$ , for all  $i \in I, x \in \mathcal{C}$  (equivalently, if  $\tilde{\varphi}_i(x) < wt_i(x)$ ).
- Then, remove  $i$ -labelled edges that have  $i$ -labelled loops on both ends.

**Theorem (Cain, M., Rodrigues, Rodrigues '23)**

$\mathcal{Q}$  is a seminormal quasi-crystal that satisfies the local axioms.


**Corollary (Cain, M., Rodrigues, Rodrigues '23)**

If  $\mathcal{C}$  has highest weight  $\lambda$ , the number of connected components of  $\mathcal{Q}$  is given by the number of standard Young tableaux of shape  $\lambda$ .



$$s_{21} = F_{21} + F_{12}.$$

# From crystals to quasi-crystals



$$s_{211} = F_{211} + F_{121} + F_{112}$$

# Some references



A.J. Cain, R.P. Guilherme, A. Malheiro “Quasi-crystals for arbitrary root systems and associated generalizations of the hypoplactic monoid”. arXiv:2301.00271.



A.J. Cain, A. Malheiro “Crystallizing the hypoplactic monoid: from quasi-Kashiwara operators to the Robinson–Schensted–Knuth-type correspondence for quasi-ribbon tableaux”, J. Algebr. Comb. **45** (2), 475–524 (2017).



D. Krob, J.-Y. Thibon “Noncommutative symmetric functions. IV: Quantum linear groups and Hecke Algebras at  $q = 0$ ”, J. Algebr. Comb. **6** (4), 339 – 376 (1997).



J.-C. Novelli “On the hypoplactic monoid”, Discrete Math., **217** (1–3), 315–336 (2000).



J.R. Stembridge “A local characterization of simply-laced crystals”, Trans. Am. Math. Soc. **355** (12), 4807–4823 (2003).

Thank you!