

## Lecture 5: Vector spaces (class)

Admin: Reading:

<b>Matrix Analysis and Applied Linear Algebra</b> Carl D. Meyer	<b>Linear Algebra and Its Applications</b> Fourth Edition Gilbert Strang
<b>Contents</b>	<b>Contents</b>
<b>Preface</b> 1. Linear Equations 1.1 Elimination 1.2 Elementary Row Operations 1.3 Gaussian Method 1.4 LU Factorization 1.5 Computing Determinants 1.6 Inverse Matrices 1.7 Making Computations Work 1.8 Making Gaussian Elimination Work  2. Matrige Systems and Echelon Form 2.1 Row Echelon Form and Pivot 2.2 Row Operations 2.3 Consistency of Linear Systems 2.4 Rank 2.5 Homogeneous Systems  3. Matrix Algebra 3.1 From Ax = b to Ax = b 3.2 Matrix Multiplication 3.3 Powers of Matrices 3.4 Matrix Inverses 3.5 Matrix Factorizations 3.6 Matrix Functions  4. Vector Spaces 4.1 Vectors and Subspaces 4.2 Linear Independence 4.3 Basis and Dimension  LPP  Index  Preface Contents Index	<b>1 Matrices and Gaussian Elimination</b> 1.1 Introduction ..... 1.2 The Geometry of Linear Equations ..... 1.3 An Example of Gaussian Elimination ..... 1.4 Matrix Notation and Matrix Multiplication ..... 1.5 Triangular Factors and Row Exchanges ..... 1.6 Inverse Matrices and Determinants ..... 1.7 Special Matrices and Applications ..... Review Exercises .....  <b>2 Vector Spaces</b> 2.1 Vector Spaces and Subspaces ..... 2.2 Solving Ax = 0 and Ax = b ..... 2.3 Linear Transformations, Roots, and Eigenvectors ..... 2.4 Change of Basis, Similarity ..... 2.5 Graphs and Networks ..... 2.6 Linear Transformations ..... Review Exercises .....  <b>3 Orthogonality</b> 3.1 Orthogonal Vectors and Subspaces ..... 3.2 Cosines and Projections onto Lines ..... 3.3 Perpendicular Lines and Angles ..... 3.4 Orthogonal Bases and Gram-Schmidt ..... 3.5 The Fast Fourier Transform ..... Review Exercises .....  <b>4</b>

## VECTOR SPACES

Why? They're everywhere!

$$\mathbb{R}^4 \quad (a, b, c, d)$$

$$\text{cubic polynomials} \quad ax^3 + bx^2 + cx + d$$

$$2 \times 2 \text{ matrices} \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\text{multilinear polynomials in } x \text{ and } y \quad axy + bx + cy + d$$

:

So it makes sense to abstract their properties and study them together.

And, the best way to understand matrices is as linear transformations on vector spaces.

Main properties: "LINEARITY"

$$\begin{aligned} \text{Addition: } & (a, b, c, d) + (e, f, g, h) \\ &= (a+e, b+f, c+g, d+h) \\ & \begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} e & f \\ g & h \end{pmatrix} = \begin{pmatrix} a+e & b+f \\ c+g & d+h \end{pmatrix} \end{aligned}$$

$$(ax^3 + bx^2 + cx + d) + (ex^3 + fx^2 + gx + h) = (a+e)x^3 + (b+f)x^2 + \dots$$

Scalar multiplication:  $\mathbb{R}, \mathbb{C}, \mathbb{F}$

$$5(a, b, c, d) = (5a, 5b, 5c, 5d)$$

$$5 \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 5a & 5b \\ 5c & 5d \end{pmatrix}$$

$$5(ax^3 + bx^2 + cx + d) = \dots$$



These all behave the same way!

Note: Only under scalar multiplication, not arbitrary multiplication.

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} e & f \\ g & h \end{pmatrix} = \begin{pmatrix} ae+bg & af+bh \\ ce+dg & cf+dh \end{pmatrix}$$

$$(ax^3 + bx^2 + cx + d)(ex^3 + fx^2 + gx + h) = ae x^6 + (af+be)x^5 + \dots$$

$(a, b, c, d)(e, f, g, h)$  is not defined!

(there's just the dot product  $ae+bf+cg+dh$ )

(there's just the dot product  $a\mathbf{e} + b\mathbf{f} + c\mathbf{g} + d\mathbf{h}$ )

Definition: A **vector space** consists of

- a set of "vectors"  $V$
- a field  $\mathbb{F}$  (often the reals  $\mathbb{R}$  or complex #'s  $\mathbb{C}$ )
- operations of
  - vector addition  $V \times V \rightarrow V$ , denoted  $\vec{x} + \vec{y}$
  - scalar multiplication  $\mathbb{F} \times V \rightarrow V$ , denoted  $\alpha \vec{x}$

that satisfy:

- closure under addition & scalar multiplication:

for all  $\alpha \in \mathbb{F}$   
 $\vec{x}, \vec{y} \in V$

$$\vec{x} + \vec{y} \in V$$

$$\alpha \vec{x} \in V$$

interesting

- existence of  $\vec{0} \in V$

$$\vec{0} + \vec{x} = \vec{x} \text{ for all } \vec{x}$$

- additive inverses

for all  $\vec{x} \in V$ , there exists  $\vec{y} \in V$   
 s.t.  $\vec{x} + \vec{y} = \vec{0}$

for all  $\alpha, \beta \in \mathbb{F}$ ,  $\vec{x}, \vec{y}, \vec{z} \in V$ :

$$\vec{x} + \vec{y} = \vec{y} + \vec{x}$$

$$\vec{x} + (\vec{y} + \vec{z}) = (\vec{x} + \vec{y}) + \vec{z}$$

$$\alpha(\beta \vec{x}) = (\alpha \beta) \vec{x}$$

$$(\alpha + \beta) \vec{x} = \alpha \vec{x} + \beta \vec{x}$$

$$\alpha(\vec{x} + \vec{y}) = \alpha \vec{x} + \alpha \vec{y}$$

- $1\vec{x} = \vec{x}$  (identity for multiplication)

Note: The most important properties to check are closure under addition and scalar multiplication.

The other properties are usually automatic.

Examples: Vector spaces are everywhere!

①  $\mathbb{R}^n$ : real vectors  $(x_1, x_2, \dots, x_n)$   
 coordinate-wise addition & multiplication

②  $\mathbb{C}^n$  matrices  $\mathbb{R}^{m \times n}$  or  $\mathbb{C}^{m \times n}$

③ the single-point sets  $\{\vec{0}\}$  or  $\{(0, 0, \dots, 0)\}$   
 (trivially closed under addition & multiplication)

But these are NOT vector spaces:

$$\{(1, 0, 0)\}$$

$$\{(0, 0), (1, 0)\}$$

the interval  $[0, 1]$

④ function spaces, e.g.,

all functions  $\mathbb{R} \rightarrow \mathbb{R}$

all functions  $[0, 1] \rightarrow \mathbb{R}$

addition  $(f+g)(x) = f(x) + g(x)$

multiplication  $(cf)(x) = c f(x)$

$$g = 5f \text{ means}$$

$$g(x) = 5f(x)$$

$$(f+g)(x) = f(x) + g(x)$$

LUM is NOT a vector space!

closed under scalar multiplication ✓

NOT closed under addition X

Subspaces!


⑤  $\{(x, 2x) : x \in \mathbb{R}\}$   
 includes  $(0,0)$ , closed under +,  $\times$  ✓

$\mathbb{R}^2$


in  $\mathbb{R}^3$ :

$\{\vec{0} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}\}, \mathbb{R}^3$   
 ↑ all lines thru  $\vec{0}$

- ②  $\{(x, 2x) : x \in \mathbb{R}\}$   
 includes  $(0,0)$ , closed under  $+$ ,  $\times$  ✓  
All subspaces of  $\mathbb{R}^2$ :  
 $\{\vec{o}\}$ , lines through  $\vec{o}$   
 $\{(x,y) : ax+by=0\}$ ,  $\mathbb{R}^2$  itself  
Not subspaces:  
 other lines: curves:



Important: Lines/planes/hyperplanes that don't go through the origin ( $\vec{o}$ ) are NOT subspaces!!



### ③ The SPAN of any (finite or infinite) set of points S

Span(S) is defined to be the set of all finite linear combinations of elements from S  
 i.e., all sums  $d_1 \vec{v}_1 + d_2 \vec{v}_2 + \dots + d_r \vec{v}_r$  for scalars  $d_i$  and  $\vec{v}_i \in S$

By definition, this is closed under  $+$ ,  $\times$ , and hence is a vector space.

$$\subseteq \left( \sum_{i=1}^r d_i \vec{v}_i \right) = \sum_{j=1}^r (c_j d_j) \vec{v}_j$$

Examples: What are:

- $\text{Span} \underbrace{\{(1, 2)\}}_{S} \subseteq \mathbb{R}^2 = \{c(1, 2) | c \in \mathbb{R}\}$

$\text{Span}(\{(1, 2)\}) = \{\vec{o}\}$

- $\text{Span} \{(1, 2), (-1, -2)\} = \text{Span}(\{(1, 2)\}) = \text{Span}(\{(-1, -2)\}) = \text{Span}(\{(1, 2), (2, 4)\}) = L$

$$\begin{aligned} \mathbb{R}^2 &= \text{Span}(\{(0, 1), (1, 0)\}) \\ &= \{c(0, 1) + d(1, 0) | c, d \in \mathbb{R}\} \\ &= \{(c, d) | c, d \in \mathbb{R}\} \end{aligned}$$

$\text{Span} \left( \begin{array}{l} \text{any two non zero} \\ \text{vectors in } \mathbb{R}^2 \text{ that} \\ \text{are not multiples of each other} \end{array} \right) = \mathbb{R}^2$

- $\text{Span} \{1, x, x^2, x^3, \dots\} = \left\{ \sum_{j=0}^K c_j x^j | c_j \in \mathbb{R}, K < \infty \right\}$

- $\text{Span} \left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 1 \end{pmatrix} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\} = \left\{ \begin{pmatrix} x \\ z \end{pmatrix} | x, z \in \mathbb{R} \right\} = xz\text{-plane} \subseteq \mathbb{R}^3$

$$\begin{array}{c} \text{C } \{ z \mid \\ = xz - \text{plane} \subseteq \mathbb{R}^3 \end{array}$$

- Span  $\{(1, 2, 1, 1, 5) = \vec{v}, (-2, -4, 0, 4, -2) = \vec{w}, (1, 2, 2, 4, 9) = \vec{x}\}$

-this is a subspace of  $\mathbb{R}^5$  — it must be either a line, a plane, or a 3D hyperplane

Two approaches to find out:

① Add vectors one at a time

$$\vec{v} = \begin{pmatrix} 1 \\ 2 \\ 1 \\ 1 \\ 5 \end{pmatrix} : \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 1 \\ 1 \\ 5 \end{pmatrix} \right\} = \text{a line} \dots$$



$$\vec{w} : \text{Span} \left\{ \vec{v}, \vec{w} \right\} = \text{a plane} \dots = \left\{ c\vec{v} + d\vec{w} \mid c, d \in \mathbb{R} \right\}$$

$\vec{w} \neq c\vec{v}$  for any  $c \in \mathbb{R}$

$$\vec{x} : \text{Span} \left\{ \vec{v}, \vec{w}, \vec{x} \right\}$$

? Is  $\vec{x} \in \text{Span} \left\{ \vec{v}, \vec{w} \right\}$ ?

do there exist  $c, d \in \mathbb{R}$  s.t.  $\vec{x} = c\vec{v} + d\vec{w}$ ?

is there any solution to

$$\begin{pmatrix} 1 & -2 \\ 2 & -4 \\ 1 & 0 \\ 1 & 4 \\ 5 & -2 \end{pmatrix} \begin{pmatrix} c \\ d \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 4 \\ 9 \end{pmatrix} ?$$

$$\begin{pmatrix} c=2 \\ d=\frac{1}{2} \end{pmatrix}$$

$$\vec{x} = 2\vec{v} + \frac{1}{2}\vec{w} \in \text{Span} \left\{ \vec{v}, \vec{w} \right\} = \text{Span} \left\{ \vec{v}, \vec{w}, \vec{x} \right\}$$

$$\begin{aligned} \text{Span} \left\{ \vec{v}_1, \dots, \vec{v}_k \right\} \\ = \left\{ \vec{x} \mid \begin{array}{l} \text{there is a solution to} \\ \left( \begin{array}{cccc|c} 1 & 1 & \dots & 1 & \\ \vec{v}_1 & \vec{v}_2 & \dots & \vec{v}_k & \\ \hline 1 & 1 & \dots & 1 & \end{array} \right) \vec{y} = \vec{x} \end{array} \right\} \end{aligned}$$

② Start with all the vectors, and try to simplify them  
This is what Gaussian elimination does:

$$\text{Let } M = \begin{pmatrix} 1 & 2 & 1 & 1 & 5 \\ -2 & -4 & 0 & 4 & -2 \\ 1 & 2 & 2 & 4 & 9 \end{pmatrix} \xrightarrow{-1}$$

$$M \xrightarrow{\text{GE}} \begin{pmatrix} 1 & 2 & 1 & 1 & 5 \\ 0 & 0 & 2 & 6 & 8 \\ 0 & 0 & 1 & 3 & 4 \end{pmatrix} \xrightarrow{\text{GE}} \begin{pmatrix} 1 & 2 & 1 & 1 & 5 \\ 0 & 0 & 1 & 3 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \xrightarrow{-1/2}$$

$$\xrightarrow{\text{GE}} \begin{pmatrix} 1 & 2 & 0 & -2 & 1 \\ 0 & 0 & 1 & 3 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow \text{Span} \left\{ \vec{v}, \vec{w}, \vec{x} \right\} = \text{Span} \left\{ \begin{pmatrix} 1 \\ 2 \\ 0 \\ -2 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right\}$$

$$\begin{pmatrix} 1 \\ 2 \\ 0 \\ -2 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\left( \begin{array}{ccc} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \\ \vdots & \vdots & \vdots \\ \vec{v}_n & \vec{v}_n & \vec{v}_n \end{array} \right) \xrightarrow{\text{GE}} \left( \begin{array}{ccc} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 \\ \vdots & \vdots & \vdots \\ \vec{v}_n & \vec{v}_n & \vec{v}_n \end{array} \right)$$

$$\text{Span} \left\{ \vec{v}_j \right\} = \text{Span} \left\{ \vec{v}_j \right\}$$

GE does not change the span of the rows

GE does not change the span of the rows

Why?

$$\text{Span} \left\{ \vec{v}_1, \vec{v}_2, \dots, \vec{v}_k \right\} = \text{Span} \left\{ \underbrace{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_k}_{\sum_{j=1}^k a_j \vec{v}_j} \right\}$$

$$\underbrace{a_1(c\vec{v}_1) + a_2\vec{v}_2 + \dots + a_k\vec{v}_k}$$

$$\text{Span} \left\{ \vec{v}_1, \vec{v}_2, \dots, \vec{v}_k \right\} = \text{Span} \left\{ \vec{v}_1 + \vec{v}_2, \vec{v}_2, \vec{v}_3, \dots, \vec{v}_k \right\}$$

$$\underbrace{a_1\vec{v}_1 + a_2\vec{v}_2 + \dots + a_k\vec{v}_k}$$

$$\underbrace{a_1(\vec{v}_1 + \vec{v}_2) + (a_2 - a_1)\vec{v}_2 + a_3\vec{v}_3 + \dots + a_k\vec{v}_k}$$

Note: Adding a multiple of one row to another does not change the span of the rows.

$$\text{Span} \left\{ \vec{v}_1, \vec{v}_2, \vec{v}_3, \dots \right\}$$

$$= \text{Span} \left\{ \vec{v}_1, \vec{v}_2 + \beta \vec{v}_1, \vec{v}_3, \dots \right\}$$

since anything you can reach with a linear combination

$$d_1\vec{v}_1 + d_2\vec{v}_2 + d_3\vec{v}_3 + \dots$$

can also be reached with a linear combination of the new vectors

$$(d_1 - \beta d_2)\vec{v}_1 + d_2(\vec{v}_2 + \beta \vec{v}_1) + d_3\vec{v}_3 + \dots,$$

and vice versa.

⇒ The nonzero rows left over after Gaussian elimination (are a minimal set of vectors) that span the same set as the original rows.

Claim:  $\text{Span}(S)$  is the **smallest** vector space that contains all the points in  $S$ .

Proof:

- Let  $T$  be a vector space containing all of  $S$ .

Goal: Show  $\text{Span}(S) \subseteq T$ .

- Let  $\vec{v} \in \text{Span}(S)$ . Goal: Show  $\vec{v} \in T$ .

$$\Rightarrow \vec{v} = c_1\vec{v}_1 + c_2\vec{v}_2 + \dots + c_r\vec{v}_r, \text{ with all } \vec{v}_j \in S, c_j \in \mathbb{R}$$

$\Rightarrow$  all  $\vec{v}_j \in T \Rightarrow c_j\vec{v}_j \in T$  by closure under multiplication

$\Rightarrow c_1\vec{v}_1 + \dots + c_r\vec{v}_r \in T$  by closure under addition

$\Rightarrow \vec{v} \in T$   $\square$

$$\text{④ Polynomials} = \text{Span} \left\{ 1, x, x^2, x^3, \dots \right\}$$

Continuous functions ✓

Differentiable functions ✓

Functions  $f$  with  $f(1) = 0$  ✓

...

Definition: **Affine subspace** = translated subspace

i.e., a set  $\vec{u} + V$  for a vector  $\vec{u} \neq \vec{0}$   
and subspace  $V$ .


⑧ The **Sum** of two subspaces is a subspace.

Definition: For subsets  $X$  and  $Y$  of a vector space,


$$X + Y = \{x + y \mid x \in X, y \in Y\}.$$

(In English: all sums of a vector in  $X$  and a vector in  $Y$ .)

Example:



Example:



Claim 1: If  $X$  and  $Y$  are subspaces,  
then  $X+Y$  is a subspace.

Proof: The key properties to check are closure under  $+$  and  $\cdot$ .

Closure under addition:

Want to show (WTS) if  $a, b \in X+Y$ , then  $a+b \in X+Y$ .

$$\begin{aligned} a \in X+Y &\Rightarrow a = x+y \text{ for some } x \in X, y \in Y \\ b \in X+Y &\Rightarrow b = x'+y' \quad " \quad x' \quad " \quad y' \\ a+b &= (x+y) + (x'+y') \\ &= (x+x') + (y+y') \in X+Y \quad \checkmark \end{aligned}$$

Closure under multiplication:

WTS: If  $a \in X+Y$ , then for all scalars  $\alpha$ ,  $\alpha a \in X+Y$ :

$$\begin{aligned} a &= x+y \\ \alpha a &= \alpha(x+y) = (\alpha x) + (\alpha y) \quad \checkmark \end{aligned}$$

□

Claim 2: For subsets  $S$  and  $T$  of a vector space  $V$ ,

$$\text{Span}(S) + \text{Span}(T) = \text{Span}(S \cup T)$$

Proof:  $\text{Span}(S) = \{\text{finite linear combinations of elts of } S\}$


$\text{Span}(T) = \{\text{finite linear combinations of elts of } T\}$

$\therefore x \in \text{Span}(S) + \text{Span}(T)$

$$\Leftrightarrow x = \sum_{j=1}^k \alpha_j s_j + \sum_{j=1}^l \beta_j t_j$$

$\Leftrightarrow x$  is a finite linear combination of elements of  $S \cup T$ .  $\checkmark$  □

Example: From before



More examples of vector spaces

Space	Closed under addition?	Closed under multiplication?	Vector space?
$\mathbb{C}^{n \times n}$ in $\mathbb{M}^n$	✓	✓	✓

Space      Closed under addition?      multiplication?      Vector space?


$$V_1 = \left\{ (b_1, b_2, b_3) \in \mathbb{R}^3 \mid \begin{array}{l} \text{st. } \\ b_1 - 2b_2 + 3b_3 = 0 \end{array} \right\}$$

Yes      Yes      ✓

$$V_2 = \left\{ (b_1, b_2, b_3) \in \mathbb{R}^3 \mid \begin{array}{l} \text{st. } \\ b_2 b_3 = 0 \end{array} \right\}$$


No      Yes      No

$$\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \in V_2$$



$$V_3 = \left\{ \vec{b} \in \mathbb{R}^2 \mid b_1 + b_2 = 1 \right\}$$

No      No      No



$$V_h = \left\{ \vec{b} \in \mathbb{R}^n \mid A\vec{b} = \vec{0} \right\}$$

solutions to these  
homogeneous equations

Yes      Yes

closure under addition:  
 $\vec{b}, \vec{c} \in V_h \Rightarrow \vec{b} + \vec{c} \in V_h$  ✓

$$A\vec{b} = \vec{0} = A\vec{c}$$

$$A(\vec{b} + \vec{c}) = \vec{0}$$

closure under multiplication  
 $c \in \mathbb{R}, \vec{b} \in V_h \Rightarrow c\vec{b} \in V_h$  ✓

$$A\vec{b} = \vec{0} \Rightarrow A(c\vec{b}) = cA\vec{b} = c\cdot\vec{0} = \vec{0}$$

No

$$V = \left\{ \vec{b} \in \mathbb{R}^n \mid A\vec{b} = \vec{c} \right\}$$

where  $A \in \mathbb{R}^{m \times n}$  are given  
and  $\vec{c} \neq \vec{0}$

addition      multiplication

$\vec{0} \notin V$   
because  
vector space?

$$A\vec{0} = \vec{0} \neq \vec{c}$$

$$V_4 = \text{Span}(\{(1,1,0), (2,0,1)\})$$

✓

✓

✓

$$V_5 = \left\{ \text{upper-triangular } m \times m \text{ matrices} \right\}$$

✓

✓

✓

$$V_6 = \left\{ \text{diagonal } n \times n \text{ matrices} \right\}$$

✓

✓

✓

$$V_7 = \left\{ 10 \times 10 \text{ matrices } A \mid \text{rank}(A) \geq 10 \right\}$$

✓

✓

✓

$V_6 = \{ \text{all } 10 \times 10 \text{ matrices } A \}$

$$V_7 = \left\{ \begin{array}{l} 10 \times 10 \text{ matrices } A \\ \text{with } \text{Trace}(A) = \sum_{i=1}^{10} a_{ii} = 0 \end{array} \right\}$$

$$V_8 = \left\{ \begin{array}{l} 10 \times 10 \text{ matrices } A \\ \text{with } \text{Tr}(A) = 1 \end{array} \right\}$$

$$V_9 = \left\{ \begin{array}{l} 3 \times 3 \text{ matrices } A \\ \text{with } A \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = 0 \end{array} \right\}$$

$$V_{10} = \left\{ \begin{array}{l} \text{symmetric } n \times n \text{ matrices} \\ (\text{i.e. } A = A^T) \end{array} \right\}$$

✓ ✓ ✓

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \notin V_8$$

✗

$$a_{11} + a_{12} + a_{13} = 0$$

$$a_{21} + a_{22} + a_{23} = 0$$

$$a_{31} + a_{32} + a_{33} = 0$$

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix} \quad \begin{pmatrix} a & b & c \\ b & d & e \\ c & e & f \end{pmatrix}$$

$A_{ij} \neq 0 \quad a_{ij} - a_{ji} = 0$

✓

$$\left\{ \begin{array}{l} \text{arithmetic progressions,} \\ \text{i.e., sequences } (x_1, x_2, x_3, \dots) \end{array} \right\}$$

$$\left\{ \begin{array}{l} \text{with } x_j - x_{j-1} \text{ constant} \end{array} \right\}$$

$$\text{(e.g., } (0, 2, 4, 6, 8, 10, \dots)$$

$$\left\{ \begin{array}{l} \text{differentiable functions} \\ f: \mathbb{R} \rightarrow \mathbb{R} \text{ with } f'(2) = 3 \end{array} \right\}$$

$$+ (1, 2, 3, 4, 5, 6, \dots)$$

$$= (1, 4, 7, 10, 13, 16, \dots)$$

✓

$$\left\{ (0, 0), (2, 1), (4, 2), (6, 3), \dots, (94, 47), (96, 48) \right\}$$

$$\subset (\mathbb{Z}/97\mathbb{Z}) \times (\mathbb{Z}/97\mathbb{Z})$$

(97 is prime)

$$c(0, 2, 4, 6, 8, \dots)$$

$$= (c, 2c, 4c, 6c, \dots)$$

✓

$$\left\{ \begin{array}{l} \text{all matrices that commute} \\ \text{with a given matrix } A \end{array} \right\}$$

✓

*Proof.* To prove (4.1.1), demonstrate that the two closure properties **(A1)** and **(M1)** hold for  $\mathcal{S} = \mathcal{X} + \mathcal{Y}$ . To show **(A1)** is valid, observe that if  $\mathbf{u}, \mathbf{v} \in \mathcal{S}$ , then  $\mathbf{u} = \mathbf{x}_1 + \mathbf{y}_1$  and  $\mathbf{v} = \mathbf{x}_2 + \mathbf{y}_2$ , where  $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}$  and  $\mathbf{y}_1, \mathbf{y}_2 \in \mathcal{Y}$ . Because  $\mathcal{X}$  and  $\mathcal{Y}$  are closed with respect to addition, it follows that  $\mathbf{x}_1 + \mathbf{x}_2 \in \mathcal{X}$  and  $\mathbf{y}_1 + \mathbf{y}_2 \in \mathcal{Y}$ , and therefore  $\mathbf{u} + \mathbf{v} = (\mathbf{x}_1 + \mathbf{x}_2) + (\mathbf{y}_1 + \mathbf{y}_2) \in \mathcal{S}$ . To verify **(M1)**, observe that  $\mathcal{X}$  and  $\mathcal{Y}$  are both closed with respect to scalar multiplication so that  $\alpha\mathbf{x}_1 \in \mathcal{X}$  and  $\alpha\mathbf{y}_1 \in \mathcal{Y}$  for all  $\alpha$ , and consequently  $\alpha\mathbf{u} = \alpha\mathbf{x}_1 + \alpha\mathbf{y}_1 \in \mathcal{S}$  for all  $\alpha$ . To prove (4.1.2), suppose  $\mathcal{S}_{\mathcal{X}} = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_r\}$  and  $\mathcal{S}_{\mathcal{Y}} = \{\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_t\}$ , and write

$$\begin{aligned} \mathbf{z} \in \text{span}(\mathcal{S}_{\mathcal{X}} \cup \mathcal{S}_{\mathcal{Y}}) &\iff \mathbf{z} = \sum_{i=1}^r \alpha_i \mathbf{x}_i + \sum_{i=1}^t \beta_i \mathbf{y}_i = \mathbf{x} + \mathbf{y} \text{ with } \mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{Y} \\ &\iff \mathbf{z} \in \mathcal{X} + \mathcal{Y}. \blacksquare \end{aligned}$$

### Example 4.1.8

If  $\mathcal{X} \subseteq \mathbb{R}^2$  and  $\mathcal{Y} \subseteq \mathbb{R}^2$  are subspaces defined by two different lines through the origin, then  $\mathcal{X} + \mathcal{Y} = \mathbb{R}^2$ . This follows from the parallelogram law—sketch a picture for yourself.

### Exercises for section 4.1

4.1.1. Determine which of the following subsets of  $\mathbb{R}^n$  are in fact subspaces of  $\mathbb{R}^n$  ( $n > 2$ ).

- $\{\mathbf{x} \mid x_1 \geq 0\}$ ,  $\{\mathbf{x} \mid x_1 = 0\}$ ,  $\{\mathbf{x} \mid x_1 x_2 = 0\}$ ,
- $\left\{ \mathbf{x} \mid \sum_{j=1}^n x_j = 0 \right\}$ ,  $\left\{ \mathbf{x} \mid \sum_{j=1}^n x_j = 1 \right\}$ ,
- $\{\mathbf{x} \mid \mathbf{A}\mathbf{x} = \mathbf{b}, \text{ where } \mathbf{A}_{m \times n} \neq \mathbf{0} \text{ and } \mathbf{b}_{m \times 1} \neq \mathbf{0}\}$ .

4.1.2. Determine which of the following subsets of  $\mathbb{R}^{n \times n}$  are in fact subspaces of  $\mathbb{R}^{n \times n}$ .

- (a) The symmetric matrices. (b) The diagonal matrices.  
 (c) The nonsingular matrices. (d) The singular matrices.  
 (e) The triangular matrices. (f) The upper-triangular matrices.  
 (g) All matrices that commute with a given matrix  $\mathbf{A}$ .  
 (h) All matrices such that  $\mathbf{A}^2 = \mathbf{A}$ .  
 (i) All matrices such that  $\text{trace}(\mathbf{A}) = 0$ .

4.1.3. If  $\mathcal{X}$  is a plane passing through the origin in  $\mathbb{R}^3$  and  $\mathcal{Y}$  is the line through the origin that is perpendicular to  $\mathcal{X}$ , what is  $\mathcal{X} + \mathcal{Y}$ ?

- 4.1.4. Why must a real or complex nonzero vector space contain an infinite number of vectors?

- 4.1.5. Sketch a picture in  $\mathbb{R}^3$  of the subspace spanned by each of the following.
- (a)  $\left\{ \begin{pmatrix} 1 \\ 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 6 \\ 4 \end{pmatrix}, \begin{pmatrix} -3 \\ -9 \\ -6 \end{pmatrix} \right\}$ , (b)  $\left\{ \begin{pmatrix} -4 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 5 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \right\}$ , *try plane*  
 (c)  $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$ , *line*

- 4.1.6. Which of the following are spanning sets for  $\mathbb{R}^3$ ?
- (a)  $\left\{ \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \right\}$ , (b)  $\left\{ \begin{pmatrix} 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \end{pmatrix} \right\}$ ,  
 (c)  $\left\{ \begin{pmatrix} 1 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \right\}$ ,  
 (d)  $\left\{ \begin{pmatrix} 1 & 2 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 1 \end{pmatrix} \right\}$ , *first+second*  
 (e)  $\left\{ \begin{pmatrix} 1 & 2 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 & -1 \end{pmatrix}, \begin{pmatrix} 4 & 4 & 0 \end{pmatrix} \right\}$ .

- 4.1.7. For a vector space  $\mathcal{V}$ , and for  $\mathcal{M}, \mathcal{N} \subseteq \mathcal{V}$ , explain why  
*skip*  $\text{span}(\mathcal{M} \cup \mathcal{N}) = \text{span}(\mathcal{M}) + \text{span}(\mathcal{N})$ .

- 4.1.8. Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two subspaces of a vector space  $\mathcal{V}$ .
- (a) Prove that the intersection  $\mathcal{X} \cap \mathcal{Y}$  is also a subspace of  $\mathcal{V}$ . ✓  
 (b) Show that the union  $\mathcal{X} \cup \mathcal{Y}$  need not be a subspace of  $\mathcal{V}$ .

- 4.1.9. For  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\mathcal{S} \subseteq \mathbb{R}^{n \times 1}$ , the set  $\mathbf{A}(\mathcal{S}) = \{\mathbf{Ax} \mid \mathbf{x} \in \mathcal{S}\}$  contains all possible products of  $\mathbf{A}$  with vectors from  $\mathcal{S}$ . We refer to  $\mathbf{A}(\mathcal{S})$  as the set of *images* of  $\mathcal{S}$  under  $\mathbf{A}$ .
- (a) If  $\mathcal{S}$  is a subspace of  $\mathbb{R}^n$ , prove  $\mathbf{A}(\mathcal{S})$  is a subspace of  $\mathbb{R}^m$ . ✓  
 (b) If  $\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_k$  spans  $\mathcal{S}$ , show  $\mathbf{As}_1, \mathbf{As}_2, \dots, \mathbf{As}_k$  spans  $\mathbf{A}(\mathcal{S})$ . ✓

- 4.1.10. With the usual addition and multiplication, determine whether or not the following sets are vector spaces over the real numbers.
- (a)  $\mathbb{R}$ , ✓ (b)  $\mathbb{C}$ , ✓ (c) The rational numbers, *X*

- 4.1.11. Let  $\mathcal{M} = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_r\}$  and  $\mathcal{N} = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_r, \mathbf{v}\}$  be two sets of vectors from the same vector space. Prove that  $\text{span}(\mathcal{M}) = \text{span}(\mathcal{N})$  if and only if  $\mathbf{v} \in \text{span}(\mathcal{M})$ .

- 4.1.12. For a set of vectors  $\mathcal{S} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ , prove that  $\text{span}(\mathcal{S})$  is the intersection of all subspaces that contain  $\mathcal{S}$ . Hint: For  $\mathcal{M} = \bigcap_{\mathcal{S} \subseteq \mathcal{V}} \mathcal{S}$ , prove that  $\text{span}(\mathcal{S}) \subseteq \mathcal{M}$  and  $\mathcal{M} \subseteq \text{span}(\mathcal{S})$ .

## ⑥ Examples over other fields

$\mathbb{F}_p$  = field of numbers mod p (for a prime p)

$\mathbb{F}_2 = \{0, 1\}$

Bit strings of length n form a vector space:

n=3:  $(0,0,0), (0,0,1), (0,1,0), (0,1,1)$   
 $(1,0,0), (1,0,1), (1,1,0), (1,1,1)$

addition is coordinate-wise, mod 2

$$(0,0,1) + (0,1,1) = (0,1,0)$$

subspace, e.g.,

$$\text{Span}(\{(0,0,1), (1,0,1)\}) \\ = \{(0,0,0), (0,0,1), (1,0,1), (1,0,0)\}$$

Problems:

1. How many subspaces are there of  $\mathbb{R}^2$ ?

Answer: Infinitely many!  
 (Lines through the origin)

2. How many subspaces are there of  $\mathbb{R}$ ?

Answer: Two!  $\{0\}$  and  $\mathbb{R}$  itself.

3. How many subspaces are there of  $\mathbb{Z}_2^2$ ?

$\{(0,0)\}$ , everything  $\mathbb{Z}_2^2$

$\{(0,0), (0,1)\}, \{(0,0), (1,0)\}$

$\{(0,0), (1,0)\}, \{(0,0), (1,1)\}$

and that's it!

$$\{(0,0), (0,1)\}, \{(0,0), (1,0)\}$$

$$\{(0,0), (1,0)\}, \{(0,0), (1,1)\}$$

and that's it!

if a subspace contains two of the nonzero points, then it also includes their sum, which is the last nonzero point  $\therefore (1,0) + (0,1) + (1,1) = (0,0)$   
means that any two sum to the third

$$1 + 1 + 1 = 6$$

4. How many subspaces are there of  $\mathbb{R}^3$ ?  
We'll answer this later! Definitely  $\infty$  though

vector  $(1,1,-1)$  and automatically contains any multiple  $(c,c,-c)$ :

$$\text{Nullspace is a line } \begin{bmatrix} 1 & 0 & 1 \\ 5 & 4 & 9 \\ 2 & 4 & 0 \end{bmatrix} \begin{bmatrix} c & c & -c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

The nullspace of  $B$  is the line of all points  $x = c, y = c, z = -c$ . (The line goes through the origin, as any subspace must.) We want to be able, for any system  $Ax = b$ , to find  $C(A)$  and  $N(A)$ : all attainable right-hand sides  $b$  and all solutions to  $Ax = 0$ .

The vectors  $b$  are in the column space and the vectors  $x$  are in the nullspace. We shall compute the dimensions of those subspaces and a convenient set of vectors to generate them. We hope to end up by understanding all four of the subspaces that are intimately related to each other and to  $A$ —the column space of  $A$ , the nullspace of  $A$ , and their two perpendicular spaces.

### Problem Set 2.1

- Construct a subset of the  $x$ - $y$  plane  $\mathbb{R}^2$  that is
  - closed under vector addition and subtraction, but not scalar multiplication.
  - closed under scalar multiplication but not under vector addition.

*Hint:* Starting with  $u$  and  $v$ , add and subtract for (a). Try  $cu$  and  $cv$  for (b).
- Which of the following subsets of  $\mathbb{R}^3$  are actually subspaces?
  - The plane of vectors  $(b_1, b_2, b_3)$  with first component  $b_1 = 0$ .
  - The plane of vectors  $b$  with  $b_1 = 1$ .
  - The vectors  $b$  with  $b_2 b_3 = 0$  (this is the union of two subspaces, the plane  $b_2 = 0$  and the plane  $b_3 = 0$ ).
  - All combinations of two given vectors  $(1, 1, 0)$  and  $(2, 0, 1)$ .
  - The plane of vectors  $(b_1, b_2, b_3)$  that satisfy  $b_3 - b_2 + 3b_1 = 0$ .
- Describe the column space and the nullspace of the matrices
 
$$A = \begin{bmatrix} 1 & -1 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 0 & 3 \\ 1 & 2 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$
- What is the smallest subspace of 3 by 3 matrices that contains all symmetric matrices and all lower triangular matrices? What is the largest subspace that is contained in both of those subspaces?
- Addition and scalar multiplication are required to satisfy these eight rules:

- $x + y = y + x$ .
- $x + (y + z) = (x + y) + z$ .
- There is a unique “zero vector” such that  $x + 0 = x$  for all  $x$ .
- For each  $x$  there is a unique vector  $-x$  such that  $x + (-x) = 0$ .
- $1x = x$ .
- $(c_1 c_2)x = c_1(c_2x)$ .
- $c(x + y) = cx + cy$ .
- $(c_1 + c_2)x = c_1x + c_2x$ .
  - Suppose addition in  $\mathbb{R}^2$  adds an extra 1 to each component, so that  $(3, 1) + (5, 0)$  equals  $(9, 2)$  instead of  $(8, 1)$ . With scalar multiplication unchanged, which rules are broken?
  - Show that the set of all positive real numbers, with  $x + y$  and  $cx$  redefined to equal the usual  $xy$  and  $x'$ , is a vector space. What is the “zero vector”?
  - Suppose  $(x_1, x_2) + (y_1, y_2)$  is defined to be  $(x_1 + y_2, x_2 + y_1)$ . With the usual  $cx = (cx_1, cx_2)$ , which of the eight conditions are not satisfied?

1.  $x+y=y+x$ .  
 2.  $x+(y+z)=(x+y)+z$ .  
 3. There is a unique “zero vector” such that  $x+0=x$  for all  $x$ .  
 4. For each  $x$  there is a unique vector  $-x$  such that  $x+(-x)=0$ .  
 5.  $1x=x$ .  
 6.  $(c_1c_2)x=c_1(c_2x)$ .  
 7.  $c(x+y)=cx+cy$ .  
 8.  $(c_1+c_2)x=c_1x+c_2x$ .
- (a) Suppose addition in  $\mathbf{R}^2$  adds an extra 1 to each component, so that  $(3, 1) + (5, 0)$  equals  $(9, 2)$  instead of  $(8, 1)$ . With scalar multiplication unchanged, which rules are broken?  
 (b) Show that the set of all positive real numbers, with  $x+y$  and  $cx$  redefined to equal the usual  $xy$  and  $x'$ , is a vector space. What is the “zero vector”?  
 (c) Suppose  $(x_1, x_2) + (y_1, y_2)$  is defined to be  $(x_1 + y_2, x_2 + y_1)$ . With the usual  $cx = (cx_1, cx_2)$ , which of the eight conditions are not satisfied?
6. Let  $\mathbf{P}$  be the plane in 3-space with equation  $x+2y+z=6$ . What is the equation of the plane  $\mathbf{P}_0$  through the origin parallel to  $\mathbf{P}$ ? Are  $\mathbf{P}$  and  $\mathbf{P}_0$  subspaces of  $\mathbf{R}^3$ ?  
 7. Which of the following are subspaces of  $\mathbf{R}^\infty$ ?  
 (a) All sequences like  $(1, 0, 1, 0, \dots)$  that include infinitely many zeros.  
 (b) All sequences  $(x_1, x_2, \dots)$  with  $x_j=0$  from some point onward.  
 (c) All decreasing sequences:  $x_{j+1} \leq x_j$  for each  $j$ .  
 (d) All convergent sequences: the  $x_j$  have a limit as  $j \rightarrow \infty$ .  
 (e) All arithmetic progressions:  $x_{j+1} - x_j$  is the same for all  $j$ .  
 (f) All geometric progressions  $(x_1, kx_1, k^2x_1, \dots)$  allowing all  $k$  and  $x_1$ .
8. Which of the following descriptions are correct? The solutions  $x$  of

$$Ax = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

form

- (a) a plane.  
 (b) a line.  
 (c) a point.  
 (d) a subspace.

- (e) the nullspace of  $A$ .  
 (f) the column space of  $A$ .  
 9. Show that the set of nonsingular 2 by 2 matrices is not a vector space. Show also that the set of *singular* 2 by 2 matrices is not a vector space.  
 10. The matrix  $A = \begin{bmatrix} 2 & -2 \\ 2 & -2 \end{bmatrix}$  is a “vector” in the space  $\mathbf{M}$  of all 2 by 2 matrices. Write the zero vector in this space, the vector  $\frac{1}{2}A$ , and the vector  $-A$ . What matrices are in the smallest subspace containing  $A$ ?  
 11. (a) Describe a subspace of  $\mathbf{M}$  that contains  $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  but not  $B = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix}$ .  
 (b) If a subspace of  $\mathbf{M}$  contains  $A$  and  $B$ , must it contain  $I$ ?  
 (c) Describe a subspace of  $\mathbf{M}$  that contains no nonzero diagonal matrices.  
 12. The functions  $f(x) = x^2$  and  $g(x) = 5x$  are “vectors” in the vector space  $\mathbf{F}$  of all real functions. The combination  $3f(x) - 4g(x)$  is the function  $h(x) = \underline{\hspace{2cm}}$ . Which rule is broken if multiplying  $f(x)$  by  $c$  gives the function  $f(cx)$ ?  
 13. If the sum of the “vectors”  $f(x)$  and  $g(x)$  in  $\mathbf{F}$  is defined to be  $f(g(x))$ , then the “zero vector” is  $g(x) = x$ . Keep the usual scalar multiplication  $cf(x)$ , and find two rules that are broken.  
 14. Describe the smallest subspace of the 2 by 2 matrix space  $\mathbf{M}$  that contains  
 (a)  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ .      (b)  $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ .  
 (c)  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ .      (d)  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$ ,  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$ .  
 15. Let  $\mathbf{P}$  be the plane in  $\mathbf{R}^3$  with equation  $x+y-2z=4$ . The origin  $(0, 0, 0)$  is not in  $\mathbf{P}$ ! Find two vectors in  $\mathbf{P}$  and check that their sum is not in  $\mathbf{P}$ .  
 16.  $\mathbf{P}_0$  is the plane through  $(0, 0, 0)$  parallel to the plane  $\mathbf{P}$  in Problem 15. What is the equation for  $\mathbf{P}_0$ ? Find two vectors in  $\mathbf{P}_0$  and check that their sum is in  $\mathbf{P}_0$ .  
 17. The four types of subspaces of  $\mathbf{R}^3$  are planes, lines,  $\mathbf{R}^3$  itself, or  $\mathbf{Z}$  containing only  $(0, 0, 0)$ .  
 (a) Describe the three types of subspaces of  $\mathbf{R}^2$ .  
 (b) Describe the five types of subspaces of  $\mathbf{R}^4$ .  
 18. (a) The intersection of two planes through  $(0, 0, 0)$  is probably a       but it could be a      . It can’t be the zero vector  $\mathbf{Z}$ !  
 (b) The intersection of a plane through  $(0, 0, 0)$  with a line through  $(0, 0, 0)$  is probably a       but it could be a      .

18. (a) The intersection of two planes through  $(0,0,0)$  is probably a \_\_\_\_\_ but it could be a \_\_\_\_\_. It can't be the zero vector  $\mathbf{Z}$ !  
 (b) The intersection of a plane through  $(0,0,0)$  with a line through  $(0,0,0)$  is probably a \_\_\_\_\_ but it could be a \_\_\_\_\_.

- (c) If  $\mathbf{S}$  and  $\mathbf{T}$  are subspaces of  $\mathbf{R}^5$ , their intersection  $\mathbf{S} \cap \mathbf{T}$  (vectors in both subspaces) is a subspace of  $\mathbf{R}^5$ . Check the requirements on  $x+y$  and  $cx$ .
19. Suppose  $\mathbf{P}$  is a plane through  $(0,0,0)$  and  $\mathbf{L}$  is a line through  $(0,0,0)$ . The smallest vector space containing both  $\mathbf{P}$  and  $\mathbf{L}$  is either \_\_\_\_\_ or \_\_\_\_\_.
20. True or false for  $\mathbf{M} = \text{all } 3 \times 3 \text{ matrices}$  (check addition using an example)?  
 (a) The skew-symmetric matrices in  $\mathbf{M}$  (with  $A^T = -A$ ) form a subspace.  
 (b) The unsymmetric matrices in  $\mathbf{M}$  (with  $A^T \neq A$ ) form a subspace.  
 (c) The matrices that have  $(1,1,1)$  in their nullspace form a subspace.

**Problems 21–30 are about column spaces  $C(A)$  and the equation  $Ax = b$ .**

21. Describe the column spaces (lines or planes) of these particular matrices:

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 0 \\ 2 & 0 \\ 0 & 0 \end{bmatrix}.$$

22. For which right-hand sides (find a condition on  $b_1, b_2, b_3$ ) are these systems solvable?

$$(a) \begin{bmatrix} 1 & 4 & 2 \\ 2 & 8 & 4 \\ -1 & -4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}. \quad (b) \begin{bmatrix} 1 & 4 \\ 2 & 9 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

23. Adding row 1 of  $A$  to row 2 produces  $B$ . Adding column 1 to column 2 produces  $C$ . A combination of the columns of \_\_\_\_\_ is also a combination of the columns of  $A$ . Which two matrices have the same column \_\_\_\_\_?

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 \\ 3 & 6 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}.$$

24. For which vectors  $(b_1, b_2, b_3)$  do these systems have a solution?

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}.$$

25. (Recommended) If we add an extra column  $b$  to a matrix  $A$ , then the column space gets larger unless \_\_\_\_\_. Give an example in which the column space gets larger and an example in which it doesn't. Why is  $Ax = b$  solvable exactly when the column space doesn't get larger by including  $b$ ?

26. The columns of  $AB$  are combinations of the columns of  $A$ . This means: *The column space of  $AB$  is contained in (possibly equal to) the column space of  $A$ .* Give an example where the column spaces of  $A$  and  $AB$  are not equal.

27. If  $A$  is any  $8 \times 8$  invertible matrix, then its column space is \_\_\_\_\_. Why?  
 28. True or false (with a counterexample if false)?  
 (a) The vectors  $b$  that are not in the column space  $C(A)$  form a subspace.  
 (b) If  $C(A)$  contains only the zero vector, then  $A$  is the zero matrix.  
 (c) The column space of  $2A$  equals the column space of  $A$ .  
 (d) The column space of  $A - I$  equals the column space of  $A$ .  
 29. Construct a  $3 \times 3$  matrix whose column space contains  $(1,1,0)$  and  $(1,0,1)$  but not  $(1,1,1)$ . Construct a  $3 \times 3$  matrix whose column space is only a line.  
 30. If the  $9 \times 12$  system  $Ax = b$  is solvable for every  $b$ , then  $C(A) = _____$ .  
 31. Why isn't  $\mathbf{R}^2$  a subspace of  $\mathbf{R}^3$ ?

## 2.2 Solving $Ax = 0$ and $Ax = b$

Chapter 1 concentrated on square invertible matrices. There was one solution to  $Ax = b$  and it was  $x = -A^{-1}b$ . That solution was found by elimination (not by computing  $A^{-1}$ ). A rectangular matrix brings new possibilities— $U$  may not have a full set of pivots. This section goes onward from  $U$  to a reduced form  $R$ —the simplest matrix that elimination can give.  $R$  reveals all solutions immediately.

For an invertible matrix, the nullspace contains only  $x = 0$  (multiply  $Ax = 0$  by  $A^{-1}$ ). The column space is the whole space ( $Ax = b$  has a solution for every  $b$ ). The new questions appear when the nullspace contains more than the zero vector and/or the column space contains less than all vectors:

- Any vector  $x_n$  in the nullspace can be added to a particular solution  $x_p$ . The solutions to all linear equations have this form,  $x = x_p + x_n$ :

**Complete solution**  $Ax_p = b$  and  $Ax_n = 0$  produce  $A(x_p + x_n) = b$ .

space contains *less than all* vectors:

- Any vector  $x_n$  in the nullspace can be added to a particular solution  $x_p$ . The solutions to all linear equations have this form,  $x = x_p + x_n$ :

**Complete solution**  $Ax_p = b$  and  $Ax_n = 0$  produce  $A(x_p + x_n) = b$ .

- When the column space doesn't contain every  $b$  in  $\mathbb{R}^m$ , we need the conditions on  $b$  that make  $Ax = b$  solvable.

A 3 by 4 example will be a good size. We will write down all solutions to  $Ax = 0$ . We will find the conditions for  $b$  to lie in the column space (so that  $Ax = b$  is solvable). The 1 by 1 system  $0x = b$ , one equation and one unknown, shows two possibilities:

$0x = b$  has *no solution* unless  $b = 0$ . The column space of the 1 by 1 zero matrix contains only  $b = 0$ .

$0x = 0$  has *infinitely many solutions*. The nullspace contains *all*  $x$ . A particular solution is  $x_p = 0$ , and the complete solution is  $x = x_p + x_n = 0 + (\text{any } x)$ .

## More important examples: Subspaces of a Matrix

Let  $A \in \mathbb{R}^{m \times n}$  be an  $m \times n$  real-valued matrix.

- $\text{Range}(A) = \{\vec{A}\vec{x} \mid \vec{x} \in \mathbb{R}^n\}$   
= Span(columns of  $A$ )  
AKA "column space" of  $A$

Observe:  $\vec{b} \in \text{Range}(A) \Leftrightarrow A\vec{x} = \vec{b}$  has a solution

- $\text{Range}(A^T) = \text{Span}(\text{rows of } A)$   
"row space"
- $\text{Kernel}(A) = \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \vec{0}\}$  (solutions to the homogeneous equations)  
AKA "null space" of  $A$

Successive Fourier approximations to the step function

