

A128 · A novel RNA motif discovery pipeline to elucidate GLDR-2 target recognition

Presenter: Melissa Badendieck — Department of Biology & Biotechnology and Department of Chemistry & Biochemistry, Worcester Polytech

Authors: Melissa Badendieck, Shuling Lin, Lyah Esplana, Karl-Frédéric Vieux, Aya Narunsky

Session: Day 1 · Thu Oct 1, 2026 · 2:15–4:15 PM

RNA tailing is the non-templated, post-transcriptional addition of nucleotides to the 3'-ends of transcripts carried out by terminal nucleotidyl transferases (TNTs). GLDR-2 is a non-canonical TNT, lacking an RNA recognition domain, and is ubiquitously expressed in *C. elegans*. GLDR-2's closest homolog is GLD-2, a cytoplasmic poly(A) polymerase expressed in both *C. elegans* and mammals that plays a crucial role in gametogenesis and development. GLD-2-depleted *C. elegans* are completely sterile. Similarly, GLDR-2-depleted male *C. elegans* are sub-fertile. The exact mechanism underlying GLDR-2 targeting to specific RNAs remains unknown. We hypothesize that RNA-binding proteins (RBPs) mediate interactions between GLDR-2 and specific RNAs by binding conserved motifs shared among targets. We combine computational and experimental approaches to identify these motifs using a novel in silico motif-discovery pipeline. We developed a computational pipeline to identify candidate RNA motifs that mediate GLDR-2 association with its target RNAs, as identified by Vieux et al., 2021. We combine HMM-based sequence searches with structural modelling to identify recurring motifs among RNAs that interact with GLDR-2. Motifs are ranked based on their prevalence among targets and the conservation of their predicted secondary structures. The highest-ranking motifs will be experimentally validated using immunoprecipitation coupled with high-throughput sequencing to assess their association with GLDR-2. We will then disrupt candidate RBPs associated with these motifs using CRISPR/Cas9-based genome editing to evaluate their requirement for GLDR-2 targeting to specific RNAs. Ultimately, establishing the molecular mechanisms that determine GLDR-2 target recognition will advance our understanding of its roles in fertility and development.

A novel RNA motif discovery pipeline to elucidate GLDR-2 target recognition

Melissa Badendieck^{*,1,2}, Shuling Lin¹, Lyah Esplana², Karl-Frédéric Vieux¹, Aya Narunsky²

¹ Department of Biology & Biotechnology, Worcester Polytechnic Institute, Worcester, MA

² Department of Chemistry & Biochemistry, Worcester Polytechnic Institute, Worcester, MA

RNA tailing is the non-templated, post-transcriptional addition of nucleotides to the 3'-ends of transcripts carried out by terminal nucleotidyl transferases (TNTs). GLDR-2 is a non-canonical TNT, lacking an RNA recognition domain, and is ubiquitously expressed in *C. elegans*. GLDR-2's closest homolog is GLD-2, a cytoplasmic poly(A) polymerase expressed in both *C. elegans* and mammals that plays a crucial role in gametogenesis and development. GLD-2-depleted *C. elegans* are completely sterile. Similarly, GLDR-2-depleted male *C. elegans* are sub-fertile. The exact mechanism underlying GLDR-2 targeting to specific RNAs remains unknown. We hypothesize that RNA-binding proteins (RBPs) mediate interactions between GLDR-2 and specific RNAs by binding conserved motifs shared among targets. We combine computational and experimental approaches to identify these motifs using a novel *in silico* motif-discovery pipeline.

We developed a computational pipeline to identify candidate RNA motifs that mediate GLDR-2 association with its target RNAs, as identified by Vieux et al., 2021. We combine HMM-based sequence searches with structural modelling to identify recurring motifs among RNAs that interact with GLDR-2. Motifs are ranked based on their prevalence among targets and the conservation of their predicted secondary structures. The highest-ranking motifs will be experimentally validated using immunoprecipitation coupled with high-throughput sequencing to assess their association with GLDR-2. We will then disrupt candidate RBPs associated with these motifs using CRISPR/Cas9-based genome editing to evaluate their requirement for GLDR-2 targeting to specific RNAs. Ultimately, establishing the molecular mechanisms that determine GLDR-2 target recognition will advance our understanding of its roles in fertility and development.