

A155 · Multimodal data-driven approaches for discovery and validation of pneumonia sub-phenotypes

Presenter: Amulya Shastry — Boston University

Authors: Amulya Shastry

Session: Day 2 · Fri Oct 2, 2026 · 2:15–4:15 PM

Rationale: Elucidating pneumonia sub-phenotypes can improve host-based therapeutics. Pneumonia involves heterogeneous and dysregulated histopathology responses, but these histopathological changes within the lungs have not been explored for sub-phenotyping. **Objective:** To investigate whether pneumonic lungs can be clustered into sub-phenotypes based on histopathology and examine whether these sub-phenotypes re-emerge in a new cohort. **Methods:** H&E slides from subjects with pneumonia diagnosis (n=276) or with no known lung disease (n=16) at autopsy were scored across 18 histopathology features (Necrosis, Fibrosis, Edema etc.) by two pathologists (discovery cohort). Consensus clustering was used to group samples into clusters, and distance and variation-based clustering metrics were used to select the optimum number of clusters. 5 different leukocytes were quantified in a subset of patients (n=159) using multispectral immunofluorescence assay. A new cohort of samples were scored (n=104) using the same framework (validation cohort). Both cohorts were co-clustered to validate sub-phenotypes in the second cohort. **Results:** Our approach showed pneumonia samples cluster into seven different sub-phenotypes with distinct histopathology patterns including necrosuppurative, suppurative & chronic interstitial pneumonias. These sub-phenotypes also vary in their composition of immune cell patterns. Validation using a different cohort showed that when co-clustered with the original cohort, all seven sub-phenotypes re-emerge, with consistent histopathology signatures. **Conclusions:** Human pneumonias segregate into sub-phenotypes based on host responses and sub-phenotype markers could be further explored for therapeutic purposes. By illuminating a spectrum of histopathologies and discriminating discrete sub-phenotypes of pneumonia, a foundational framework emerges for developing and testing host-directed therapies for subsets of pneumonia patients.

Multimodal data-driven approaches for discovery and validation of pneumonia sub-phenotypes

Amulya Shastry^{1,2,3†*}, Bradley E. Hiller^{1,3†}, Nathan L. Sanders^{1,4}, Anna E. Tseng^{4,5}, Jeet Kothari⁵, Aoife K. O'Connell⁵, Hans P. Gertje⁵, Catherine T. Ha¹, Ekaterina Murzin⁹, Katrina E. Traber^{1,3}, James A. Lederer⁹, Matthew R. Jones^{1,3}, Florian Douam^{4,5}, Thomas G. Beach⁸, Daniel G. Remick⁷, Nicholas A. Crossland^{4,5,7}, Stefano Monti^{2,10}, Joshua D. Campbell², Joseph P. Mizgerd^{1,3,4,6*}

¹Pulmonary Center, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

²Section of Computational Biomedicine, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

³Department of Medicine, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

⁴Department of Virology, Immunology, and Microbiology, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

⁵National Emerging Infectious Diseases Laboratories (NEIDL), Boston University; Boston, MA, United States

⁶Department of Biochemistry and Cell Biology, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

⁷Department of Pathology & Laboratory Medicine, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

⁸Banner Sun Health Research Institute; Sun City, AZ, United States

⁹Department of Surgery, Brigham and Women's Hospital and Harvard Medical School; Boston, MA, United States

¹⁰Department of Biostatistics, School of Public Health, Boston University Chobanian & Avedisian School of Medicine; Boston, MA, United States

Rationale: Elucidating pneumonia sub-phenotypes can improve host-based therapeutics. Pneumonia involves heterogeneous and dysregulated histopathology responses, but these histopathological changes within the lungs have not been explored for sub-phenotyping.

Objective: To investigate whether pneumonic lungs can be clustered into sub-phenotypes based on histopathology and examine whether these sub-phenotypes re-emerge in a new cohort.

Methods: H&E slides from subjects with pneumonia diagnosis (n=276) or with no known lung disease (n=16) at autopsy were scored across 18 histopathology features (Necrosis, Fibrosis, Edema etc..) by two pathologists (discovery cohort). Consensus clustering was used to group samples into clusters, and distance and variation-based clustering metrics were used to select the optimum number of clusters. 5 different leukocytes were quantified in a subset of patients (n=159) using multispectral immunofluorescence assay. A new cohort of samples were scored (n=104) using the same framework (validation cohort). Both cohorts were co-clustered to validate sub-phenotypes in the second cohort.

Results: Our approach showed pneumonia samples cluster into seven different sub-phenotypes with distinct histopathology patterns including necrosuppurative, suppurative & chronic interstitial pneumonias. These sub-phenotypes also vary in their composition of immune cell patterns. Validation using a different cohort showed that when co-clustered with the original cohort, all seven sub-phenotypes re-emerge, with consistent histopathology signatures.

Conclusions: Human pneumonias segregate into sub-phenotypes based on host responses and sub-phenotype markers could be further explored for therapeutic purposes. By illuminating a spectrum of histopathologies and discriminating discrete sub-phenotypes of pneumonia, a foundational framework emerges for developing and testing host-directed therapies for subsets of pneumonia patients.