# **Seiad Valley Groundwater Basin**

Groundwater Basin Number: 1-16

County: Siskiyou

• Surface Area: 2,250 acres (4 square miles)

## **Basin Boundaries and Hydrology**

The Seiad Valley Groundwater Basin is an irregularly shaped basin that includes the valleys of Canyon and Grider Creeks and a short reach of the Klamath River. The basin is bounded to the south by granitic rocks of Slinkard Peak . The elongated shape of the Canyon Creek valley is controlled by a fault and is bounded to the south by Pre-Cretaceous metavolcanic rocks and to the north by Pre-Silurian metamorphic rocks (Strand 1962). The basin fill is comprised of Quaternary alluvium. Annual precipitation in the basin ranges from 29- to 41-inches, increasing from east to west, and increases to 53-inches to the south in the Grider Creek area.

### **Hydrogeologic Information**

Hydrogeologic information was not available for the following:

Water-Bearing Formations Groundwater Level Trends Groundwater Storage

## Groundwater Budget (Type B)

The estimate of groundwater extraction for the Seiad Valley basin is based on a survey conducted during 1991 by the California Department of Water Resources. The survey included landuse and sources of water. Groundwater extraction for municipal and industrial uses is estimated to be 29 acre-feet. Deep percolation of applied water is estimated to be 29 acre-feet.

# **Groundwater Quality**

# Water Quality in Public Supply Wells

Constituent Group <sup>1</sup>	Number of wells sampled <sup>2</sup>	Number of wells with a concentration above an MCL <sup>3</sup>
Inorganics – Primary	1	0
Radiological	0	0
Nitrates	2	0
Pesticides	0	0
VOCs and SVOCs	0	0
Inorganics – Secondary	1	1

<sup>&</sup>lt;sup>1</sup> A description of each member in the constituent groups and a generalized discussion of the relevance of these groups are included in *California's Groundwater* – *Bulletin 118* by DWR (2003).

<sup>&</sup>lt;sup>2</sup> Represents distinct number of wells sampled as required under DHS Title 22 program from 1994 through 2000.

<sup>&</sup>lt;sup>3</sup> Each well reported with a concentration above an MCL was confirmed with a second detection above an MCL. This information is intended as an indicator of the

types of activities that cause contamination in a given basin. It represents the water quality at the sample location. It does not indicate the water quality delivered to the consumer. More detailed drinking water quality information can be obtained from the local water purveyor and its annual Consumer Confidence Report.

#### **Well Production characteristics**

Well yields (gal/min)				
Municipal/Irrigation	NKD.			
Total depths (ft)				
Domestic	Range: 22 - 325	Average: 123 (42 Well		
Municipal/Irrigation	NKD	Completion Reports)		

NKD - No known data

### **Active Monitoring Data**

Agency	Parameter	Number of wells /measurement frequency
	Groundwater levels	NKD
DWR	Miscellaneous water quality	2 wells biennial
Department of Health Services	Miscellaneous water quality	2

### **Basin Management**

Groundwater management:	The County of Siskiyou adopted a groundwater management ordinance in 1998.
Water agencies	
Public	None
Private	None

#### **Selected References**

- Cornwall HR. 1981. Chromite Deposits in the Seiad Valley and Scott Bar Quadrangles, Siskiyou County, California. USGS.
- Davis GA. 1966. Metamorphic and Granitic History of the Klamath Mountains. California Division of Mines and Geology. Bulletin 190. 39-50 p.
- Davis GA, Lipman PW. 1962. Revised Structural Sequences of Pre-Cretaceous Metamorphic Rocks in the Southern Klamath Mountains, California. Geological Society of America 73(12):1547-1552.
- Irwin WP. 1960. Geologic Reconnaissance of the Northern Coast Ranges and Klamath Mountains, California. California Division of Mines and Geology. Bulletin 179.
- Irwin WP. 1966. Geology of the Klamath Mountain Province. California Division of Mines and Geology. Bulletin 190. 19-38 p.
- Irwin WP. 1981. Tectonic Accretion of the Klamath Mountains. In: Ernst WG, editor. The Geotectonic Development of California: Prentice Hall, Inc. p 29-49.
- Pessagno EA, Blome CD. 1990. Implications of New Jurassic Stratigraphic, Geochronometric, and Paleolatitudinal Data from the Western Klamath Terrane (Smith River and Rogue Valley Subterranes). Geology.

Strand RG. 1962. Geologic Map of California [Weed Sheet]. Scale 1:250,000. California Division of Mines and Geology.

### **Bibliography**

- Bailey EH. 1966. Geology of Northern California. California Division of Mines and Geology. Bulletin 190.
- California Department of Water Resources. 1975. California's Ground Water. California Department of Water Resources. Bulletin 118.
- California Department of Water Resources. 1980. Ground Water Basins in California. California Department of Water Resources. Bulletin 118-80.
- Dickinson WR, Ingersoll RV, Grahm SA. 1979. Paleogene Sediment Dispersal and Paleotectonics in Northern California. Geological Society of America Bulletin 90:1458-1528.
- Planert M, Williams JS. 1995. Ground Water Atlas of the United States, Segment 1, California, Nevada. USGS. HA-730-B.
- Smith JG, Page NJ, Johnson MG, Moring BC, Gray F. 1982. Preliminary Geologic Map of the Medford 1 x 2 Degree Quadrangle, Oregon and California. USGS. OF-82-955.
- Strand RG. 1969. Geologic Atlas of California [Weed Sheet]. California Division of Mines and Geology.
- Wagner DL, Saucedo GJ. 1987. Geologic Map of the Weed Quadrangle, California, Regional Geologic Map Series 4A. California Division of Mines and Geology.

#### **Errata**

Changes made to the basin description will be noted here.