

SOCIETY FOR ECOSYSTEM RESTORATION
IN NORTHERN BRITISH COLUMBIA

Restoring Fish Passage in the Peace Region - 2025

PEA-F26-F-4279-DCA

**Prepared for
Fish and Wildlife Compensation Program
and
Fish Passage Technical Working Group**

**Prepared by
Al Irvine, B.Sc., R.P.Bio. and Lucy Schick, B.Sc.
New Graph Environment Ltd.
on behalf of
Society for Ecosystem Restoration in Northern BC**

**Prepared with financial support of the Fish and Wildlife Compensation Program on
behalf of its program partners BC Hydro, the Province of BC, Fisheries and Oceans
Canada, First Nations and Public Stakeholders.**

Version 0.5.1 2026-05-14

***Claude Opus 4.7 (Anthropic) assisted with literature synthesis, drafting, and technical writing. All
scientific interpretation, data analysis, and conclusions are the responsibility of the authors.***



new graph environment

Table of Contents

List of Figures	iv
Acknowledgement	vi
Executive Summary	1
1 Introduction	3
2 Background	5
3 Methods	17
4 Results and Discussion	31
5 Recommendations	45
Appendix - Fish Species by Watershed Group	47
Appendix - Assessment Data Summary	51
Appendix - Climate Departure	55
Appendix - Floodplain Delineation	79
Appendix - Phase 1 Fish Passage Assessment Data and Photos	83
Tributary To Parsnip River - 199663 - Appendix	115
Tributary To Nation River - 203597 - Appendix	125
Tributary To Williston Reservoir - 203605 - Appendix	135
Appendix - Environmental DNA Results	145
Tributary to Missinka River - 125179 - Monitoring Appendix	149
Tributary to the Table River - 125231 - Monitoring Appendix	153
Tributary to Kerry Lake - 198692 - Monitoring Appendix	157
Appendix - UAV Imagery	161
Appendix - Collaborative GIS Layers	163
References	165
Session Info	171
Attachment - Phase 1 Data and Photos	175
Attachment - Habitat Assessment and Fish Sampling Data	177

List of Figures

2.1 Overview map of Study Area	7
2.2 Hydrograph for Parsnip River above Misinchinka River (Station 07EE007)	8
2.3 Summary of historic fish observations vs. stream gradient category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.	13
2.4 Summary of historic fish observations vs. channel width category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.	15
2.5 Summary of historic fish observations vs. watershed size category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.	16
4.1 Summary of potential rearing and spawning habitat upstream of habitat confirmation assessment sites. Bull trout rearing and spawning models used for habitat estimates (total length of stream network <10.5% and <5.5% gradient, respectively)	39
4.2 Boxplots of densities (fish/100m ²) of fish captured downstream (ds) and upstream (us) by electrofishing.	41
4.3 Histograms of fish lengths by species. Fish captured by electrofishing during habitat confirmation assessments.	42
5.1 FWCP Peace Region area of interest in northeastern British Columbia, coloured by ecoregion. Williston Reservoir dominates the basin; the climate-departure pipeline is run on the regional outline (heavy blue) and again on each ecoregion polygon.	56
5.2 Annual mean temperature anomaly for the FWCP Peace Region relative to the 1951-1980 baseline. Bars are degrees above (red) or below (blue) the pre-warming average; the two trend lines are the 75-year (1951-present, dashed) and 45-year (1981-present, solid) Theil-Sen slopes.	59
5.3 Annual precipitation anomaly (% of the 1951-1980 baseline) for the FWCP Peace Region. The long-term trend is positive but not statistically significant at the regional scale; significance does appear in the two northernmost ecoregions (see Per-Ecoregion section)	60
5.4 Annual daytime maximum temperature (tmax) anomaly relative to the 1951-1980 baseline.	61
5.5 Annual overnight minimum temperature (tmin) anomaly relative to the 1951-1980 baseline.	62
5.6 Diurnal temperature range (daytime maximum minus overnight minimum) annual mean for the FWCP Peace Region. The downward slope is the day-night asymmetry — overnight lows are warming faster than daytime highs.	63
5.7 Annual peak snow water equivalent (swe_max) for the FWCP Peace Region. ERA5-Land mm snow water equivalent, regional spatial mean.	65
5.8 Day of year when half the annual snowmelt has accumulated (snowmelt_doy_50). Earlier dates indicate an earlier freshet centroid — directly relevant to crossing-design hydrology and to migration timing.	66
5.9 Annual maximum of 7-day rolling daily snowmelt (snowmelt_rate_peak). Higher values indicate more concentrated freshet pulses on the snowpack side, upstream of routing through soil and channel storage.	67
5.10 Annual snowfall fraction: the percent of annual precipitation that fell as snow. Anomaly is in percentage points relative to the 1951-1980 baseline.	68

List of Figures

5.11 Spatial pattern of annual mean temperature departure across the FWCP Peace Region (2015-2025 mean minus 1951-1980 mean, degrees C) . Warming is broad but uneven — the west-of-Rockies pattern is consistent with windward-slope amplification along the Continental Divide. . . .	69
5.12 Annual mean temperature anomaly relative to the 1951-1980 baseline, by ecoregion. Dashed line is the 75-year Theil-Sen trend (1951-present) ; solid line is the 45-year trend (1981-present) . A solid line steeper than the dashed line indicates accelerating warming.	70
5.13 Annual precipitation anomaly (% of the 1951-1980 baseline) by ecoregion. The two northernmost ecoregions (BMP, NRM) show statistically significant precipitation increases over the full record; the others do not.	71
5.14 Annual peak snow water equivalent anomaly by ecoregion, relative to the 1951-1980 baseline. Bars are mm of water-equivalent snowpack departure from baseline.	72
5.15 Snowmelt 50 % day-of-year (DOY-50) anomaly by ecoregion, relative to the 1951-1980 baseline. Negative values (red) mean the freshet midpoint shifted earlier in the year.	73
5.16 Watershed groups intersecting the FWCP Peace Region (full extent, including parts spilling outside the FWCP boundary) , labelled with their codes, on top of ecoregion fills.	75
5.17 Parsnip River Watershed Group functional floodplain (green) over hillshaded terrain. Bull trout accessible order-3+ streams in blue; lakes and wetlands in light blue; parks light green; First Nations reserves light grey with diamond markers and labels; municipalities outlined in red; forest service / resource roads grey; railways black dashed; watershed boundary heavy black.	80
5.18 South-east corner of the Parsnip River Watershed Group at full resolution — the headwaters around Arctic Lake on the continental divide. Symbology as Figure 5.17 ; named streams labelled in italic blue; municipalities outlined and labelled in red where present.	81
5.19 Map of Tributary To Parsnip River	115
5.20 Left: Typical habitat downstream of PSCIS crossing 199663. Right: Typical habitat upstream of PSCIS crossing 199663.	123
5.21 Map of Tributary To Nation River	125
5.22 Left: Typical habitat downstream of PSCIS crossing 203597. Right: Wetland habitat located 350m upstream of PSCIS crossing 203597.	133
5.23 Map of Tributary To Williston Reservoir	135
5.24 Left: Typical habitat downstream of PSCIS crossing 203605. Right: Typical habitat upstream of PSCIS crossing 203605.	143
5.25 Densities of fish (fish/100m ²) captured upstream and downstream of PSCIS crossing 125179 in 2025.	151
5.26 Densities of fish (fish/100m ²) captured upstream and downstream of PSCIS crossing 125231 in 2025.	155
5.27 Densities of fish (fish/100m ²) captured upstream and downstream of PSCIS crossing 198692 in 2025. Densities are shown where wetted-area data are available; some surveys at this site had no recorded width.	158

Acknowledgement

At New Graph Environment, we understand our well-being as inseparable from the health of the land and waters we work within. When we care for ecosystems, we care for ourselves and for the communities connected to them.

Executive Summary

 [Executive Summary \(PDF\)](#)

This report is available as a PDF and as an online interactive report at https://www.newgraphenvironment.com/fish_passage_peace_2025_reporting. We recommend viewing online as the web-hosted HTML version contains more features and is more easily navigable. Please reference the website for the latest version stamped PDF from [fish_passage_peace_2025_reporting.pdf](#).

The Society for Ecosystem Restoration in Northern BC (SERNbc) is working together with the McLeod Lake Indian Band, the Peace Region Fish and Wildlife Compensation Program (FWCP), the Provincial Fish Passage Technical Working Group (FPTWG), road/rail tenure holders and other stakeholders/partners to prioritize, plan and fund the restoration of fish passage at road crossing structure barriers within the FWCP Peace Region.

The primary objective of this project is to identify and prioritize fish passage barriers within study areas, develop comprehensive restoration plans to address these barriers, and foster momentum for broader ecosystem restoration initiatives. While the primary focus is on fish passage, this work also serves as a lens through which to view the broader ecosystems, leveraging efforts to build capacity for ecosystem restoration and improving our understanding of watershed health. We recognize that the health of life - such as our own - and the health of our surroundings are interconnected, with our overall well-being dependent on the health of our environment.

Although the main purpose of this report is to document 2025 field work data and results, it also builds on multiple prior years of SERNbc fish passage reporting; per-site links to the relevant prior reports are in the [Assessment Data Summary \(page 51\)](#) appendix.

Fish passage assessment procedures conducted through SERNbc in the FWCP Peace Region since 2020 are amalgamated online within the Results and Discussion section of the report found [here](#), which includes links to project reporting for each site.

In 2025, fieldwork activities took place within the Parsnip River, Parsnip Arm, Carp River, and Nation River watershed groups. Fish passage assessments and habitat confirmations were completed at new and previously identified crossings, with site-level results documented per-crossing in the appendices. Two crossings advanced to FWCP-funded 2026 implementation following this year's work: PSCIS 199663 (Kennedy Siding, tributary to the Parsnip River — replacement with a streambed simulation in coordination with BC Timber Sales) and PSCIS 203597 (THUT15000 Deactivation, tributary to the Nation River — riparian enhancements including soil decompaction, native planting, and coarse woody debris placement).

Environmental DNA (eDNA) sampling was incorporated at program scale this year. A total of 35 samples were collected across 10 streams in partnership with the University of Northern British Columbia (UNBC) for ddPCR analysis. Per-species detection results informed crossing prioritization and partner conversations, and are summarized in [Appendix - Environmental DNA Results \(page 145\)](#) and visualized in the interactive Peace eDNA map.

Fish sampling and PIT-tagging was completed at three monitoring sites using a custom fish-passage effectiveness monitoring form standardized across the program. Ongoing effectiveness monitoring continued at PSCIS 125179 (Tributary to Missinka River) and PSCIS 125131 (Tributary to Table River), extending multi-year baselines from prior reports; baseline monitoring was initiated this year at PSCIS 198692 (Kerry Lake) ahead of planned remediation. eDNA sampling was integrated at all three sites as part of both the baseline and the ongoing effectiveness components of the monitoring protocol.

Two analytical dimensions are new to the program this year. A floodplain delineation pilot was completed for the Parsnip River Watershed Group using the [flooded](#) R package, mapping the functional floodplain footprint and the off-channel habitat units (lakes, wetlands, side channels) that lateral connectivity supports — see [Appendix - Floodplain Delineation \(page 79\)](#). A climate departure analysis was completed for the FWCP Peace Region using the [cd](#) R package, quantifying temperature, precipitation, snowpack, and atmospheric drying trends since 1951 across the region's ecoregions — see [Appendix - Climate Departure \(page 55\)](#). Both layers complement the structural barrier inventory by providing the watershed context that prioritization decisions depend on.

Fish passage work in this region operates inside the constraints described in the [Approach \(page 5\)](#) section of the background: thousands of structures across multiple jurisdictions and tenure holders, six-to-eight-figure remediation costs, and a finite funding envelope. Decisions about which crossings to advance are made opportunistically — pursuing simpler, lower-cost options to maintain momentum, while building the multi-year relationships and shared data that larger commitments require. The breadth of 2025 partner engagement is summarised in [Engage Partners \(page 31\)](#).

1 Introduction

This report is available as a PDF and as an online interactive report at https://www.newgraphenvironment.com/fish_passage_peace_2025_reporting. We recommend viewing online as the web-hosted HTML version contains more features and is more easily navigable. Please reference the website for the latest PDF from [fish_passage_peace_2024_reporting.pdf](#).

The Society for Ecosystem Restoration in Northern BC (SERNbc) is working together with the McLeod Lake Indian Band, the Peace Region Fish and Wildlife Compensation Program (FWCP), the Provincial Fish Passage Technical Working Group (FPTWG), road/rail tenure holders and other stakeholders/partners to prioritize, plan and fund the restoration of fish passage at road crossing structure barriers within the FWCP Peace Region.

The primary objective of this project is to identify and prioritize fish passage barriers within study areas, develop comprehensive restoration plans to address these barriers, and foster momentum for broader ecosystem restoration initiatives. While the primary focus is on fish passage, this work also serves as a lens through which to view the broader ecosystems, leveraging efforts to build capacity for ecosystem restoration and improving our understanding of watershed health. We recognize that the health of life - such as our own - and the health of our surroundings are interconnected, with our overall well-being dependent on the health of our environment.

This project builds on prior years of Society for Ecosystem Restoration Northern BC (SERNbc) fish passage reporting (Irvine 2020; Irvine 2022; Irvine and Winterscheidt 2023, 2024; Irvine and Schick 2025); per-site links to the relevant prior reports are in the [Assessment Data Summary \(page 51\)](#) appendix.

The health and viability of freshwater fish populations can depend on access to tributary and off channel areas which provide refuge during high flows, opportunities for foraging, overwintering habitat, spawning habitat and summer rearing habitat (Bramblett et al. 2002; Swales and Levings 1989; Diebel et al. 2015). Culverts can present barriers to fish migration due to low water depth, increased water velocity, turbulence, a vertical drop at the culvert outlet and/or maintenance issues (Slaney and Zaldokas 1997; Cote et al. 2005). As road crossing structures are commonly upgraded or removed there are numerous opportunities to restore connectivity by ensuring that fish passage considerations are incorporated into repair, replacement, relocation and deactivation designs.

Although remediation and replacement of stream crossing structures can have benefits to local fish populations, the costs of remedial works can be significant and the impacts of the work often complex to evaluate and quantify. Additionally, allocation of ecosystem restoration funding towards infrastructure upgrades on transportation right of ways are not always considered ethical under all

1 Introduction

circumstances from all perspectives. When funds are finite and invested groups are engaged in fund raising, cost benefits and the ethics of crossing replacements should be explored collaboratively alongside the cost benefits and ethics of alternative investment activities including transportation corridor relocation/deactivation, land procurement/covenant, cattle exclusion, riparian/floodplain restoration, habitat complexing, water conservation, commercial/recreational fishing management, salt water interventions and research.

Please note that at the time of reporting this document can be considered a living document. See the [Open Source Reporting Framework \(page 28\)](#) section for details on version tracking and project documentation.

2 Background

2.1 Approach

Fish passage assessments in British Columbia follow the provincial protocol (MoE 2011), which scores barrier risk at road-stream crossings from structural parameters — culvert embedment, slope, outlet drop, and culvert diameter relative to channel width. The “Barrier” and “Potential Barrier” classifications represent passability for juvenile salmon or small resident rainbow trout under any flow conditions that may occur throughout the year (Clarkin et al. 2005; Bell 1991; Thompson 2013). The thresholds are deliberately conservative, so the classifications are risk scores rather than determinations of whether fish actually move through. Many structures classified as “Barrier” still pass some species and life stages at some flows, and natural features in adjacent reaches can be as limiting as the structure itself. Passability can be quantified in other ways — fish length, swim speed, and species physiology all affect connectivity outcomes — and finer-grained scoring methods exist for structures already classified as barriers (Bourne et al. 2011; Kemp and O’Hanley 2010; Washington Department of Fish & Wildlife 2009). Whether a remediation changes outcomes for fish therefore depends on the habitat upstream, the species and life stages in question, and the broader condition of the reach the structure connects to — not the classification alone.

The provincial inventory contains thousands of structures on fish-bearing streams. Replacement at a typical forest road culvert is in the six-figure range; structural replacements on highway and rail corridors run into seven and eight figures. Funding for restoration is finite and split across multiple programs and partners. Prioritization across the inventory therefore carries more weight than any individual assessment — the assessment establishes what could be fixed, while prioritization decides what should be.

Authority and funding for these structures are distributed across rail authorities, highway ministries, forestry licensees, and private landowners. Each operates under its own mandate, performance metrics, and budget cycle. Advancing restoration involves working inside organizations whose primary mandate is moving goods, maintaining roads, or harvesting timber — not aquatic ecological outcomes. The work proceeds through shared data, sustained relationships, and incremental progress across multiple budget years and field seasons.

Many road and rail corridors were built without consideration of fish passage or floodplain function. Damage frequently extends beyond the structure at the crossing — channelized streams, dykes that sever lateral connection, drained wetlands, cleared riparian buffers. Replacing a culvert in-kind can restore longitudinal passage while leaving the broader corridor-scale disconnection unaddressed. Understanding where infrastructure constrains watershed function helps distinguish cases where structural replacement is sufficient from cases where remediation needs to address more than the structure itself.

2 Background

This report compiles the layers used for prioritization in the watershed groups where the work is active: structural assessment data, habitat modelling, climate trajectory, and floodplain extent. Each layer contributes a piece of the picture — what is present, what it could support, how it is changing, and the surrounding watershed context. The work happens over many years across many partners; this report documents one year's contribution to that effort.

2.2 Project Location

The study area includes the FWCP Peace Region with a focus to date on traditional territories of the Tse'khene First Nations. In 2025, field assessments were completed with the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups (Figure [2.1](#)).

In 2019/2020, following a literature review, analysis of fish habitat modelling data, the Provincial Stream Crossing Inventory System (PSCIS) and a community scoping exercise within the McLeod Lake Indian Band habitat confirmation assessments were conducted at 17 sites throughout the Parsnip River watershed with 10 crossings rated as high priorities for rehabilitation and three crossings rated as moderate priorities for restoration. An engineering design for site 125179 on a tributary to the Missinka River was also completed through the 2019/2020 project. In 2021/2022, project activities reconvened through FWCP directed project PEA-F22-F-3577-DCA. Partners were engaged, funding was raised, planning was conducted and reporting was completed to initiate restoration activities of high priority crossings. Materials were purchased and permitting was put in place to prepare for replacement of the twin culverts on the Missinka River tributary with a clear-span bridge. In 2025/2026, this collaborative project leveraged ongoing connectivity restoration initiatives in the province and engaged multiple partners to catalyze fish passage restoration activities at high-priority sites identified in from 2019-2024.

2.3 Tse'khene




Figure 2.1: Overview map of Study Area

2.3 Tse'khene

The Parsnip River watershed is located within the south-eastern portion of the 108,000 km² traditional territory of the Tse'khene from the [McLeod Lake Indian Band](#). The Tse'khene "*People of the Rocks*" are a south westerly extension of the Athabaskan speaking people of northern Canada. They were nomadic hunters whose language belongs to the Beaver-Sarcee-Tse'khene branch of Athapaskan ("History Who We Are" 2023). Extensive work is underway to preserve the language with resources such as First Voices available [online](#) and in [app form](#) for iphone and ipad devices.

The continental divide separates watersheds flowing north into the Arctic Ocean via the Mackenzie River and south and west into the Pacific Ocean via the Fraser River (Figure 2.1). The Parsnip River is a 6th order stream with a watershed that drains an area of 5597km². The mainstem of the river flows within the Rocky Mountain Trench in a north direction into Williston Reservoir starting from the continental divide adjacent to Arctic Lakes. Major tributaries include the Misinchinka, Colbourne, Reynolds, Anzac, Table, Hominka and Missinka sub-basins which drain the western slopes of the Hart Ranges of the Rocky Mountains. The Parsnip River has a mean annual discharge of 150.5 m³/s with flow patterns typical of high elevation watersheds on the west side of

2 Background

the northern Rocky Mountains which receive large amounts of precipitation as snow leading to peak levels of discharge during snowmelt, typically from May to July (Figure 2.2).

Construction of the 183m high and 2134m long W.A.C. Bennett Dam was completed in 1967 at Hudson's Hope, BC, creating the Williston Reservoir (Hirst 1991). Filling of the 375km² reservoir was complete in 1972 and flooded a substantial portion of the Parsnip River and major tributary valleys forming what is now known as the Peace and Parsnip reaches. The replacement of riverine habitat with an inundated reservoir environment resulted in profound changes to the ecology, resource use and human settlement patterns in these systems (Hagen et al. 2015; Pearce et al. 2019; Stamford et al. 2017). Prior to the filling of the reservoir, the Pack River, into which McLeod Lake flows, was a major tributary to the Parsnip River. The Pack River currently enters the Williston Reservoir directly as the historic location of the confluence of the two rivers lies within the reservoir's footprint.




Figure 2.2: Hydrograph for Parsnip River above Misinchinka River (Station #07EE007).

2.4 Floodplains and Watershed Function

Floodplains are the low-lying ground adjacent to stream channels that water reclaims during high-flow events. They are not idle margins — they are where rivers do geomorphic work: storing and sorting sediment, exchanging nutrients between terrestrial and aquatic systems, recruiting riparian vegetation, and delivering large woody debris to the channel. For fish, floodplain habitats — side channels, sloughs, beaver complexes, oxbow ponds, and connected wetlands — provide the off-channel conditions that drive freshwater productivity. Juvenile salmonids depend on these areas for low-velocity rearing, overwinter refuge when mainstem conditions become harsh, and high-flow shelter during freshet. In many systems, floodplain habitats are where freshwater survival is won or lost.

Floodplains are often the part of the watershed most compromised by the infrastructure legacy described above. Dykes sever lateral connection; channelization confines the active footprint; drainage eliminates wetland complexes. Mapping floodplain extent at watershed scale therefore complements the linear barrier inventory — prioritization can consider not just whether a structure passes fish but the condition and extent of the off-channel habitat in the reach the structure connects to.

2.5 Fisheries

Fish species recorded in the BC Fisheries Information Summary System (FISS) for the watershed groups in this study area are summarised in [Appendix - Fish Species by Watershed Group \(page 47\)](#) (MoE 2024a). In addition to flooding related to the formation of the Williston Reservoir, transmission lines, gas pipelines, rail, road networks, forestry, elevated water temperatures, interrupted connectivity, invasion from non-native species and insect infestations affecting forested areas pose threats to fisheries values in the Parsnip River watershed (Hagen et al. 2015; Stamford et al. 2017; Hagen and Weber 2019; COSEWIC 2012). A brief summary of trends and knowledge status related to Arctic grayling, bull trout, kokanee, mountain whitefish and rainbow trout in Williston Watershed streams is provided in Fish and Wildlife Compensation Program (2020) with a more detailed review of the state of knowledge for Parsnip River watershed populations of Arctic grayling and bull trout provided below.

Bull Trout - sa'ba

Tse'khene Elders from the McLeod Lake Indian Band report that sa'ba (bull trout) size and abundance has decreased in all rivers and tributaries from the reservoir with more injured and diseased fish captured in recent history than was common in the past (Pearce et al. 2019) .

Bull trout populations of the Williston Reservoir watershed are included within the Western Arctic population 'Designatable Unit 10', which, in 2012, received a ranking of 'Special Concern' by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC 2012). They were added to Schedule 1 under the Species at Risk Act in 2019 (Species Registry Canada 2020) and are also

2 Background

considered of special concern (blue-listed) provincially (BC Species & Ecosystem Explorer 2020). Some or all of the long-term foot survey index sections of four Williston Reservoir spawning tributaries (Davis Creek, Misinchinka River, Point Creek, and Scott Creek), have been surveyed within 16 of the 19 years between 2001 and 2019 (16 of 19 in Davis River, 10 years over a 13-year period in the Misinchinka River, 11 years over a 14-year period for Point Creek, and 9 years over an 11-year period for Scott Creek (Hagen et al. 2020).

A study of bull trout critical habitats in the Parsnip River was conducted in 2014 with the Misinchinka and Anzac systems identified as the most important systems for adfluvial (large bodied) bull trout spawners. The Table River was also highlighted as an important spawning destination. Other watersheds identified as containing runs of large bodied bull trout spawners included the Colbourne, Reynolds, Hominka and Missinka River with potentially less than 50 spawners utilizing each sub-basin (Hagen et al. 2015). Hagen and Weber (2019) have synthesized a large body of information regarding limiting factors, enhancement potential, critical habitats and conservation status for bull trout of the Williston Reservoir and the reader is encouraged to review this work for context. They have recommended experimental enhancements within a monitoring framework for Williston Reservoir bull trout (some spawning and rearing in Parsnip River mainstem and tributaries) which include stream fertilization, side channel development, riparian restoration and fish access improvement.

In 2018, sub-basins of the Anzac River watershed, Homininka River, Missinka River and Table River watersheds were designated as fisheries sensitive watersheds under the authority of the *Forest and Range Practices Act* due to significant downstream fisheries values and significant watershed sensitivity (Beaudry 2013a, 2014, 2013b; P. Beaudry 2014). Special management is required in these watersheds to protect habitat for fish species including bull trout and Arctic grayling including measures (among others) to limit equivalent clearcut area, reduce impacts to natural stream channel morphology, retain old growth attributes and maintain fish habitat/movement (Forest and Range Practices Act 2018).

Arctic Grayling - dusk'ihje

Tse'khene Elders from the McLeod Lake Indian Band report that dusk'ihje (Arctic grayling) numbers have declined dramatically since the flooding of the reservoir and that few dusk'ihje have been caught in the territory in the past 30 years (Pearce et al. 2019).

Since impoundment of the Williston Reservoir, it appears that physical habitat and ecological changes have been the most significant factors limiting Arctic grayling productivity. Although these changes are not well understood they have likely resulted in the inundation of key low gradient juvenile rearing and overwintering habitats, isolation of previously connected populations and increases in abundance of predators such as bull trout (Shrimpton et al. 2012; Hagen et al. 2018). Rapid increases in industrial activity and angler access in the Parsnip River watershed pose

2.5 Fisheries

significant risks to Arctic Grayling productivity with these threats primarily linked to forestry and pipeline initiatives (Hagen and Stamford 2021).

A detailed review of Arctic grayling life history is available in Stamford et al. (2017). Migration of mature adult Arctic grayling occurs in the spring, typically coinciding with water temperatures of 4°C. In the Parsnip watershed, spawning appears to occur between late May and late June, primarily within the lower reaches of the Anzac and Table rivers and the Parsnip River mainstem. Spawning habitat is generally associated with side-channel and multi-channel areas containing small gravels. The current distribution of Williston Arctic grayling is concentrated in fourth-order and larger streams (Williamson and Zimmerman 2005; Stamford et al. 2017). According to Stewart et al. (2007), Arctic grayling spawn in both large and small tributaries to rivers and lakes, intermittent streams, mainstem rivers, and lakes—most commonly at tributary mouths. Although earlier studies suggest that 0+ grayling overwinter in the lower reaches of larger tributaries (e.g., Table and Anzac rivers) and the Parsnip River, and that few age-1+ grayling have been sampled in tributaries, habitat use in small tributaries and the extent to which they are connected with mainstem habitats across all core areas remains poorly understood. Between 1995 and 2019, population monitoring of Arctic grayling was conducted in the Table River in 9 out of 25 years, and in the Anzac River for 8 years, using snorkel surveys. Results from 2018 and 2019 support efforts to assess the conservation status of the species in the Parsnip Core Area (Hagen et al. 2018). Preliminary telemetry data from 2019 indicate that both Arctic grayling and bull trout rely on the Parsnip River mainstem for overwinter residency. Arctic grayling typically begin moving into tributaries in April and are widespread across the watershed by June, with larger adults observed in headwater habitats by August.

A five-year study on Arctic grayling abundance and trends in the Parsnip River watershed is presented in Hagen and Stamford (2023), which reports that the most productive summer rearing habitats for Arctic grayling are within the Anzac and Table rivers. Although estimated abundance is lower than in the Anzac and Table, productive summer rearing habitats for adult Arctic grayling in the upper Parsnip River watershed are distributed between 36–25km of the Missinka River and 48–32km of the Hominka River. In the Anzac River, Hagen and Stamford (2021) report that a 30km stretch from 47km to 16km supports high Arctic grayling abundance and is assumed to provide productive rearing habitat for adults. A chute obstruction located at 47km likely limits further upstream movement. Although the spatial distribution of high Arctic grayling abundance in the Table River has not been delineated through reconnaissance surveys, it has been observed to span at least a 20km zone from the waterfall migration barrier at 37km to 18km.

Spatial ecology studies in the Parsnip River between 2018 and 2021 has been reported on by Martins et al. (2022) with results related to:

- temperature modeling and spatio-temporal patterns in thermal habitat,
- telemetry data modeling and arctic grayling spatial ecology, and
- trophic relationships between Arctic grayling and bull trout

Fish presence by Habitat Type

A review of available fisheries data for the Parsnip River watershed stratified by different habitat characteristics (stream gradient, channel width, and watershed size) can provide insight into which habitats may provide the highest intrinsic value for fish species based on the number of fish captured in those habitats in past assessment work (Tables [2.1](#) - [2.3](#) and Figures [2.3](#) - [2.5](#)). It should be noted however that it should not be assumed that all habitat types have been sampled in a non-biased fashion or that particular sites selected do not have a disproportionate influence on the overall dataset composition (ie. fish salvage sites are often located adjacent to construction sites which are more commonly located near lower gradient stream reaches).

Table 2.1: Summary of historic fish observations vs. stream gradient category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

species_code	Gradient	Count	total_spp	Percent
BT	0 - 3 %	411	614	67
BT	03 - 5 %	86	614	14
BT	05 - 8 %	58	614	9
BT	08 - 15 %	50	614	8
BT	15 - 22 %	9	614	1
GR	0 - 3 %	249	251	99
GR	03 - 5 %	2	251	1
KO	0 - 3 %	145	150	97
KO	03 - 5 %	2	150	1
KO	05 - 8 %	2	150	1
KO	22+ %	1	150	1
RB	0 - 3 %	2035	2619	78
RB	03 - 5 %	275	2619	11
RB	05 - 8 %	186	2619	7
RB	08 - 15 %	103	2619	4
RB	15 - 22 %	14	2619	1
RB	22+ %	6	2619	0

2.5 Fisheries




Figure 2.3: Summary of historic fish observations vs. stream gradient category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

Table 2.2: Summary of historic fish observations vs. channel width category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

species_code	Width	Count	total_spp	Percent
BT	0 - 2m	30	614	5
BT	02 - 04m	96	614	16
BT	04 - 06m	81	614	13
BT	06 - 10m	82	614	13
BT	10 - 15m	59	614	10
BT	15m+	248	614	40
BT	—	18	614	3
GR	02 - 04m	6	251	2
GR	06 - 10m	6	251	2

2 Background

species_code	Width	Count	total_spp	Percent
GR	10 - 15m	6	251	2
GR	15m+	231	251	92
GR	–	2	251	1
KO	0 - 2m	4	150	3
KO	02 - 04m	22	150	15
KO	04 - 06m	3	150	2
KO	06 - 10m	11	150	7
KO	10 - 15m	5	150	3
KO	15m+	31	150	21
KO	–	74	150	49
RB	0 - 2m	656	2619	25
RB	02 - 04m	520	2619	20
RB	04 - 06m	257	2619	10
RB	06 - 10m	222	2619	8
RB	10 - 15m	122	2619	5
RB	15m+	454	2619	17
RB	–	388	2619	15

2.5 Fisheries




Figure 2.4: Summary of historic fish observations vs. channel width category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

Table 2.3: Summary of historic fish observations vs. watershed size category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

species_code	Watershed	count_wshd	total_spp	Percent
BT	0 - 25km2	252	614	41
BT	25 - 50km2	93	614	15
BT	50 - 75km2	40	614	7
BT	75 - 100km2	11	614	2
BT	100km2+	218	614	36
GR	0 - 25km2	8	251	3
GR	50 - 75km2	2	251	1
GR	75 - 100km2	4	251	2
GR	100km2+	237	251	94

2 Background

species_code	Watershed	count_wshd	total_spp	Percent
KO	0 - 25km2	105	150	70
KO	25 - 50km2	6	150	4
KO	50 - 75km2	5	150	3
KO	75 - 100km2	3	150	2
KO	100km2+	31	150	21
RB	0 - 25km2	1802	2619	69
RB	25 - 50km2	171	2619	7
RB	50 - 75km2	97	2619	4
RB	75 - 100km2	48	2619	2
RB	100km2+	501	2619	19




Figure 2.5: Summary of historic fish observations vs. watershed size category for the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups.

3 Methods

3.1 Climate Departure

Climate departure measures how far recent climate has shifted from a historical baseline — in this case, how much warmer, drier, or snowpack-altered the FWCP Peace Region has become relative to the pre-warming period (1951–1980). Where traditional fish-passage assessments focus on the physical structures in and around streams, climate departure adds the atmospheric and hydrologic context those structures sit inside: warming air temperatures that drive warmer water, shifting snowpack that reshapes the freshet, and atmospheric drying that reduces late-summer flows. Understanding these departures at the watershed scale lets us ask whether the habitat above a barrier is gaining or losing value as climate changes — and weight barrier prioritisation accordingly.

We examined 15 climate variables spanning temperature (annual mean, daytime maximum, overnight minimum), moisture (precipitation, vapour pressure deficit, soil moisture, relative humidity), and snowpack (snow water equivalent, snowfall, snowmelt, snow cover, snowfall fraction, peak snow water equivalent, snowmelt timing, and peak melt rate). Each variable was assessed at annual and seasonal scales, across the full region and within each of five ecoregions.

Climate departures were computed with the [cd](#) R package, which reads ERA5-Land hourly reanalysis (1950–present, ~9 km native grid) aggregated to monthly, seasonal, and annual periods on a single grid covering British Columbia, served as Cloud-Optimised GeoTIFFs from a public S3 catalog. For each variable the package crops the published rasters to the area of interest, takes a yearly spatial mean, and computes anomalies against the 1951–1980 pre-warming reference period — the same base period Hansen et al. (2012) used for global temperature emergence. Long-term trends are tested with Mann-Kendall significance and quantified with Theil-Sen slopes; the recent-decade (2015–2025) versus pre-warming window comparison uses a Welch two-sample t-test on the annual values. The same pipeline runs at any polygon scale — region, ecoregion, or watershed group. The regional and ecoregion-level analyses are summarised in this section; the watershed-group breakdown feeds into the per-watershed barrier prioritisation that follows. [Appendix - Climate Departure \(page 55\)](#) carries the full set of plots and tables.

3.2 Planning — Habitat, Connectivity and Floodplain Modelling

3.2.1 Habitat Modelling

Habitat modelling used to help guide planning for field assessments is generated by `bcfishpass` (Norris [2020] 2024) which has been designed to prioritize potential fish passage barriers for assessment or remediation by generating a simple model of aquatic habitat connectivity. We utilize the `bcfishpass` access model, linear spawning/rearing habitat model and lateral habitat connectivity for planning purposes. These models provide a valuable starting point, but their results are not definitive and should always be considered with professional judgment. Detailed information regarding model methodology, select parameters and known model limitations are detailed in Norris ([2020] 2024) with key documentation linked below:

- [Access model](#)
- [Linear spawning/rearing habitat models](#)
- [Lateral habitat model](#)

3 Methods

Table 3.1 documents the custom species-specific thresholds for stream gradient and channel width applied to the linear spawning and rearing habitat model for this year’s project planning.

Table 3.1: Stream gradient and channel width thresholds used to model potentially highest value fish habitat.

Variable	Bull Trout	Arctic Grayling
Spawning Gradient Max (%)	5.5	2.5
Spawning Width Min (m)	2	4
Rearing Width Min (m)	1.5	1.5
Rearing Gradient Max (%)	10.5	3.5
*		

3.2.1.1 Statistical Support for bcfishpass Fish Habitat Modelling Updates

This project provided the statistical background for updates to bcfishpass that facilitated incorporation of channel width (observed or predicted) into species specific linear spawning/rearing habitat models. In early 2021, Bayesian statistical methods were developed to predict channel width in all provincial freshwater atlas stream segments where width measurements had not previously been measured in the field. The model was based on the relationship between watershed area and mean annual precipitation weighted by upstream watershed area (Thorley and Irvine 2021). In December of 2021, Thorley and Irvine (2021) methods were updated using a power model derived by Finnegan et al. (2005) which relates stream discharge to watershed area and mean annual precipitation resulting in Thorley et al. (2021) which was utilized for channel width estimates within bcfishpass modelling at the time of reporting. More detailed documentation of the methodology used to facilitate both the data collection and statistical analysis can be sourced in Irvine ([2021] 2022) and Thorley et al. (2021).

In 2024, in collaboration with Poisson Consulting - stream discharge and temperature causal effects pathways were mapped with the intent of focusing aquatic restoration actions in areas of highest potential for positive impacts on fisheries values (ie. elimination of areas from intrinsic models where water temperatures are likely too cold to support fish production). The project began with a custom mechanistic model (visually represented [here](#)), but the model struggled to converge. The project then shifted to the air2stream model, which offers a middle ground between fully mechanistic models—often data-intensive and reliant on quantities that are difficult to measure or estimate—and purely statistical models, which lack physical justification and perform poorly when extrapolated to new conditions (Toffolon and Piccolroaz (2015)). After several adaptations, the expected stream temperatures were best modeled using the four-parameter version of the air2stream model, with added random effects by site for each of the four parameters (Hill et al. (2024)). The data used for the model were sourced from the following locations, for years 2019-2021:

- Water temperature data collected in the Nechako Watershed were downloaded from [Zenodo](#) (Gilbert et al. 2022).
- Hourly air temperature data were obtained from the ERA-5-Land dataset via the Copernicus Climate Change Service (Muñoz Sabater (2019))
- Daily baseflow and surface runoff data were sourced from the [Pacific Climate Impacts Consortium's Gridded Hydrologic Model Output](#) using the ACCESS1-0_rcp85 scenario (Pacific Climate Impacts Consortium (n.d.)).

3.2.2 Bull Trout and Arctic Grayling Critical Habitat

We collaborated with Eduardo Martins (Associate Professor at UNBC) to incorporate Arctic grayling into our collaborative GIS project so it could be used to inform field work activities and potential remediation site prioritization. Similarly, critical bull trout habitat data was added to the GIS project after being obtained from Luc Turcotte (WLRS - Regional Aquatic Specialist) and stored securely. Bull trout data acquisition required prior completion of the Access to Secure Species and Ecosystem Data and Information (SEDIS) training conducted through WLRS.

3.2.3 Floodplain Delineation

To complement the linear barrier inventory, we mapped functional floodplain extent at watershed scale for the Parsnip River Watershed Group. Overlaying floodplain footprint on the crossing inventory lets us ask not just whether fish can pass a structure, but how much off-channel habitat sits within the affected corridor and whether infrastructure has compromised the lateral connectivity that sustains it.

We used the [flooded](#) R package (Nagel et al. 2014), which models the area a river actively inundates during regular high-flow events — the **functional floodplain**. This is distinct from the narrow active channel (the wetted perimeter at low flow) and the broader geologic valley bottom (which includes terraces untouched by water for centuries). The functional floodplain is the scale where off-channel habitat is created and maintained: where side channels form, seasonal flooding sustains wetlands, and riparian forests recruit. Bankfull depth is estimated from the regional regression of Hall et al. (2007), calibrated against field-mapped floodplain sites in Pacific Northwest streams. Inputs were the [MRDEM-30](#) 30 m DEM, the bull trout accessible stream network (stream order ≥ 3) from [bcfishpass](#), and BC Freshwater Atlas lakes and wetlands on that network. Full methodology and watershed-scale mapping are in [Appendix - Floodplain Delineation \(page 79\)](#).

3.3 Fish Passage Assessments

3.3.1 Natural Barriers to Fish Passage

Our assessments may include natural features such as waterfalls that could limit fish passage. This informs whether upstream culvert upgrades would restore access for anadromous species (e.g., salmon) or primarily benefit resident fish already upstream. We document these features by measuring height, gradient, and pool depth, recording field observations, capturing site photographs, and reviewing background sources for context.

3.3.2 Road Stream Crossings

In the field, crossings prioritized for follow-up were first assessed for fish passage following the procedures outlined in “Field Assessment for Determining Fish Passage Status of Closed Bottomed Structures” (MoE 2011). The reader is referred to (MoE 2011) for detailed methodology. Crossings surveyed included closed bottom structures (CBS), open bottom structures (OBS) and crossings considered “other” (i.e. fords). Photos were taken at surveyed crossings and when possible included images of the road, crossing inlet, crossing outlet, crossing barrel, channel downstream and channel upstream of the crossing and any other relevant features. The following information was recorded for all surveyed crossings: date of inspection, crossing reference, crew member initials, Universal Transverse Mercator (UTM) coordinates, stream name, road name and kilometer, road tenure information, crossing type, crossing subtype, culvert diameter or span for OBS, culvert length or width for OBS. A more detailed “full assessment” was completed for all closed bottom structures and included the following parameters: presence/absence of continuous culvert embedment (yes/no), average depth of embedment, whether or not the culvert bed resembled the native stream bed, presence of and percentage backwatering, road fill depth, outlet drop, outlet pool depth, inlet drop, culvert slope, average downstream channel width, stream slope, presence/absence of beaver activity, presence/absence of fish at time of survey, type of valley fill,

3.3 Fish Passage Assessments

and a habitat value rating. Habitat value ratings were based on channel morphology, flow characteristics (perennial, intermittent, ephemeral), fish migration patterns, the presence/absence of deep pools, un-embedded boulders, substrate, woody debris, undercut banks, aquatic vegetation and overhanging riparian vegetation (Table [3.2](#)).

Table 3.2: Habitat value criteria (Fish Passage Technical Working Group, 2011).

Habitat Value	Fish Habitat Criteria
High	The presence of high value spawning or rearing habitat (e.g., locations with abundance of suitably sized gravels, deep pools, undercut banks, or stable debris) which are critical to the fish population.
Medium	Important migration corridor. Presence of suitable spawning habitat. Habitat with moderate rearing potential for the fish species present.
Low	No suitable spawning habitat, and habitat with low rearing potential (e.g., locations without deep pools, undercut banks, or stable debris, and with little or no suitably sized spawning gravels for the fish species present).

3 Methods

Fish passage potential was determined for each stream crossing identified as a closed bottom structure as per MoE (2011). The combined scores from five criteria: depth and degree to which the structure is embedded, outlet drop, stream width ratio, culvert slope, and culvert length were used to screen whether each culvert was a likely barrier to some fish species and life stages (Table 3.3, Table 3.4). These criteria were developed based on data obtained from various studies and reflect an estimation for the passage of a juvenile salmon or small resident rainbow trout (Clarkin et al. 2005; Bell 1991; Thompson 2013). For crossings determined to be potential barriers or barriers based on the data, a culvert fix and recommended diameter/span was proposed.

Table 3.3: Fish Barrier Risk Assessment (MoE 2011).

Risk	LOW	MOD	HIGH
Embedded	>30cm or >20% of diameter and continuous	<30cm or 20% of diameter but continuous	No embedment or discontinuous
Value	0	5	10
Outlet Drop (cm)	<15	15-30	>30
Value	0	5	10
SWR	<1.0	1.0-1.3	>1.3
Value	0	3	6
Slope (%)	<1	1-3	>3
Value	0	5	10
Length (m)	<15	15-30	>30
Value	0	3	6

Table 3.4: Fish Barrier Scoring Results (MoE 2011).

Cumulative Score	Result
0-14	passable
15-19	potential barrier
>20	barrier

The habitat gain index is the quantity of modelled habitat upstream of the subject crossing and represents an estimate of habitat gained with remediation of fish passage at the crossing. For this project, a gradient threshold between accessible and non-accessible habitat was set at 25% (for a minimum length of 100m) intended to represent the maximum gradient of which the strongest swimmers of anadromous species (bull trout) are likely to be able to migrate upstream.

3.3 Fish Passage Assessments

For reporting of Phase 1 - fish passage assessments within the body of this report (Table 3.3), a “total” value of habitat <20% output from *bcfishpass* was used to estimate the amount of habitat upstream of each crossing less than 25% gradient before a falls of height >5m - as recorded in MoE (2024b) or documented in other *bcfishpass* online documentation. For Phase 2 - habitat confirmation sites, conservative estimates of the linear quantity of habitat to be potentially gained by fish passage restoration, bull trout rearing maximum gradient threshold (10.5%) was used. To generate estimates for area of habitat upstream (m^2), the estimated linear length was multiplied by half the downstream channel width measured (overall triangular channel shape) as part of the fish passage assessment protocol. Although these estimates are not generally conservative, have low accuracy and do not account for upstream stream crossing structures they allow a rough idea of the best candidates for follow up.

Potential options to remediate fish passage were selected from MoE (2011) and included:

- Removal (RM) - Complete removal of the structure and deactivation of the road.
- Open Bottom Structure (OBS) - Replacement of the culvert with a bridge or other open bottom structure. Based on consultation with FLNR road crossing engineering experts, for this project we considered bridges as the only viable option for OBS type .
- Streambed Simulation (SS) - Replacement of the structure with a streambed simulation design culvert. Often achieved by embedding the culvert by 40% or more. Based on consultation with FLNR engineering experts, we considered crossings on streams with a channel width of <2m and a stream gradient of <8% as candidates for replacement with streambed simulations.
- Additional Substrate Material (EM) - Add additional substrate to the culvert and/or downstream weir to embed culvert and reduce overall velocity/turbulence. This option was considered only when outlet drop = 0, culvert slope <1.0% and stream width ratio < 1.0.
- Backwater (BW) - Backwatering of the structure to reduce velocity and turbulence. This option was considered only when outlet drop < 0.3m, culvert slope <2.0%, stream width ratio < 1.2 and stream profiling indicates it would be effective..

3.3.3 Cost Estimates

Cost estimates for structure replacement with bridges and embedded culverts were generated based on the channel width, slope of the culvert, depth of fill, road class and road surface type. Road details were sourced from FLNRORD (2020b) and FLNRORD (2020a) through *bcfishpass*. Interviews with Phil MacDonald, Engineering Specialist FLNR - Kootenay, Steve Page, Area Engineer - FLNR - Northern Engineering Group and Matt Hawkins - MoTi - Design Supervisor for Highway Design and Survey - Nelson were utilized to help refine estimates which have since been adjusted for inflation in 2020 and based on past experience.

Base costs for installation of bridges on forest service roads and permit roads with surfaces specified in provincial GIS road layers as rough and loose was estimated at \$30000/linear m and

3 Methods

assumed that the road could be closed during construction and a minimum bridge span of 15m. For streams with channel widths <2m, embedded culverts were reported as an effective solution with total installation costs estimated at \$100k/crossing (pers. comm. Phil MacDonald, Steve Page then adjusted for inflation in 2020). For larger streams (>6m), estimated span width increased proportionally to the size of the stream. For crossings with large amounts of fill (>3m), the replacement bridge span was increased by an additional 3m for each 1m of fill >3m to account for cutslopes to the stream at a 1.5:1 ratio. To account for road type, a multiplier table was generated to estimate incremental cost increases with costs estimated for structure replacement on paved surfaces, railways and arterial/highways costing up to 15 times more than forest service roads due to expenses associate with design/engineering requirements, traffic control and paving. The cost multiplier table (Table 3.5) should be considered very approximate with refinement recommended for future projects.

Table 3.5: Cost multiplier table based on road class and surface type.

Class	Surface	Class Multiplier	Surface Multiplier	Bridge \$/15m	Streambed Simulation \$
FSR	Rough	1	1	450,000	100,000
FSR	Loose	1	1	450,000	100,000
Resource	Loose	1	1	450,000	100,000
Collector	Loose	1	1	450,000	100,000
Recreation	Loose	1	1	450,000	100,000
Resource	Rough	1	1	450,000	100,000
Permit	Unknown	1	1	450,000	100,000
Permit	Loose	1	1	450,000	100,000
Permit	Rough	1	1	450,000	100,000
Unclassified	Loose	1	1	450,000	100,000
Unclassified	Rough	1	1	450,000	100,000
Unclassified	Paved	1	2	750,000	150,000
Unclassified	Unknown	1	2	750,000	150,000
Local	Loose	4	1	1,500,000	200,000
Local	Paved	4	2	3,000,000	400,000
Collector	Paved	4	2	3,000,000	400,000
Arterial	Paved	15	2	11,250,000	1,500,000
Highway	Paved	15	2	11,250,000	1,500,000
Rail	Rail	15	2	11,250,000	1,500,000

3.3 Fish Passage Assessments

3.3.4 Climate Change Risk Assessment

In collaboration with the Ministry of Transportation and Infrastructure (MoTi), a new climate change replacement program aims to prioritize vulnerable culverts for replacement (pers. comm Sean Wong, 2022) based on data collected and ranked related to three categories - culvert condition, vulnerability and priority. Within the “condition” risk category - data was collected and crossings were ranked based on erosion, embankment and blockage issues. The “climate” risk category included ranked assessments of the likelihood of both a flood event affecting the culvert as well as the consequence of a flood event affecting the culvert. Within the “priority” category the following factors were ranked - traffic volume, community access, cost, constructability, fish bearing status and environmental impacts (Table 3.6). This project is still in its early stages with methodology changes going forward.

Table 3.6: Climate change data collected at MoTi culvert sites

Parameter	Description
erosion_issues	Erosion (scale 1 low - 5 high)
embankment_fill_issues	Embankment fill issues 1 (low) 2 (medium) 3 (high)
blockage_issues	Blockage Issues 1 (0-30%) 2 (>30-75%) 3 (>75%)
condition_rank	Condition Rank = embankment + blockage + erosion
condition_notes	Describe details and rational for condition rankings
likelihood_flood_event_affecting_culvert	Likelihood Flood Event Affecting Culvert (scale 1 low - 5 high)
consequence_flood_event_affecting_culvert	Consequence Flood Event Affecting Culvert (scale 1 low - 5 high)
climate_change_flood_risk	Climate Change Flood Risk (likelihood x consequence) 1-6 (low) 6-12 (medium) 10-25 (high)
vulnerability_rank	Vulnerability Rank = Condition Rank + Climate Rank
climate_notes	Describe details and rational for climate risk rankings
traffic_volume	Traffic Volume 1 (low) 5 (medium) 10 (high)
community_access	Community Access - Scale - 1 (high - multiple road access) 5 (medium - some road access) 10 (low - one road access)
cost	Cost (scale: 1 high - 10 low)
constructability	Constructability (scale: 1 difficult -10 easy)
fish_bearing	Fish Bearing 10 (Yes) 0 (No) - see maps for fish points
environmental_impacts	Environmental Impacts (scale: 1 high -10 low)
priority_rank	Priority Rank = traffic volume + community access + cost + constructability + fish bearing + environmental impacts
overall_rank	Overall Rank = Vulnerability Rank + Priority Rank
priority_notes	Describe details and rational for priority rankings

3.4 Habitat Confirmation Assessments

Following fish passage assessments, habitat confirmations were completed in accordance with procedures outlined in the document “A Checklist for Fish Habitat Confirmation Prior to the Rehabilitation of a Stream Crossing” (Fish Passage Technical Working Group 2011). The main objective of the field surveys was to document upstream habitat quantity and quality and to determine if any other obstructions exist above or below the crossing. Habitat value was assessed based on channel morphology, flow characteristics (perennial, intermittent, ephemeral), the presence/absence of deep pools, un-embedded boulders, substrate, woody debris, undercut banks, aquatic vegetation and overhanging riparian vegetation. Criteria used to rank habitat value was based on guidelines in Fish Passage Technical Working Group (2011) (Table [3.2](#)).

During habitat confirmations, to standardize data collected and facilitate submission of the data to provincial databases, information was collected on digital field forms adapted from provincial [“Site Cards”](#). Habitat characteristics recorded included channel widths, wetted widths, residual pool depths, gradients, bankfull depths, stage, temperature, conductivity, pH, cover by type, substrate and channel morphology (among others). When possible, the crew surveyed downstream of the crossing to a minimum distance 300m and upstream to a minimum distance of 500 - 600m. Any potential obstacles to fish passage were inventoried with photos, physical descriptions and locations recorded on site cards. Surveyed routes were recorded with time-signatures on handheld GPS units.

3.4.1 Fish Sampling

3.4.1.1 Electrofishing

Fish sampling was conducted on a subset of sites when biological data was considered to add significant value to the physical habitat assessment information. Electrofishing was utilized for fish sampling according to stream inventory standards and procedures found in the Reconnaissance (1:20 000) Fish and Fish Habitat Inventory Manual (Resource Inventory Standards Committee 2001). A Haltech 2000 backpack electrofisher was used within discrete site units both upstream and downstream of the subject crossing with electrofisher settings and seconds, water quality parameters (i.e. conductivity, temperature and pH), start and end locations, length of site and wetted widths (average of a minimum of three) recorded.

3.4.1.2 Fish Handling and Processing

Captured fish were held in buckets with sufficient water to minimize stress until processing, and multiple buckets were used when catch numbers were high. For each fish captured, fork length, weight and species was recorded with results documented in the fish data submission spreadsheet.

3.4 Habitat Confirmation Assessments

3.4.1.3 Pit Tagging

Fish with a fork length greater than 60 mm and belonging to species approved under the scientific fish collection permit PG25-983916 were tagged with Passive Integrated Transponders (PIT tags) using the [Abdominal Cavity](#) method outlined by Biomark. To anesthetize fish prior to pit tagging, we used a solution of approximately 0.1 mL of clove oil per 1 L of water (1:10,000). This concentration was selected for its efficiency in providing effective sedation with minimal residual effects, making it ideal for studies in which fish are released back into their natural habitats (Fernandes et al. 2017). The clove oil solution was prepared in advance by dissolving pure clove oil in ethyl alcohol in a 1:9 ratio (clove oil: ethyl alcohol) to enhance solubility, then mixed into the water bucket (Fernandes et al. 2017). Fish were immersed in this solution until they reached an appropriate level of anesthesia for handling and then were tagged. To maintain needle sharpness and minimize injury risk, needles were replaced approximately every 10 fish. Each tagged fish was scanned with the PIT reader, and both the PIT tag ID and row ID were recorded. Once tagged, fish were placed into a bucket of fresh water and allowed to recover before being released back into the stream. Fish information and habitat data will be submitted to the province under scientific fish collection permit PG25-983916.

3.4.2 Environmental DNA (eDNA) Sampling

3.4.2.1 Sample Collection and Filtration

Samples were collected using two approaches informed by Hobbs et al. (2025): in-stream filtration with a Haltech EMOS sampler and grab-and-go bottle collection. For both methods, all water collected at a site (typically 1–3L) was filtered onto a single filter, yielding one replicate per site. Mixed cellulose ester (MCE) filters with 1 μ m pore size were used most commonly, with 0.45 μ m and 1.5 μ m filters of other materials used occasionally. Samples were typically collected upstream and downstream of each road–stream crossing, though in some cases only downstream samples were obtained.

For in-stream filtration, Haltech reusable aluminum housings were pre-loaded with filters. In high-turbidity conditions, a Haltech pre-filter with a 100 μ m nitex screen was installed upstream to remove coarse material. For grab-and-go sampling, stream water was collected in sanitized 1L HDPE Nalgene bottles and filtered at the end of each day.

After filtration, filters were removed from housings in a clean environment, placed in coin envelopes with silica desiccant inside labelled ziplock bags, and frozen until shipment. Aluminum housings and Nalgene bottles were sanitized with bleach solution between uses.

3.4.2.2 Quality Assurance

Approximately 10% of samples were collected for quality assurance. Field blanks using distilled water were collected daily. Office blanks were processed at field accommodations by filtering clean water when processing grab-and-go samples. Positive controls were obtained at select sites where electrofishing confirmed species presence prior to eDNA sampling.

3.4.2.3 Laboratory Analysis

Samples were shipped frozen to the University of Northern British Columbia (UNBC) for extraction and analysis. DNA was extracted and analyzed using quantitative polymerase chain reaction (qPCR) with species-specific assays following protocols developed by Murray and Booth (2023). Assays targeted mitochondrial genes (COI, CytB) using fluorescent dyes (FAM, VIC, HEX), allowing multiple species to be tested per sample through multiplexing. Target species included Chinook, Coho, Sockeye, Rainbow Trout, Bull Trout, Arctic Grayling, and Kokanee, with additional assays for Coastal Cutthroat and White Sturgeon at select sites.

Due to close evolutionary relationships and hybridization, some species pairs cannot be distinguished using mtDNA-based assays. The Bull Trout assay also detects Dolly Varden, requiring geographic context to interpret results in areas where ranges overlap. Similarly, Rainbow Trout and Coastal Cutthroat assays may cross-react where these species co-occur or hybridize.

To optimize laboratory efficiency, we collaborated with UNBC to develop a sample run plan that grouped samples by compatible dye types, minimizing the number of qPCR runs required. An assay mapping table cross-referencing species codes to assay identifiers and dye types was developed, along with processing scripts to generate sample groupings and a run order configuration specifying species order within each group.

3.4.2.4 eDNA Extraction

eDNA was extracted in the dedicated eDNA lab at UNBC and analysed with the protocols developed by Murray and Booth (2023).

3.4.3 Open Source Reporting Framework

This report is produced using open-source tools (R, bookdown) with full version control via git, enabling iterative updates. All code, data, and revision history are publicly accessible in the [project repository](#), with ongoing tasks tracked via GitHub [issues](#) and version history maintained in the [project changelog](#).

3.4.4 Aerial Imagery

Scripted processing and serving of UAV imagery collected during the project is available at https://github.com/NewGraphEnvironment/stac_uav_bc/ (Irvine [2025] 2025). [OpenDroneMap](#) was utilized to produce orthomosaics, digital surface models (DSMs), and digital terrain models (DTMs) (OpenDroneMap Authors [2014] 2025). To support efficient web-based access - imagery products were converted to cloud-optimized GeoTIFFs (COGs) using `rio-cogeo`, then collated according to the [SpatioTemporal Asset Catalog \(STAC\)](#) specification with `pystac` and uploaded to S3 storage Amazon Web Services (2025). A `titiler` tile server was set up to facilitate interactive viewing of the orthoimagery and an Application Program Interface (API) leveraging `stac-fastapi-pgstac` is served at <https://images.a11s.one> to enable linking of collection images through QGIS as well as remote spatial and temporal querying using open source software such as `rstac` (Development Seed [2019] 2025; `stac-utils` 2025; Simoes et al. 2021).

3.5 Engineering Design

3.5 Engineering Design

Engineering designs were signed and sealed by professional engineers. When possible - completed designs are loaded to the PSCIS data portal.

3.6 Remediations

Structure replacement was conducted by project specific contractors. If not already completed, as-built drawings will be loaded to the PSCIS data portal.

3.7 Monitoring

Monitoring of fish passage restoration sites — both proposed and completed - is essential to ensure restoration investments lead to meaningful ecological outcomes. Monitoring enables evaluation of whether remediation actions improve connectivity for fish and provides critical feedback to refine future prioritization and restoration strategies.

Baseline data collection, including fish sampling and aerial surveys via drone, are core components of this monitoring. While detailed methods for these activities are included in the previous habitat confirmation section, they are also fundamental to effectiveness monitoring, as they provide context to not only facilitate prioritization and communications but also for detecting change following restoration.

3.8 Collaborative GIS Environment

A collaborative GIS environment (sern_peace_fwcp_2023) was established using [QGIS](#) and [Mergin Maps](#) to enable project team members to view, edit, and analyze shared spatial data on desktop computers and on mobile devices in the field. The environment integrates background layers from the [BC Data Catalogue](#) and other sources with project-specific field data, providing a central platform for planning, standardized data collection, and progress tracking. A summary of layers loaded to the project is provided in [Appendix - Collaborative GIS Layers \(page 163\)](#).

To support consistent and targeted assessment, a custom field form was developed for routine effectiveness monitoring based loosely on Forest Investment Account (2003) but tailored specifically for fish passage projects. Table [3.7](#) outlines the monitoring metrics used.

Table 3.7: Description of monitoring metrics used for effectiveness monitoring.

Parameter	Description
Dewatering	Have the remediation works led to dewatering of the channel due to substrate aggradation or other factors?
Velocity	Are flow velocities similar to those within the natural channel? Are they expected to exceed swim speeds of particular fish species/life stages of interest?
Constriction	Have the remediation works led to constriction of the channel. Compare channel width underneath structure and within construction footprint to average channel widths upstream and downstream?

3 Methods

Parameter	Description
Substrate	Is the substrate within/under and adjacent to the remediated structure generally equivalent to that found upstream and downstream where natural channel conditions exist?
Riparian	What is the condition of the riparian area within the construction footprint?
UAV Flight	Was a flight conducted with unmanned aerial vehicle to document conditions at time of monitoring?
Flow_depth	What are the flow depths at the time of assessment within project footprint. Are depths expected to be sufficient to facilitate upstream passage for specific species/life stages of interest?
Stability	Does the structure appear to be stable or is there evidence of erosion/shifting?
Revegetation	How were riparian areas rehabilitated and are they improving fish habitat value?
Cover	Is cover available for fish within the construction footprint in the form of overhanging vegetation, large/small woody debris, boulders, undercut banks, etc?
Maintenance	If required, provide maintenance recommendations.
Recommendations	General recommendations for follow up. Could include revegetation, addition of substrate, fish sampling, etc.

4 Results and Discussion

4.1 Engage Partners

Project advancement in 2025 depended on engagement with partners across multiple agencies, Nations, industrial tenure holders, funders, and collaborators. Touchpoints during the year (email, video calls, meetings, field-based coordination, and contract administration) are summarised in Table 4.1.

Table 4.1: Project partners engaged in 2025.

Organization / Role	Topic / Focus
BC Timber Sales — Engineering Officer	Kennedy Siding (PSCIS 199663) FWCP-funded replacement; THUT15000 (PSCIS 203597) deactivation and riparian rehab
BC Timber Sales — Crossing CRP / bridges site plans	Kennedy Siding engineering design standards and site plan
McLeod Lake Indian Band — TLU Coordinator, Land Stewardship, LRO, Referrals, Director	NE-004 contract; land-guardian field participation; electrofishing course; collaborative QGIS/Mergin project access; council meeting planning; Azouzetta bull trout discussion
Sekani Forest Products (formerly Canfor)	Mackenzie and Peace NRD coordination; SUP spatial data; in-ground bridges KML
Canfor (prior-year coordination thread)	Fern Creek / Table FSR coordination
Sinclar Forest Group	Kerry Lake FSR; Arctic FSR; PSCIS 125000 (Trib to Parsnip) coordination
WLRS — Bull Trout WHA and Fisheries Sensitive Watershed program	Bull trout critical habitat; meeting follow-up and 2025 re-engagement
Independent fisheries biologists	eDNA methodology for bull trout and Arctic grayling; Arctic Grayling synthesis report collaboration
WLRS / Provincial Fish Passage Technical Working Group	Letters of support; FWCP F27 proposal; FPTWG and MoTi funding 2025/26 coordination
BC Hydro — FWCP grant administration	F25 close-out and F27 FWCP proposal submission; project ID corrections
Chu Cho Environmental	Upper Sustut Salmon Resiliency Fund proposal; Mesilinka River FWCP restoration planning; set up with collaborative QGIS/Mergin project access
Upper Peace Bull Trout CEMPRA Workshop (multi-agency: BC Hydro, WLRS, MoE BC, gov.ab.ca, Chu Cho, ESSA, independent consultants)	Bull trout stressor modeling and expert elicitation (March 2025)
UNBC	eDNA ddPCR analysis (program-scale 2025 submission)
UNBC NRESi and SERNbc	NRESi-FWCP Mixer event poster (Feb 2026); SERNbc coordination
Poisson Consulting	Stream Network Temperature Modelling 2025
Eclipse Geomatics / Skeena Knowledge Trust	Temperature data sharing
Stantec	Offsetting work consultation

4.2 Site Assessment Data

A summary of fish passage assessment procedures conducted through SERNbc since 2020 — including assessments, habitat confirmations, design, remediation, monitoring, fish sampling, eDNA sampling, and drone imagery — is provided in [Appendix - Assessment Data Summary \(page 51\)](#).

4.3 Climate Departure

Four findings from the regional climate-departure analysis carry to fish-passage prioritisation.

Warming is broad, fast, and significant region-wide. All five ecoregions of the FWCP Peace Region warmed about 1.8 °C since 1951 — annual mean (+1.8 °C), daytime maximum (+1.7 °C), and overnight minimum (+1.9 °C) all tell the same story. The trend is highly significant (Mann-Kendall $p < 0.001$ in every ecoregion). A subtle west-of-Rockies gradient layers on top, with the western and central ecoregions running about 0.2 °C warmer than the southeast — see [Appendix - Climate Departure \(page 55\)](#) Figure 5.11.

The atmosphere is drying despite locally rising precipitation. Vapour pressure deficit rose 0.28 hPa region-wide, significant in every ecoregion. Annual precipitation rose +4 % regionally but the long-term trend is significant only in the two northernmost ecoregions; soil moisture is essentially flat everywhere because the warmer atmosphere is drinking the extra water back.

The snowpack signal is about timing, not quantity. Total annual snowfall is essentially unchanged (-6 %); annual snow water equivalent is down 10 %. But the seasonal redistribution is sharp — summer snow water equivalent collapsed 75 %, spring snowmelt rose +37 %, and the snowmelt midpoint shifted 11 days earlier. The freshet is arriving weeks earlier and the high-elevation snowpack that historically lingered into summer no longer does.

Day-night asymmetry is present. Overnight minimums warmed 1.9 °C while daytime maximums warmed 1.7 °C — overnight outpacing daytime is the textbook fingerprint of greenhouse warming (Karl et al. 1993). Mean diurnal range has narrowed by 0.3 °C across the record.

For barrier prioritisation these signals cut both ways. Where upstream reaches are cold-limited, added growing-degree-days can raise the intrinsic potential of habitat above the barrier and *increase* the value of restoring passage; where upstream reaches sit closer to the upper edge of the cold-water thermal niche, the same warming reduces usable habitat and lowers the payoff. The watershed-group breakdown in [Appendix - Climate Departure \(page 55\)](#) maps the five-ecoregion signal onto the 16 FWCP Peace watershed groups.

4.4 Planning — Habitat, Connectivity and Floodplain Modelling

4.4.1 Habitat Modelling

Habitat modelling from `bcfishpass` including access model, linear spawning/rearing habitat model and lateral habitat connectivity models for watershed groups within our study area were updated for the spring of 2025 and are included spatially in the collaborative GIS project. A snapshot of these outputs related to each modeled and PSCIS stream crossing structure are also included within an `sqlite` database within this year's project reporting/code repository [here](#).

4.5 Fish Passage Assessments

4.4.1.1 Statistical Support for bc fishpass Fish Habitat Modelling Updates

Initial mapping of stream discharge and temperature causal effects pathways for the future purpose of focusing aquatic restoration actions in areas of highest potential for positive impacts on fisheries values (ie. elimination of areas from intrinsic models where water temperatures are likely too cold to support fish production) are detailed in Hill et al. (2024) which is included as [Attachment - Water Temperature Modelling \(page 0\)](#).

4.4.2 Floodplain Delineation

Modelled functional floodplain in the Parsnip River Watershed Group covers 48,116 ha — 8.6 % of the 5,597 km² watershed group — attached to 1,580 km of bull trout accessible order-3+ streams. Within that footprint sit 198 lakes and 464 mapped wetland polygons — the off-channel inventory that crossing remediations in these reaches stand to reconnect or leave severed, depending on whether design accounts for the floodplain context around the structure. The full metric table, watershed-scale map, and south-east detail map are in [Appendix - Floodplain Delineation \(page 79\)](#).

4.5 Fish Passage Assessments

Field assessments were conducted between September 21, 2025 and September 23, 2025 by Allan Irvine, R.P.Bio. and Lucy Schick, B.Sc. A total of 16 Fish Passage Assessments were completed, including 15 Phase 1 assessments and 1 reassessments.

Of the 16 sites where fish passage assessments were completed, 15 were not yet inventoried in the PSCIS system. This included 3 crossings considered “passable”, 1 crossings considered a “potential” barrier, and 11 crossings were considered “barriers” according to threshold values based on culvert embedment, outlet drop, slope, diameter (relative to channel size) and length (MoE 2011).

A summary of crossings assessed, a rough cost estimate for remediation, and a priority ranking for follow-up for Phase 1 sites is presented in Table [4.2](#). Detailed data with photos are presented in [Attachment - Phase 1 Data and Photos \(page 175\)](#).

4 Results and Discussion

Table 4.2: Upstream habitat estimates and cost benefit analysis for Phase 1 assessments ranked as a ‘barrier’ or ‘potential’ barrier. Bull trout network model used for habitat estimates (total length of stream network <25% gradient).

PSCIS ID	External ID	Priority	Stream	Road	Barrier Result	Habitat value	Habitat Upstream (km)	Stream Width (m)	Fix	Cost Est (\$K)
6559	–	low	Tributary To Williston Reservoir	Finlay Community Connector	Barrier	Low	17.4	1.0	SS-CBS	100
203597	15201563	high	Tributary To Nation River	Thut 15000	Barrier	High	16.8	2.5	OBS	450
203598	15201146	moderate	Tributary To Nation River	Thut 15000	Barrier	Medium	9.2	1.6	SS-CBS	100
203599	15202249	moderate	Tributary To Nation River	Thut 14000	Barrier	Medium	14.2	3.0	OBS	450
203600	15200985	moderate	Tributary To Nation River	Finlay-Nation FSR	Barrier	High	30.6	4.1	OBS	450
203601	15201007	moderate	Tributary To Nation River	Finlay-Nation FSR	Barrier	High	15.9	3.5	OBS	450
203602	15201006	high	Tributary To Nation River	Finlay-Nation FSR	Barrier	High	17.5	3.4	OBS	495
203603	16400714	moderate	Tributary To Williston Reservoir	Finlay-Nation FSR	Barrier	High	106.7	7.4	OBS	450
203604	16401990	moderate	Tributary To Williston Reservoir	Finlay FSR	Barrier	Medium	26.6	10.0	OBS	810

4.6 Habitat Confirmation Assessments

During the 2025 field assessments, habitat confirmation assessments were conducted at 3 sites within the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toodoggone River, and Upper Omineca River watershed groups. A total of approximately 2km of stream was assessed. Georeferenced field maps are provided in [Attachment - Maps \(page 0\)](#).

As collaborative decision-making was ongoing at the time of reporting, site prioritization can be considered preliminary. Results are summarized in Table 4.3 with raw habitat and fish sampling data included in [Attachment - Data \(page 177\)](#). A summary of preliminary modelling results illustrating quantities of bull trout spawning and rearing habitat potentially available upstream of each crossing as estimated by measured/modelled channel width and upstream accessible stream length are presented in Figure 4.1. Detailed information for each site assessed with Phase 2 assessments (including maps) are presented within site-specific appendices to this document.

4.6 Habitat Confirmation Assessments

Table 4.3: Overview of habitat confirmation sites. Bull trout rearing model used for habitat estimates (total length of stream network <10.5% gradient).

PSCIS ID	Stream	Road	Tenure	UTM	UTM zone	Fish Species	Habitat Gain (km)	Habitat Value	Priority	Comments
199663	Tributary To Parsnip River	Chco 12000	BCTS	515416 6094963	10	RB	0.0	Medium	Moderate	Low gradient stream with significant riparian tree blow down throughout. Very difficult walking so most was done outside of the active channel with spot visits to the stream for measurements/photos completed intermittently for the 600m that was assessed. Wide floodplain within well defined canyon (approximately 15m deep with slopes of ~45°). Healthy riparian, consisting of herb and shrub cover within sparse spruce forest that appears to have been impacted by some sort of beetle or other affliction. Most of the spruce trees were dead with many already fallen down.
203597	Tributary To Nation River	Thut 15000	Sekani R07214 8	425959 6122703	10	RB	5.6	High	High	Surveyed 350m upstream to the beginning of the wetlands. Very nice stream with good flow, although recent rainfall may have had an influence. Pockets of gravels throughout the area surveyed. Although the area viewed was limited, the wetlands appeared shallow so freezing to the bottom may occur during the winter.
203605	Tributary To Williston Reservoir	Finlay FSR	MoF	486504 6124990	10	RB	35.9	High	High	Large stream with complex habitat and diverse cover including deep pools, boulders, large and small woody debris, and undercut banks. A canyon section approximately 100m long was located 100m upstream of the Modeste FSR bridge, containing multiple cascade steps up to 1m high with deep downstream pools. At the upper end of the site, the channel deflected against steep gully walls. Abundant gravel was present, suitable for adfluvial and resident rainbow trout as well as bull trout. eDNA sample collected.

4 Results and Discussion

Table 4.4: Summary of Phase 2 fish passage reassessments.

PSCIS ID	Embedded	Outlet Drop (m)	Diameter (m)	SWR	Slope (%)	Length (m)	Final score	Barrier Result
PSCIS ID	Embedded	Outlet Drop (m)	Diameter (m)	SWR	Slope (%)	Length (m)	Final score	Barrier Result
199663	No	0.60	1.2	0.00	3	13	30	Barrier
203597	No	0.25	0.8	3.12	1	19	29	Barrier
203605	No	0.95	3.9	1.54	2	46	37	Barrier

4.6 Habitat Confirmation Assessments

Table 4.5: Cost benefit analysis for Phase 2 assessments. Bull trout rearing model used for habitat estimates (total length of stream network <10.5% gradient).

PSCIS ID	Stream	Road	Barrier Result	Habitat value	Stream Width (m)	Fix	Cost Est (\$K)	Habitat Upstream (m)	Cost Benefit (m / \$K)	Cost Benefit (m2 / \$K)
199663	Tributary To Parsnip River	Chco 12000	Barrier	Medium	3.0	SS-CBS	–	0	–	–
203597	Tributary To Nation River	Thut 15000	Barrier	High	2.5	OBS	450	5570	12377.8	15472.2
203605	Tributary To Williston Reservoir	Finlay FSR	Barrier	High	6.7	OBS	450	35932	79848.9	239546.7

4 Results and Discussion

Table 4.6: Summary of Phase 2 habitat confirmation details.

PSCIS ID	Length surveyed upstream (m)	Average Channel Width (m)	Average Wetted Width (m)	Average Pool Depth (m)	Average Gradient (%)	Total Cover	Habitat Value
-	-	-	-	-	-	-	-
---	-----	-----	-----	-----	-----	-----	-----

4.6 Habitat Confirmation Assessments

Table 4.7: Summary of watershed area statistics upstream of Phase 2 crossings.

Site	Area Km	Elev Site	Elev Min	Elev Max	Elev Median	Elev P60	Aspect
199663	1.0	713	707	849	784	774	SW
203597	8.2	900	907	1065	980	971	S
203605	55.6	681	671	1644	980	937	SE
203611	55.6	682	671	1644	980	937	SE

* Elev P60 = Elevation at which 60% of the watershed area is above




Figure 4.1: Summary of potential rearing and spawning habitat upstream of habitat confirmation assessment sites. Bull trout rearing and spawning models used for habitat estimates (total length of stream network <10.5% and <5.5% gradient, respectively).

4.6.1 Fish Sampling

Fish sampling was conducted at 8 sites within 6 streams, with a total of 85 fish captured. Fork length, weight, and species were documented for each fish. Salmonids with fork lengths >60mm were PIT-tagged to facilitate long-term tracking of health and movement. Fork length data was used to delineate salmonids based on life stages: fry (0 to 65mm), parr (>65 to 110mm), juvenile (>110mm to 140mm), and adult (>140mm) by visually assessing the histograms presented in Figure 4.3. A summary of sites assessed is included in Table 4.8, and raw data is provided in [Attachment - Data \(page 177\)](#). A summary of density results is also presented in Figure 4.2.

Table 4.8: Summary of electrofishing sites.

site	stream	passes	ef_length_m	ef_width_m	area_m2	enclosure
125179_us_ef1	Tributary To Missinka River	1	45	2.2	99.0	partial enclosure
125179_ds_ef1	Tributary To Missinka River	1	35	2.7	94.5	partial enclosure
125231_ds_ef1	Tributary To Table River	1	50	2.8	140.0	partial enclosure
125231_us_ef1	Tributary To Table River	1	50	1.6	80.0	partial enclosure
198692_us_ef1	Tributary To Kerry Lake	1	16	–	–	partial enclosure
198692_us_ef2	Tributary To Kerry Lake	1	12	–	–	open
198692_ds_ef1	Tributary To Kerry Lake	1	34	–	–	partial enclosure
198692_ds_ef2	Tributary To Kerry Lake	1	4	–	–	open

4.6 Habitat Confirmation Assessments




Figure 4.2: Boxplots of densities (fish/100m²) of fish captured downstream (ds) and upstream (us) by electrofishing.

4 Results and Discussion




Figure 4.3: Histograms of fish lengths by species. Fish captured by electrofishing during habitat confirmation assessments.

4.6.2 Environmental DNA (eDNA) sampling

New in 2025, environmental DNA (eDNA) sampling was incorporated into the program at both previously assessed and newly added sites. A total of 35 collection events were undertaken across 10 streams, including 28 real environmental samples, 3 field blanks, and 4 office blanks for quality assurance.

Per-species detection results across the 28 real environmental samples are summarized in Table [4.9](#). Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; sub-threshold signals (1-3 droplets) are tracked separately as they are not interpretable as species presence per ddPCR convention. Site-level detail (per-site detection table, field blanks, and retests) is provided in [Appendix - Environmental DNA Results \(page 145\)](#), and the [interactive Peace eDNA map](#) shows toggleable per-species layers.

4.7 Engineering Design

Table 4.9: Per-species summary of eDNA results across real environmental samples in the Peace project area (excludes field and office blanks).

Species	Code	Sites tested	Detected (≥4 droplets)	Sub-threshold only (1-3 droplets)	No detection (0 droplets)
Species	Code	Sites tested	Detected (≥4 droplets)	Sub-threshold only (1-3 droplets)	No detection (0 droplets)
Rainbow Trout	RAIN	28	22	2	4
Bull Trout	BULT	28	1	8	19
Burbot	BURB	2	1	0	1
Arctic Grayling	GRAY	24	0	1	23
Kokanee	SOCK	22	0	0	22

4.7 Engineering Design

MIGHT NEED TO UPDATE

No new designs were commissioned in due to uncertainty related to forest harvesting activities and lack of funding for the 50% costs of replacing structures. All sites with a design for remediation of fish passage can be found online within the Results and Discussion section of the report found [here](#).

4.8 Monitoring

In 2025, effectiveness monitoring was conducted at two previously remediated crossings, and baseline data collection (fish sampling and eDNA only — no effectiveness form) was completed at one site ahead of planned remediation:

- **Tributary to Missinka River (PSCIS 125179)** on Chuchinka-Missinka FSR — effectiveness monitoring (form + fish sampling + eDNA), extending the multi-year baseline established in Irvine and Winterscheidt (2023) and Irvine and Schick (2025). Detail in [Appendix - Tributary to Missinka River \(page 149\)](#).
- **Tributary to the Table River (PSCIS 125231)** on Chuchinka-Table FSR — effectiveness monitoring (form + fish sampling + eDNA) following the 2023 replacement of the crossing with a clear-span bridge by Canfor with FWCP funding via SERNbc coordination (Irvine and Winterscheidt 2024). Detail in [Appendix - Tributary to the Table River \(page 153\)](#).
- **Tributary to Kerry Lake (PSCIS 198692)** — pre-remediation baseline (fish sampling and eDNA) to inform future effectiveness monitoring once the crossing is upgraded. The 2024 habitat confirmation is documented in Irvine and Schick (2025). Detail in [Appendix - Tributary to Kerry Lake \(page 157\)](#).

A summary of effectiveness monitoring form metrics across the two effectiveness sites is presented in Table [4.10](#).

Table 4.10: Summary of monitoring sites.

PSCIS ID	Stream	Road	Crossing type	Crossing subtype	Diameter or span (m)	Length or width (m)	Assessment comments
125231	Tributary to Table River	Table FSR	OBS	BRIDGE	–	–	Overall, the site looks very similar to last year, in good condition with no evidence of any structural issues and the instream portion doing quite well. As there is a large amount of rock rip wrap adjacent to the stream that was not planted the riparian vegetation within the footprint is basically nonexistent. Electrofishing was conducted at sites downstream and upstream of the bridge with fish over 65mm PIT tagged. eDNA samples were collected as well.
125179	Tributary to Missinka River	Chuchinka-Missinka FSR	OBS	BRIDGE	16	4.8	Routine effectiveness evaluation. Structure was replaced several years ago and monitoring is ongoing. Electrofishing with fish over 65mm in length PIT tagged, as well as EDNA sampling was conducted in 2025.

4.9 Collaborative GIS Environment

In addition to numerous layers documenting fieldwork activities since , a summary of background information spatial layers and tables loaded to the collaborative GIS project (sern_peace_fwcp_2023) at the time of writing (2026-05-14) is provided in [Appendix - Collaborative GIS Layers \(page 163\)](#).

5 Recommendations

A major challenge in advancing fish passage restoration is the complexity of working across jurisdictions and with multiple stakeholders—rail and highway authorities, forestry ministries, licensees, and private landowners. These partners are often being asked to accommodate priorities that originate outside their mandates and budgets. Convincing them to invest in difficult, high-cost interventions—like modifying crossings or relocating infrastructure—requires navigating uncertainty about costs and ecological outcomes, as well as a disconnect between the benefits to watershed health and the internal pressures or performance goals of these agencies. It's a tough ask: to take on massive, uncertain projects when they're already stretched thin with their own responsibilities.

Fish passage restoration across British Columbia is further complicated by the legacy of infrastructure deeply embedded in the landscape. Roads, railways, highways, community infrastructure and private assets often constrain floodplains and disrupt natural hydrological processes. While targeted repairs to individual barriers are essential, they won't resolve the broader systemic issues without rethinking and restructuring how infrastructure interacts with watershed function. Loss of riparian vegetation and intensive beaver management only add to the degradation. Addressing these challenges means making strategic, well-communicated choices—picking battles carefully, building trust, and staying committed to a longer-term transformation.

While preliminary top remediation priorities are provided by watershed group, these rankings are inherently subjective and can depend on the capacity and willingness of infrastructure owners and tenure holders to support implementation—both financially and over the often multi-year project timelines. In practice, we must often act opportunistically, pursuing simpler, lower-cost options to maintain momentum and achieve near-term progress.

Government, community groups, landowners, non-profits, industry and other stakeholders should work collaboratively to address high and moderate priority barriers identified online within the Results and Discussion section of the report found [here](#). Although the table presents many options - currently - linked reports specify whether each site is a low, moderate, or high priority. Progress on any front is meaningful, and aiming to remediate at least one high-priority site per year per watershed group—regardless of its overall rank—is a practical and effective approach.

To enhance fish passage restoration in the FWCP Peace Region:

- Maintain strong partnerships to support funding, site selection, remediation, and monitoring through adaptive management informed by traditional knowledge and real-time data.
- Prioritize detailed assessments in areas with blockages and high habitat potential, especially near McLeod Lake.
- Use climate modeling to prioritize crossings that enable access to cold, drought-resistant habitats.
- Secure financial commitments for Fern Creek remediation despite uncertainties in harvest planning.

5 Recommendations

- Continue effectiveness monitoring at key sites using fish sampling, eDNA, PIT tagging, temperature data, and aerial imagery.
- Continue to develop a cost-effective monitoring framework to assess productivity gains from improved passage.
- Collaborate with WLRS, UNC, local fisheries experts, FWCP, and the CEMPRA Project working group.
- Utilize environmental DNA (eDNA) to better understand bull trout and Arctic grayling habitat use at both potential and remediated sites. This will refine prioritization and assess fish passage effectiveness.

Appendix - Fish Species by Watershed Group

Historical fish observations by watershed group are recorded in the BC Fisheries Information Summary System (FISS) (MoE 2024a). Per-species presence across the watershed groups in this study area is summarised in Table 5.1. In the interactive (gitbook) version of this report the table shows one column per watershed group; in the PDF version the watershed-group columns are collapsed to a single comma-separated list of watershed-group codes per species to fit page width.

Table 5.1: Fish species recorded in the Carp Lake, Crooked River, Finlay Arm, Finlay River, Firesteel River, Fox River, Ingenika River, Lower Omineca River, Mesilinka River, Nation River, Ospika River, Parsnip Arm, Parsnip River, Peace Arm, Toadoggone River, and Upper Omineca River watershed groups.

Scientific Name	Species Name	BC List	COSEWIC	Present in WSGs
<i>Catostomus catostomus</i>	Longnose Sucker	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Catostomus columbianus</i>	Bridgelip Sucker	Yellow	–	Crooked
<i>Catostomus commersonii</i>	White Sucker	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Fox, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Upper Omineca
<i>Catostomus macrocheilus</i>	Largescale Sucker	Yellow	–	Carp Lake, Crooked, Finlay Arm, Ingenika, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toadoggone
<i>Coregonus clupeaformis</i>	Lake Whitefish	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Cottus aleuticus</i>	Coastrange Sculpin (formerly Aleutian Sculpin)	Yellow	–	Parsnip
<i>Cottus asper</i>	Prickly Sculpin	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Cottus cognatus</i>	Slimy Sculpin	Yellow	–	Carp Lake, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Cottus ricei</i>	Spoonhead Sculpin	Yellow	NAR (May 1989)	Finlay Arm, Lower Omineca, Parsnip Arm, Upper Omineca
<i>Couesius plumbeus</i>	Lake Chub	Yellow	DD	Carp Lake, Crooked, Finlay Arm, Finlay, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Esox lucius</i>	Northern Pike	Yellow	–	Crooked
<i>Hybognathus hankinsoni</i>	Brassy Minnow	No Status	–	Carp Lake, Crooked, Finlay Arm, Nation, Parsnip Arm

Appendix - Fish Species by Watershe...

Scientific Name	Species Name	BC List	COSEWIC	Present in WSGs
<i>Lota lota</i>	Burbot	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Mylocheilus caurinus</i>	Peamouth Chub	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Ingenika, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Upper Omineca
<i>Oncorhynchus clarkii</i>	Cutthroat Trout	No Status	–	Ospika
<i>Oncorhynchus clarkii clarkii</i>	Coastal Cutthroat Trout	Blue	–	Finlay, Lower Omineca
<i>Oncorhynchus kisutch</i>	Coho Salmon	Not Reviewed	–	Mesilinka
<i>Oncorhynchus mykiss</i>	Rainbow Trout	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Oncorhynchus mykiss</i>	Steelhead	Yellow	–	Mesilinka
<i>Oncorhynchus mykiss</i>	Steelhead (Winter-run)	Yellow	–	Finlay Arm
<i>Oncorhynchus nerka</i>	Kokanee	Not Reviewed	–	Finlay Arm, Finlay, Firesteel, Ingenika, Lower Omineca, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone
<i>Oncorhynchus nerka</i>	Sockeye Salmon	Not Reviewed	–	Mesilinka
<i>Osmerus dentex</i>	Rainbow Smelt	Unknown	–	Firesteel, Nation, Parsnip
<i>Prosopium coulterii</i>	Pygmy Whitefish	Yellow	NAR (Nov 2016)	Finlay Arm, Fox, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Upper Omineca
<i>Prosopium cylindraceum</i>	Round Whitefish	Yellow	–	Parsnip
<i>Prosopium williamsoni</i>	Mountain Whitefish	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Ptychocheilus oregonensis</i>	Northern Pikeminnow	Yellow	–	Carp Lake, Crooked, Finlay Arm, Lower Omineca, Nation, Parsnip Arm, Parsnip, Peace Arm
<i>Rhinichthys cataractae</i>	Longnose Dace	Yellow	–	Crooked, Finlay Arm, Finlay, Firesteel, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Toadoggone, Upper Omineca
<i>Richardsonius balteatus</i>	Redside Shiner	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Fox, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Upper Omineca
<i>Salvelinus confluentus</i>	Bull Trout	Blue	SC (Nov 2012)	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca
<i>Salvelinus fontinalis</i>	Brook Trout	Exotic	–	Crooked, Parsnip Arm, Parsnip
<i>Salvelinus malma</i>	Dolly Varden	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toadoggone, Upper Omineca

Scientific Name	Species Name	BC List	COSEWIC	Present in WSGs
Salvelinus namaycush	Lake Trout	Yellow	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toodoggone, Upper Omineca
Thymallus arcticus	Arctic Grayling	Yellow	–	Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toodoggone, Upper Omineca
–	Arctic Char	–	–	Finlay
–	Chub (General)	–	–	Carp Lake, Crooked, Parsnip Arm, Parsnip
–	Dace (General)	–	–	Finlay Arm, Parsnip Arm, Parsnip
–	Lamprey (General)	–	–	Nation
–	Minnow (General)	–	–	Carp Lake, Crooked, Finlay Arm, Fox, Parsnip Arm, Parsnip, Upper Omineca
–	Mottled Sculpin	–	–	Firesteel, Nation, Parsnip
–	Salmon (General)	–	–	Nation
–	Sculpin (General)	–	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Fox, Ingenika, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm, Toodoggone
–	Smelt (General)	–	–	Finlay Arm, Parsnip Arm
–	Squanga	–	–	Crooked
–	Sucker (General)	–	–	Carp Lake, Crooked, Finlay Arm, Finlay, Firesteel, Lower Omineca, Mesilinka, Nation, Parsnip Arm, Parsnip, Peace Arm, Toodoggone
–	Whitefish (General)	–	–	Carp Lake, Crooked, Lower Omineca, Mesilinka, Nation, Ospika, Parsnip Arm, Parsnip, Peace Arm
* COSEWIC abbreviations : SC - Special concern DD - Data deficient NAR - Not at risk E - Endangered T - Threatened BC List definitions : Yellow - Species that is apparently secure Blue - Species that is of special concern Exotic - Species that have been moved beyond their natural range as a result of human activity Dolly Varden are documented in all three study watersheds — Carp, Crooked, and Parsnip — but these records are likely misidentifications of bull trout (COSEWIC 2012).				

Appendix - Assessment Data Summary

Fish passage assessment procedures conducted through SERNbc since 2020 are amalgamated in Tables [5.2](#) - [5.3](#). Since 2020, orthoimagery and elevation model rasters have been generated and stored as Cloud Optimized Geotiffs on a cloud service provider (AWS) with select imagery linked to in the collaborative GIS project. Additionally, a tile service has been set up to facilitate viewing and downloading of individual images, provided in [Appendix - UAV Imagery \(page 161\)](#).

Table 5.2:
Summary
of fish
passage
assessment
procedures
conducted
in northern

British
Columbia
through
SERNbc.

**NOTE: To
view all
columns in
the table -
please
click on
one of the
sort
arrows
within
column
headers
before
scrolling
to the
right.**

Show entries

Watershed Group	Assessment	Reassessment	Habitat Confirmation	Design	Rem
All	All	All	All	All	All

Appendix - Assessment Data Summary

Francois Lake	1	0	4	0	0
Lower Chilako River	1	0	1	0	0
Morkill River	1	0	5	0	0
Nechako River	2	0	4	0	0
Tabor River	0	1	0	0	0
Upper Fraser River	2	2	3	0	0
Total	7	3	17	0	0

Showing 1 to 7 of 7 entries

Excel

CSV

Previous

1

Next

Table 5.3:
Details of
fish
passage
assessment
procedures
conducted
in northern
British
Columbia
through
SERNbc.

Show entries

Stream Crossing Id	My Crossing Reference	Watershed Group	Stream Name	Road Name	Assessment
All	All	All	All	All	All

4931		Upper Fraser River	Crooked Creek	Tinsley Pit Road		yes
7620		Upper Fraser River	Tributary of McLennan River	CN Rail access road		yes
7622		Francois Lake	UNN Flows into Fraser Lake	Stella Rd		
196200		Tabor River	Bittner Creek	Forman Road		yes
199171		Francois Lake	Tributary to Fraser Lake	Gala Bay Road		
199172		Francois Lake	Scotch Creek	Stella Road		
199173		Nechako River	Tributary to Nechako River	Dog Creek Road		
199174		Nechako River	Tributary to Nechako River	Sutherland FSR		
199190		Nechako River	Clear Creek	Highway 27 S		
199204		Nechako River	Nine Mile Creek	Settlement Rd		

Showing 1 to 10 of 26 entries

Excel

CSV

Previous

1

2

3

Next

Appendix - Climate Departure

Climate departure and fish passage

Stream crossings are the visible structures in a fish-passage inventory: culverts, bridges, weirs, fords. The streams they cross sit inside a climate that is changing — warmer air, shifting snowpack, longer shoulder seasons — and those climate changes set the hydrologic and thermal context the structures pass fish *through*. A culvert that worked when it was installed in 1985 is now passing a different hydrograph: earlier freshet, lower late-summer flows, warmer water. The barrier inventory does not see that on its own.

Warmer air drives warmer streams. For the cold-water salmonids the Peace supports — bull trout, Arctic grayling, mountain whitefish — that can move habitat in either direction: cold-limited headwaters gain growing-degree-days (larger fish, better overwinter survival, stronger out-migration), while reaches near the upper edge of the thermal niche lose usable habitat. The recovery value of removing a downstream barrier scales with which side of that line each upstream reach sits on. Climate departure tells you which side that is, watershed by watershed.

Beyond temperature, hydrologic timing matters. Earlier and faster snowmelt shifts the freshet weeks earlier in the calendar and compresses the high-flow pulse. That changes design assumptions for crossing structures, and it changes the timing of migration, gravel mobilisation, and off-channel rearing — the in-stream processes barrier removal is meant to restore.

Quantifying departure at the watershed scale lets us carry climate context into barrier prioritisation alongside the structural and biological inputs.

Area of interest

The FWCP Peace Region is a 72,811 km² administrative boundary in northeastern British Columbia covering Williston Reservoir and surrounding watersheds. All watersheds drain to the Mackenzie River — the Peace flows east into the Slave, which flows into the Mackenzie, which empties into the Beaufort Sea on the Arctic Ocean. The region spans 5 ecoregions (Figure [5.1](#)) — broad climate-physiography zones based on latitude, elevation, and dominant vegetation — and we carry that ecoregion partition through the analysis because climate departure can vary across them in ways the regional average hides.




Figure 5.1: FWCP Peace Region area of interest in northeastern British Columbia, coloured by ecoregion. Williston Reservoir dominates the basin; the climate-departure pipeline is run on the regional outline (heavy blue) and again on each ecoregion polygon.

Recent decade versus pre-warming reference

The table below compares two windows directly — the recent decade (2015–2025) against the pre-warming reference (1951–1980). The difference is reported in the variable's native units, alongside percent change where it is meaningful (precipitation, soil moisture, vapour pressure deficit, relative humidity). Two p-values appear, answering different questions. **Δp (windows)** is the Welch t-test on the annual values within each window — this tells us if the recent decade differs from the pre-warming reference. **Trend p (75-yr)** is the Mann-Kendall test on the full 1951–present series — this tells us if there is a steady year-on-year ramp. Step changes show up as significant Δp with non-significant trend p; gradual ramps show up as both significant.

Trends over the long record

Table 5.4: Recent decade (2015-2025) versus pre-warming reference (1951-1980) for all 15 climate variables and all available periods, FWCP Peace Region. Two p-values answer different questions — see text.

Variable	Period	Recent (2015– 2025)	Pre- warming (1951– 1980)	Δ		Δ p (windows)	Trend p (75- yr)
				absolute	Δ %		
prcp	annual	835.20	803.11	32.08	4.0	0.308	0.197
prcp	fall	239.18	230.76	8.42	3.7	0.622	0.493
prcp	spring	155.72	150.62	5.10	3.4	0.643	0.164
prcp	summer	266.04	242.15	23.89	9.9	0.283	0.185
prcp	winter	174.26	179.59	-5.33	-3.0	0.674	0.661
rh	annual	74.08	74.08	0.00	0.0	0.992	0.862
rh	fall	79.38	80.16	-0.78	-1.0	0.279	0.091
rh	spring	68.43	69.49	-1.06	-1.5	0.159	0.385
rh	summer	67.25	67.63	-0.38	-0.6	0.760	0.756

The recent decade ran 1.8 °C warmer than the pre-warming reference for annual mean temperature, with both window and trend p-values below 0.001. Vapour pressure deficit rose significantly on both tests. Annual precipitation rose about 4 % but neither test confirms the shift. Soil moisture and relative humidity show no meaningful change. The snow rows (swe summer, snowmelt spring, snowmelt_doy_50) carry the headline snowpack-timing signals discussed below.

Trends over the long record

Anomalies are computed against the 1951–1980 pre-warming reference. Saying a year is “+1.5 °C” means it was 1.5 °C warmer than the average year between 1951 and 1980. The trend table below has two rows per variable — trends computed from two different start years:

- **1951–present (75 years)** — the long view. Captures the full magnitude of warming since the pre-warming reference.
- **1981–present (45 years)** — starts at the beginning of the current 30-year climate normal (1981–2010), the reference used in most published climate products.

Comparing the two slopes is informative. If the 45-year slope is steeper than the 75-year slope, warming has accelerated; if shallower, warming has slowed (though “slower” almost never means “stopped”). Similar slopes mean a roughly steady rate of change across the full record. Total change is slope multiplied by the number of years — the cumulative shift over the trend window.

Table 5.5: Trend statistics for all variables and periods, FWCP Peace Region — Theil-Sen slope, Mann-Kendall p-value, and cumulative change over each trend window.

Parameter	Period	Slope	Years	Total Change	Unit	p-value
Precipitation	Annual	0.085	75	6.4	%	0.1971
Relative humidity	Annual	0.001	75	0.1	%	0.8620
Snow cover	Annual	-0.053	75	-4.0	%	0.0026
Snowfall	Annual	-0.088	75	-6.6	%	0.2528
Snowfall fraction	Annual	-0.079	75	-5.9	%	0.0078
Snowmelt	Annual	-0.039	75	-2.9	%	0.5279
Day of 50% melt	Annual	-0.150	75	-11.3	day	0.0018
Peak weekly melt rate	Annual	0.101	75	7.6	mm/wk	0.3848




Figure 5.2: Annual mean temperature anomaly for the FWCP Peace Region relative to the 1951-1980 baseline. Bars are degrees above (red) or below (blue) the pre-warming average; the two trend lines are the 75-year (1951-present, dashed) and 45-year (1981-present, solid) Theil-Sen slopes.




Figure 5.3: Annual precipitation anomaly (% of the 1951-1980 baseline) for the FWCP Peace Region. The long-term trend is positive but not statistically significant at the regional scale; significance does appear in the two northernmost ecoregions (see Per-Ecoregion section).

Daytime highs and overnight lows

The `cd` package ships daytime maximum (`tmax`) and overnight minimum (`tmin`) temperatures alongside the daily mean. They carry distinct information. Overnight minimums warming faster than daytime maximums — the “day-night asymmetry” — is one of the textbook fingerprints of greenhouse warming. Karl et al. (1993) documented this first at the global scale, finding overnight minimums rose at roughly three times the rate of daytime maximums between 1951 and 1990.

For the FWCP Peace Region, overnight minimums are warming faster than daytime maximums. Daytime maximums warmed about $+0.027\text{ }^{\circ}\text{C}$ per year since 1951 ($+2.0\text{ }^{\circ}\text{C}$ cumulative); overnight minimums warmed about $+0.032\text{ }^{\circ}\text{C}$ per year ($+2.4\text{ }^{\circ}\text{C}$ cumulative). The overnight side warmed roughly $0.4\text{ }^{\circ}\text{C}$ more than the daytime side over the full record, narrowing the diurnal temperature range by the same amount (Figure [5.6](#)).




Figure 5.4: Annual daytime maximum temperature (tmax) anomaly relative to the 1951-1980 baseline.




Figure 5.5: Annual overnight minimum temperature (tmin) anomaly relative to the 1951-1980 baseline.




Figure 5.6: Diurnal temperature range (daytime maximum minus overnight minimum) annual mean for the FWCP Peace Region. The downward slope is the day-night asymmetry — overnight lows are warming faster than daytime highs.

Snowpack

Snowpack drives the seasonal water supply across British Columbia: winter precipitation falls as snow, accumulates on the ground, and releases as meltwater across spring and summer. That seasonal storage is the difference between a late-summer creek that still flows and one that does not. It also sets the timing of the spring freshet — the annual flood pulse that salmonids time their up-river migrations to.

For the FWCP Peace Region the snowpack signal is unambiguous and concentrated in the warm shoulders of the year. Annual snow water equivalent declined about 10 % (135 → 122 mm), but the seasonal breakdown is sharper: **summer snow water equivalent collapsed by 75 %** (21.5 → 5.3 mm) and **spring snowmelt rose 37 %** (212 → 290 mm) as the freshet shifted earlier in the calendar. Total annual snowfall barely changed (-6 %) — the snow water equivalent decline is mostly about warming removing snow earlier, not less snow falling.

Seasonal snowpack curve

Aggregated to the four standard meteorological seasons — winter (December–February), spring (March–May), summer (June–August), and fall (September–November) — the table below

compares the recent decade (2015–2025) directly against the pre-warming reference (1951–1980) for each of the four monthly-native snow variables. The headline numbers (summer snow water equivalent collapse, spring snowmelt rise) are in the **summer** and **spring** rows.

Table 5.6: Seasonal snowpack: recent decade versus pre-warming reference, FWCP Peace Region. Summer snow water equivalent collapse (-75 %) and spring snowmelt rise (+37 %) carry the headline signals.

Variable	Season	Pre-warming (1951–1980)	Recent (2015–2025)	Δ %
Snow water equivalent (mm)	winter	195.9	192.6	-1.7
Snow water equivalent (mm)	spring	292.8	259.6	-11.4
Snow water equivalent (mm)	summer	21.5	5.3	-75.5
Snow water equivalent (mm)	fall	30.2	29.1	-3.6
Snowfall (mm)	winter	176.3	170.0	-3.6
Snowfall (mm)	spring	109.0	94.1	-13.7
Snowfall (mm)	summer	12.0	4.9	-59.5
Snowfall (mm)	fall	135.0	136.6	1.2
Snowmelt (mm)	winter	1.1	1.4	37.2
Snowmelt (mm)	spring	212.2	289.9	36.6
Snowmelt (mm)	summer	165.0	63.8	-61.3
Snowmelt (mm)	fall	30.9	25.9	-16.2

Annual snowpack signals

Four derived annual scalars capture the snowpack signals the literature treats as headline metrics for snow hydrology: peak snowpack (Mote et al. 2018, 2005), the date of melt midpoint (Stewart et al. 2005; Cayan et al. 2001), freshet flashiness, and the fraction of precipitation falling as snow (Knowles et al. 2006).




Figure 5.7: Annual peak snow water equivalent (swe_max) for the FWCP Peace Region. ERA5-Land mm snow water equivalent, regional spatial mean.




Figure 5.8: Day of year when half the annual snowmelt has accumulated (snowmelt_doy_50). Earlier dates indicate an earlier freshet centroid — directly relevant to crossing-design hydrology and to migration timing.




Figure 5.9: Annual maximum of 7-day rolling daily snowmelt (snowmelt_rate_peak). Higher values indicate more concentrated freshet pulses on the snowpack side, upstream of routing through soil and channel storage.




Figure 5.10: Annual snowfall fraction: the percent of annual precipitation that fell as snow. Anomaly is in percentage points relative to the 1951-1980 baseline.

What this means for the FWCP Peace Region

Snow is leaving the region earlier each year, not falling less. Total annual snowfall is roughly stable (-6 %); the snowpack decline is dominated by *earlier* melt, not by *less* snow. The seasonal data make this concrete: spring snowmelt is up 37 % while summer snowmelt is down 61 % — same total melt, redistributed earlier into the year. This matches the broader Pacific-Northwest signal documented since Mote et al. (2005).

The freshet is shifting into spring. Spring snowmelt up 37 % is direct evidence of an earlier freshet centroid — consistent in direction and rough magnitude with the 1–4 week earlier streamflow timing across western North America in Stewart et al. (2005), and with the 10-day Fraser freshet advance in Kang et al. (2016) for a basin whose southern boundary is just south of the Peace.

Summer is becoming snow-free. Summer snow water equivalent has collapsed by 75 % and summer snowmelt is down 61 %. The high-elevation snowpack that historically lingered into summer no longer does in the recent decade. For aquatic ecosystems downstream this is a loss of late-season cold-water input to streams during the warmest part of the year — the weeks where stream-temperature stress is highest for cold-water salmonids.

Spatial pattern of warming

The regional zonal mean reduces a region this size to a single number; the spatial pattern carries the rest of the story.

Warming is not spatially uniform. Total departures range from about +1.5 °C in the southeast to over +2.1 °C in the west and centre (Figure 5.11). The dominant gradient runs east-to-west — the western half of the region averages roughly 0.2 °C more warming than the eastern half — with a smaller south-to-north component. The west-of-Rockies pattern is consistent with windward-slope amplification along the Continental Divide (Pepin et al. 2015).




Figure 5.11: Spatial pattern of annual mean temperature departure across the FWCP Peace Region (2015-2025 mean minus 1951-1980 mean, degrees C). Warming is broad but uneven — the west-of-Rockies pattern is consistent with windward-slope amplification along the Continental Divide.

Per-ecoregion variation

The regional zonal mean averages over five ecoregions with different elevations and exposures. Running the same pipeline on each ecoregion polygon tests whether the regional story holds within each ecoregion — and where it does not.

All five ecoregions warmed at roughly the same rate. Precipitation is the variable that diverges: the two northernmost ecoregions (Boreal Mountains and Plateaus, Northern Canadian Rocky Mountains) show statistically significant precipitation increases, while the southern and central ecoregions do not. Vapour pressure deficit is up significantly in all five.




Figure 5.12: Annual mean temperature anomaly relative to the 1951-1980 baseline, by ecoregion. Dashed line is the 75-year Theil-Sen trend (1951-present); solid line is the 45-year trend (1981-present). A solid line steeper than the dashed line indicates accelerating warming.

Per-ecoregion variation




Figure 5.13: Annual precipitation anomaly (% of the 1951-1980 baseline) by ecoregion. The two northernmost ecoregions (BMP, NRM) show statistically significant precipitation increases over the full record; the others do not.

Snow by ecoregion

Two snow metrics are worth viewing per-ecoregion: peak snow water equivalent (annual snowpack magnitude) and DOY-50 (the day-of-year by which half the year's snowmelt has already happened — earlier values mean an earlier freshet). Pair them with the Watershed Groups Across Ecoregions section below to read each watershed group's dominant ecoregion signal.




Figure 5.14: Annual peak snow water equivalent anomaly by ecoregion, relative to the 1951-1980 baseline. Bars are mm of water-equivalent snowpack departure from baseline.

Per-ecoregion variation




Figure 5.15: Snowmelt 50 % day-of-year (DOY-50) anomaly by ecoregion, relative to the 1951-1980 baseline. Negative values (red) mean the freshet midpoint shifted earlier in the year.

Table 5.7: Per-ecoregion roll-up over the 75-year window (1951-present): annual mean / daytime maximum / overnight minimum temperature trends (°C per decade); annual precipitation trend (mm per year) and Mann-Kendall p-value; annual vapour pressure deficit trend (hPa per decade) and p-value; and recent (2015-2025) versus pre-warming (1951-1980) percent change for precipitation and soil moisture. All temperature p-values are below 0.001. A tmin slope greater than the tmax slope indicates the day-night asymmetry.

Ecoregion	tmean °C/dec	tmax °C/dec	tmin °C/dec	prcp mm/yr	prcp p	vpd hPa/dec	vpd p	prcp % change	soil moisture % chg
FAB	0.29	0.26	0.30	0.032	0.634	0.061	0.002	0.8	-1.3
CRM	0.29	0.25	0.30	0.077	0.253	0.045	0.001	4.1	-0.5
OMM	0.32	0.29	0.34	0.052	0.421	0.054	0.000	2.5	-0.7
BMP	0.30	0.27	0.33	0.144	0.023	0.033	0.003	6.3	0.3
NRM	0.29	0.26	0.31	0.156	0.015	0.030	0.004	6.6	0.6

Ecoregion codes: FAB — Fraser Basin; CRM — Central Canadian Rocky Mountains; OMM — Omineca Mountains; BMP — Boreal Mountains and Plateaus; NRM — Northern Canadian Rocky Mountains.

Watershed groups across ecoregions

The FWCP Peace Region is reported and managed at the watershed- group scale — the British Columbia Freshwater Atlas hydrological reporting unit, and the unit the FWCP funds work in. Sixteen watershed groups make up the canonical FWCP Peace list. They span the five ecoregions in different ways: some sit entirely within one, others split across two or three (Figure [5.16](#)). The table that follows gives, for each watershed group, the share of its area falling in each ecoregion — the cross-reference that maps the regional and ecoregion climate signals onto the per-watershed prioritisation unit.




Figure 5.16: Watershed groups intersecting the FWCP Peace Region (full extent, including parts spilling outside the FWCP boundary), labelled with their codes, on top of ecoregion fills.

Table 5.8: Watershed groups in the FWCP Peace Region with the percent of each group's area falling in each of the five ecoregions. Rows where one ecoregion is at or near 100 % indicate watershed groups contained within a single climate-physiography zone; rows split across two or more ecoregions indicate watershed groups whose climate departure can pull from more than one regional pattern.

Code	Name	km ² in FWCP	FAB %	CRM %	OMM %	BMP %	NRM %
CARP	Carp Lake	1794	100.0	0.0	0.0	0.0	0.0
CRKD	Crooked River	2172	100.0	0.0	0.0	0.0	0.0
FINA	Finlay Arm	7383	0.3	20.2	61.8	10.3	7.4
FINL	Finlay River	5479	0.0	0.0	0.0	30.4	69.6
FIRE	Firesteel River	4376	0.0	0.0	1.0	99.0	0.0
FOXR	Fox River	4279	0.0	0.0	0.0	16.8	83.2
INGR	Ingenika River	5328	0.0	0.0	0.7	99.3	0.0
LOMI	Lower Omineca River	4013	0.0	0.0	100.0	0.0	0.0
MESI	Mesilinka River	3294	0.0	0.0	94.8	5.2	0.0
NATP	Nation River	6880	10.0	0.0	50.7	0.0	0.0

Interpretation for fish passage

Four findings tie the regional climate-departure analysis to per-watershed barrier prioritisation.

Warming is broad and uniform; the gradient is small. All five ecoregions warmed +1.7 to +1.9 °C since 1951 — temperature is not the variable that *separates* watershed groups within the Peace. What it *does* set is the regional context: every reach in the project area is operating in air ~1.8 °C warmer than the conditions the channel, floodplain, and fish community evolved under. The within-region thermal differentiation comes from elevation, aspect, riparian cover, and groundwater contribution — not from climate-departure magnitude.

Precipitation is increasing significantly only in the two northernmost ecoregions. Watershed groups dominated by Boreal Mountains and Plateaus (Toodoggone, Firesteel, Ingenika) or Northern Canadian Rocky Mountains (Finlay, Fox) sit on statistically defensible positive precipitation trends. The other eleven watershed groups sit in Fraser Basin, Central Canadian Rockies, or Omineca Mountains, where precipitation has not changed in a way distinguishable from natural variability. For barrier prioritisation in those eleven, the climate-driven hydrology story is dominated by *atmospheric drying* (rising VPD, flat soil moisture, snowpack timing) rather than precipitation magnitude.

The atmosphere is drying despite, in places, more precipitation. Vapour pressure deficit rose significantly in every ecoregion. Soil moisture is flat across the region even where precipitation rose

— the warmer atmosphere is pulling the extra water back through evaporation and transpiration. The ecological implication is that streamflow during the summer low-flow period — the thermal-stress weeks where cold-water rearing habitat is at the margin — is not getting the relief that the precipitation increase, where it occurred, would otherwise suggest.

Snow is leaving the region earlier, not falling less. The snowpack-timing signal is regionally uniform — DOY-50 shifted earlier in every ecoregion. The freshet — the dominant high-flow event that mobilises spawning gravels, refills off-channel rearing habitat, and that crossing structures are sized against — is arriving weeks earlier across all 16 watershed groups. Summer snow water equivalent collapsed 75 %; the cold-water input that high-elevation snowpack provides to streams during the warmest weeks of summer is dropping in parallel. Both signals act on barrier prioritisation through hydrology and stream temperature rather than through which watershed they hit hardest.

For the cold-water resident salmonids the FWCP Peace supports — bull trout, Arctic grayling, mountain whitefish, rainbow trout, kokanee — the implication for barrier prioritisation is that two fundamentally different climate-departure stories play out across the 16 watershed groups, and that the priority weight on barrier removal should respond to both: warmer headwater reaches that gain growing-degree-days have *more* habitat to recover above the barrier, while reaches sitting closer to the upper thermal niche have less. The watershed-group ecoregion mapping in the table above is the bridge between the regional climate-departure summary in the main body of this report and the per-watershed weight applied to each barrier.

Appendix - Floodplain Delineation

This appendix details the floodplain delineation for the Parsnip River Watershed Group. The rationale for mapping floodplain extent alongside the fish passage barrier inventory is described in the report background — in brief, the off-channel habitats that floodplains sustain are often as important to fish productivity as the mainstem passage that crossing remediations restore, and infrastructure corridors can compromise floodplain function in ways that structure replacement alone does not address.

Floodplain footprints were delineated using the [flooded](#) R package, which implements the Valley Confinement Algorithm of Nagel et al. (2014). The algorithm combines a digital elevation model with the stream network and a modelled bankfull flood surface to distinguish valley bottoms from confined canyon reaches. Bankfull depth — the flow depth at which a stream begins to spill onto its floodplain — is estimated from upstream watershed area and mean annual precipitation using the regional regression of Hall et al. (2007), calibrated against field-mapped floodplain sites in Pacific Northwest streams.

We mapped the **functional floodplain**: the area a river actively inundates and reworks during regular high-flow events. This sits between two other scales — the narrow active channel (the wetted perimeter at low flow) and the broader geologic valley bottom (which includes terraces untouched by water for centuries). The functional floodplain matters most for habitat planning because it is the ground where side channels form, seasonal flooding sustains wetlands, and riparian forests recruit. Inputs were the [MRDEM-30](#) 30 m DEM, the bull trout accessible stream network (stream order ≥ 3) from [bcfishpass](#), and BC Freshwater Atlas lakes and wetlands on that network. Waterbodies were included so the valley confinement model fills cells that gradient and cost-distance masks would otherwise exclude.

Table 5.9: Parsnip River Watershed Group — modelled functional floodplain and the off-channel habitat units within it.

Metric	Value
Watershed group area (km ²)	5,597
Floodplain area (ha)	48,116
Floodplain as % of watershed group	8.6
Bull trout accessible streams, order 3+ (km)	2,310
Streams within floodplain (km)	1,580
Lakes within floodplain (count)	198
Lakes within floodplain (ha)	1,603
Wetlands within floodplain (count)	464
Wetlands within floodplain (ha)	6,582

The Parsnip River Watershed Group covers roughly 5,597 km². Modelled functional floodplain (Table 5.9) covers about 48,116 ha, or 8.6 % of the watershed group, attached to 1,580 km of the 2,310 km bull trout accessible order-3+ network. Within that footprint sit 198 lakes and 464 mapped wetland polygons — the off-channel habitat units that lateral floodplain connectivity supports.

The watershed-wide footprint is shown in Figure 5.17; a detail view of the south-east headwaters around Arctic Lake is shown in Figure 5.18.

**Parsnip River Watershed Group
functional floodplain, accessible order 3+**



Figure 5.17: Parsnip River Watershed Group functional floodplain (green) over hillshaded terrain. Bull trout accessible order-3+ streams in blue; lakes and wetlands in light blue; parks light green; First Nations reserves light grey with diamond markers and labels; municipalities outlined in red; forest service / resource roads grey; railways black dashed; watershed boundary heavy black.

Parsnip River Watershed Group south-east detail



Figure 5.18: South-east corner of the Parsnip River Watershed Group at full resolution — the headwaters around Arctic Lake on the continental divide. Symbology as Figure [5.17](#); named streams labelled in italic blue; municipalities outlined and labelled in red where present.

The watershed-wide view shows the floodplain following the trunk Parsnip River north toward Williston Reservoir, with substantial off-channel area in the broader valley reaches and at tributary confluences. The detail view resolves the per-reach floodplain pattern in the Arctic Lake headwaters on the continental divide and gives a sense of the resolution available within any sub-area for downstream analyses — barrier prioritisation, land-cover change tracking, or restoration scoping. In coming years we plan to extend this mapping to additional FWCP Peace watersheds so that floodplain context can inform barrier prioritisation across the region.

Appendix - Phase 1 Fish Passage Assessment Data and Photos

Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	6559	Diameter (m)	0.9
External ID	–	Length (m)	24
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	489950	Resemble Channel	–
Northing	6112363	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay Community Connector	Fill Depth (m)	2.5
Road Tenure	Sekani R01883 83	Outlet Drop (m)	1.1
Channel Width (m)	1	Outlet Pool Depth (m)	0
Stream Slope (%)	1.5	Inlet Drop	No
Beaver Activity	Yes	Slope (%)	9
Habitat Value	Low	Valley Fill	Deep Fill
Final score	36	Barrier Result	Barrier
Fix type	Replace Structure with Streambed Simulation CBS	Fix Span / Diameter	3
Comments: Wetland upstream. Stream was not flowing at time of assessment. Small channel downstream. Beaver grate devices on inlet and outlet			
Photos: PSCIS ID 6559. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203597	Diameter (m)	0.8
External ID	15201563	Length (m)	19
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	425959	Resemble Channel	–
Northing	6122703	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Thut 15000	Fill Depth (m)	1.5
Road Tenure	Sekani R07214 8	Outlet Drop (m)	0.25
Channel Width (m)	2.5	Outlet Pool Depth (m)	0.5
Stream Slope (%)	2.5	Inlet Drop	Yes
Beaver Activity	No	Slope (%)	1
Habitat Value	High	Valley Fill	Deep Fill
Final score	29	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: High-quality stream with abundant gravels and cover, directly connected to the Nation River. Bull trout noted near the confluence. eDNA samples collected upstream and downstream of the crossing. This crossing is located on a road identified by BCTS as a candidate for potential deactivation. Photos: PSCIS ID 203597. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203598	Diameter (m)	0.8
External ID	15201146	Length (m)	10
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	426678	Resemble Channel	–
Northing	6123035	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Thut 15000	Fill Depth (m)	1.5
Road Tenure	Sekani R07214 22	Outlet Drop (m)	0.2
Channel Width (m)	1.6	Outlet Pool Depth (m)	0.1
Stream Slope (%)	3.4	Inlet Drop	No
Beaver Activity	No	Slope (%)	2
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	26	Barrier Result	Barrier
Fix type	Replace Structure with Streambed Simulation CBS	Fix Span / Diameter	3
Comments: Small stream with short channel length before entering a large wetland area. Pockets of small gravel were present throughout. This crossing is located on a road identified by BCTS as a candidate for potential deactivation. Photos: PSCIS ID 203598. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203599	Diameter (m)	2.3
External ID	15202249	Length (m)	21
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	429213	Resemble Channel	–
Northing	6126036	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Thut 14000	Fill Depth (m)	1.2
Road Tenure	MoF	Outlet Drop (m)	0
Channel Width (m)	3	Outlet Pool Depth (m)	0.4
Stream Slope (%)	1.5	Inlet Drop	No
Beaver Activity	No	Slope (%)	2
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	24	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Wetland type habitat downstream with good habitat upstream with pockets of gravel and a wide channel. Culverts beginning to sag in the middle. No outlet drop.			
Photos: PSCIS ID 203599. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203600	Diameter (m)	2.35
External ID	15200985	Length (m)	15
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	458498	Resemble Channel	–
Northing	6137654	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Finlay-Nation FSR	Fill Depth (m)	2
Road Tenure	Sekani R09952 1	Outlet Drop (m)	0.15
Channel Width (m)	4.1	Outlet Pool Depth (m)	0.1
Stream Slope (%)	6	Inlet Drop	Yes
Beaver Activity	No	Slope (%)	4
Habitat Value	High	Valley Fill	Deep Fill
Final score	34	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Large stream with abundant woody debris, pools, and gravels upstream. Outlet dropped minimally onto a boulder.			
Photos: PSCIS ID 203600. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203601	Diameter (m)	2.1
External ID	15201007	Length (m)	15
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	459772	Resemble Channel	–
Northing	6140076	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Finlay-Nation FSR	Fill Depth (m)	1.3
Road Tenure	Sekani R09952 1	Outlet Drop (m)	0
Channel Width (m)	3.5	Outlet Pool Depth (m)	0.1
Stream Slope (%)	4	Inlet Drop	No
Beaver Activity	No	Slope (%)	1
Habitat Value	High	Valley Fill	Deep Fill
Final score	24	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Large stream with pools and gravels present upstream. Significant amounts of road fill had been graded into the channel.			
Photos: PSCIS ID 203601. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203602	Diameter (m)	1.55
External ID	15201006	Length (m)	23
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	466124	Resemble Channel	–
Northing	6147887	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Finlay-Nation FSR	Fill Depth (m)	3.5
Road Tenure	Sekani R01884 61	Outlet Drop (m)	0.25
Channel Width (m)	3.4	Outlet Pool Depth (m)	0.1
Stream Slope (%)	4	Inlet Drop	No
Beaver Activity	No	Slope (%)	5.5
Habitat Value	High	Valley Fill	Deep Fill
Final score	34	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	16.5
Comments: Large, low gradient, high value system. Significant amounts of road fill is eroding from the road prism into the stream on both sides of the FSR. Habitat confirmation and electrofishing or EDNA sampling is recommended. Photos: PSCIS ID 203602. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Oval Culvert
PSCIS ID	203603	Diameter (m)	5.8
External ID	16400714	Length (m)	30
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	469020	Resemble Channel	–
Northing	6148916	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay-Nation FSR	Fill Depth (m)	2
Road Tenure	MoF	Outlet Drop (m)	0
Channel Width (m)	7.4	Outlet Pool Depth (m)	0.1
Stream Slope (%)	1.5	Inlet Drop	No
Beaver Activity	Yes	Slope (%)	2
Habitat Value	High	Valley Fill	Deep Fill
Final score	24	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Very large stream with 2.5m high beaver dam located in the middle of the culvert backing up a massive wetland upstream. First passage is definitely hindered by the beaver dam however, the wetland it creates provides likely stable hydrology for the downstream system. Photos: PSCIS ID 203603. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203604	Diameter (m)	1
External ID	16401990	Length (m)	50
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	480204	Resemble Channel	–
Northing	6138985	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay FSR	Fill Depth (m)	7
Road Tenure	Sekani R10040 25	Outlet Drop (m)	0
Channel Width (m)	10	Outlet Pool Depth (m)	1.5
Stream Slope (%)	0.5	Inlet Drop	No
Beaver Activity	Yes	Slope (%)	3
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	32	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	27
Comments: Very large system with extremely undersized pipe that is submerged at the inlet with obvious beaver activity within the pipe itself. Upstream is a huge wetland, beaver-influenced area. Culvert slope estimated as the outlet pool is extremely deep.			
Photos: PSCIS ID 203604. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203605	Diameter (m)	3.9
External ID	16401519	Length (m)	46
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	486504	Resemble Channel	–
Northing	6124990	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay FSR	Fill Depth (m)	2.5
Road Tenure	MoF	Outlet Drop (m)	0.95
Channel Width (m)	6	Outlet Pool Depth (m)	0.5
Stream Slope (%)	2	Inlet Drop	No
Beaver Activity	No	Slope (%)	2
Habitat Value	High	Valley Fill	Deep Fill
Final score	37	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: High value habitat on very large system with pipe that has large outlet drop near 1m. Habitat confirmation and eDNA sampling conducted upstream and downstream of the crossing.			
Photos: PSCIS ID 203605. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Oval Culvert
PSCIS ID	203606	Diameter (m)	4.6
External ID	16401529	Length (m)	35
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	486441	Resemble Channel	–
Northing	6119999	Backwatered	No
Stream	Scovil Creek	Percent Backwatered	–
Road	Finlay FSR	Fill Depth (m)	3
Road Tenure	MoF	Outlet Drop (m)	0.2
Channel Width (m)	9.3	Outlet Pool Depth (m)	0.7
Stream Slope (%)	2	Inlet Drop	No
Beaver Activity	No	Slope (%)	2
Habitat Value	High	Valley Fill	Deep Fill
Final score	32	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Huge stream with abundant gravels and low gradient complex habitat upstream. Smaller outlet drop however pipe is somewhat undersized for the watershed.			
Photos: PSCIS ID 203606. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	–
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203607	Diameter (m)	1.65
External ID	16400428	Length (m)	16
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	488459	Resemble Channel	–
Northing	6112370	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finl 19000	Fill Depth (m)	2
Road Tenure	BCTS	Outlet Drop (m)	0
Channel Width (m)	1.5	Outlet Pool Depth (m)	0
Stream Slope (%)	1.5	Inlet Drop	Yes
Beaver Activity	No	Slope (%)	2.5
Habitat Value	Low	Valley Fill	Deep Fill
Final score	18	Barrier Result	Potential
Fix type	Replace Structure with Streambed Simulation CBS	Fix Span / Diameter	3
Comments: Dry, poorly defined channel. Unlikely fish habitat.			
Photos: PSCIS ID 203607. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203608	Diameter (m)	2.6
External ID	16491533	Length (m)	35
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	479612	Resemble Channel	–
Northing	6141425	Backwatered	Yes
Stream	Dastaiga Creek	Percent Backwatered	50
Road	Finlay FSR	Fill Depth (m)	1.6
Road Tenure	Sekani R01886 140	Outlet Drop (m)	0
Channel Width (m)	8	Outlet Pool Depth (m)	0.25
Stream Slope (%)	3	Inlet Drop	Yes
Beaver Activity	Yes	Slope (%)	1.5
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	27	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: Very large stream with a beaver dam located 30m upstream of the crossing. No outlet drop, but culvert is well undersized for the watershed. Stream width and gradient are rough estimates.			
Photos: PSCIS ID 203608. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203609	Diameter (m)	3.5
External ID	16401530	Length (m)	30
Crew	AI	Embedded	Yes
UTM Zone	10	Depth Embedded (m)	0.27
Easting	486707	Resemble Channel	Yes
Northing	6116611	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay FSR	Fill Depth (m)	3
Road Tenure	Sekani R06855 33	Outlet Drop (m)	0
Channel Width (m)	3.7	Outlet Pool Depth (m)	0.03
Stream Slope (%)	1	Inlet Drop	No
Beaver Activity	No	Slope (%)	0.5
Habitat Value	High	Valley Fill	Deep Fill
Final score	14	Barrier Result	Passable
Fix type	–	Fix Span / Diameter	–
Comments: Low gradient, larger system with abundant gravel. Culvert is fully embedded. Culvert length estimated based on photos.			
Photos: PSCIS ID 203609. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203610	Diameter (m)	2.9
External ID	16401547	Length (m)	30
Crew	AI	Embedded	Yes
UTM Zone	10	Depth Embedded (m)	0.6
Easting	476482	Resemble Channel	yes
Northing	6147549	Backwatered	Yes
Stream	Tributary To Williston Reservoir	Percent Backwatered	50
Road	Finlay FSR	Fill Depth (m)	2
Road Tenure	Sekani R10040 19	Outlet Drop (m)	0
Channel Width (m)	2	Outlet Pool Depth (m)	0.1
Stream Slope (%)	0	Inlet Drop	Yes
Beaver Activity	Yes	Slope (%)	1
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	11	Barrier Result	Passable
Fix type	–	Fix Span / Diameter	–
Comments: Beaver dam located 1m upstream from the inlet. Pipe is embedded and the top 10m is backwatered. Smaller stream with some pockets of gravel and decent flow – although there were heavy rains yesterday. Photos: PSCIS ID 203610. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Bridge
PSCIS ID	203611	Diameter (m)	18.3
External ID	16401393	Length (m)	5
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	486421	Resemble Channel	–
Northing	6124806	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Modeste FSR	Fill Depth (m)	–
Road Tenure	MoF	Outlet Drop (m)	0
Channel Width (m)	0	Outlet Pool Depth (m)	0
Stream Slope (%)	0	Inlet Drop	No
Beaver Activity	No	Slope (%)	0
Habitat Value	–	Valley Fill	–
Final score	10	Barrier Result	Passable
Fix type	–	Fix Span / Diameter	–
Comments: Bridge			
Photos: PSCIS ID 203611. From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			



PSCIS crossing 199663 was first assessed in 2024 as part of this project and was ranked as a barrier to fish passage with medium value habitat documented (Irvine and Schick 2025). The crossing was reassessed in 2025, and a habitat confirmation was completed due to the presence of medium value habitat and a significant outlet drop at the culvert. Just upstream of the confluence of this tributary and the Parsnip River, grayling have been documented (Norris [2018] 2024; MoE 2024a).

A summary of habitat modelling outputs for the crossing are presented in Table [5.11](#).

Table 5.10: Summary of derived upstream watershed statistics for PSCIS crossing 199663.

Site	Area Km	Elev Site	Elev Min	Elev Max	Elev Median	Elev P60	Aspect
199663	1	713	707	849	784	774	SW

* Elev P60 = Elevation at which 60% of the watershed area is above

Table 5.11: Summary of fish habitat modelling for PSCIS crossing 199663.

Habitat	Potential	Remediation Gain	Remediation Gain (%)
BT Rearing (km)	1.6	1.6	100
BT Spawning (km)	0.7	0.7	100
BT Network (km)	9.9	9.3	94
BT Stream (km)	8.0	7.4	92
BT Lake Reservoir (ha)	0.5	0.5	100
BT Wetland (ha)	12.8	12.8	100
BT Slopeclass03 (km)	3.6	3.0	83
BT Slopeclass05 (km)	3.6	3.6	100
BT Slopeclass08 (km)	0.9	0.9	100
BT Slopeclass15 (km)	0.2	0.2	100

* Model data is preliminary and subject to adjustments.

Stream Characteristics at Crossing 199663

At the time of the 2025 assessment, PSCIS crossing 199663 on Chco 12000 was un-embedded, non-backwatered and ranked as barrier to upstream fish passage according to the provincial

protocol (MoE 2011) (Table [5.12](#)). The culvert had a 0.6m outlet drop and a 0.4m deep outlet pool.

The water temperature was 7°C, pH was 8.2 and conductivity was 310 uS/cm.

Table 5.12: Summary of fish passage assessment for PSCIS crossing 199663.

Location and Stream Data		Crossing Characteristics	
Date	2025-09-21	Crossing Sub Type	Round Culvert
PSCIS ID	199663	Diameter (m)	1.2
External ID	–	Length (m)	13
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	515416	Resemble Channel	–
Northing	6094963	Backwatered	No
Stream	Tributary To Parsnip River	Percent Backwatered	–
Road	Chco 12000	Fill Depth (m)	0.5
Road Tenure	BCTS	Outlet Drop (m)	0.6
Channel Width (m)	0	Outlet Pool Depth (m)	0.4
Stream Slope (%)	1	Inlet Drop	No
Beaver Activity	No	Slope (%)	3
Habitat Value	Medium	Valley Fill	Deep Fill
Final score	30	Barrier Result	Barrier
Fix type	Replace Structure with Streambed Simulation CBS	Fix Span / Diameter	3
Comments: Only photos and downstream width measurements were collected in 2025 due to limited time; all other information was transcribed from the 2024 survey. Habitat confirmations and eDNA samples were collected both upstream and downstream. A large wetland was located approximately 70–80m downstream of the culvert outlet. Abundant gravels were present within the first 20–30m below the outlet pool, where flow was strong, and a substantial outlet drop was observed. Photos: From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			

Stream Characteristics at Crossing 19...



Stream Characteristics Downstream of Crossing 199663

The stream was surveyed downstream from crossing 199663 for 80m (Figure 5.20). The habitat was rated as medium value for salmonid spawning and rearing. The stream had good flow and abundant gravels for 30–40m downstream of the culvert. A wetland was located ~80m downstream of the FSR. The average channel width was 2.6m and the average wetted width was 2.8m. The dominant substrate was cobbles with gravels sub-dominant. Total cover amount was rated as abundant with deep pools dominant. Cover was also present as small woody debris, large woody debris, undercut banks, and overhanging vegetation.

Stream Characteristics Upstream of Crossing 199663

The stream was surveyed upstream from crossing 199663 for 600m (Figure 5.20). The habitat was rated as medium value. Total cover amount was rated as abundant with overhanging vegetation dominant. Cover was also present as small woody debris, large woody debris, and undercut banks. The dominant substrate was fines with gravels sub-dominant. The average channel width was 3m, the average wetted width was 2.1m, and the average gradient was 2.8%. The stream was low gradient with significant riparian tree blowdown throughout. Wide floodplain within a well defined canyon (approximately 15m deep with slopes of ~45°). Healthy riparian consisting of herb and shrub cover within sparse spruce forest that appears to have been impacted by some sort of beetle or other affliction. Most of the spruce trees were dead with many already fallen down.

Environmental DNA

eDNA samples were collected both upstream and downstream of crossing 199663 on Chco 12000, with sampling techniques summarized in Table 5.13. Samples were submitted to UNBC for ddPCR analysis against Bull Trout, Arctic Grayling, Rainbow Trout, Kokanee, and Burbot assays (Burbot was tested only at this crossing among the Peace 2025 sites). Per-position results are summarized in Table 5.14. Detections are based on the standard ddPCR call threshold of ≥4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran.

Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#), as well as within the [interactive Peace eDNA map](#).

Table 5.13: Summary of eDNA samples collected at site 199663.

Site	Stream	Habitat Type	Sample Method	Filter Type	Filter Size (um)	Volume filtered (L)	Site Description
199663_ds_ed1	Tributary To Parsnip						

Site	Stream	Habitat Type	Sample Method	Filter Type	Filter Size (um)	Volume filtered (L)	Site Description
River	riffle	instream	mce	1	2.0		Sample collected 15m downstream of culvert right bank. Site was chosen close to the culvert because of the wetland that was just downstream of this area - and wanted some flow.
199663_us_ed1	Tributary To Parsnip River	riffle	instream	mce	1	2.9	Sample collected 25m upstream of culvert inlet on the right bank

Table 5.14: eDNA detection results at crossing 199663.

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
199663_ds_ed1	M10139	BURB(195), RAIN(112)	—	BULT(0), GRAY(0)
199663_us_ed1	M10140	RAIN(65)	—	BULT(0), BURB(0), GRAY(0)

Structure Remediation and Cost Estimate

Should restoration/maintenance activities proceed, replacement of the Chco 12000 crossing (199663) with a streambed simulation (3m span) is recommended. At the time of reporting in 2026, the cost of the work is estimated at \$300,000.

Conclusion

The habitat upstream of PSCIS crossing 199663 on Chco 12000 was documented as medium value for salmonid spawning and rearing, and the crossing is rated as a moderate priority for replacement due to the large outlet drop. Environmental DNA sampling at the site confirmed rainbow trout both upstream and downstream of the crossing; no bull trout were detected. Burbot were detected downstream but not upstream.

Since the assessment season, the FWCP 2026 proposal for replacement of this crossing was accepted and 50% project funding is confirmed. Replacement is targeted for 2026 in coordination with BC Timber Sales; an engineering firm is expected to collect the data required for the site plan.

Table 5.15: Summary of habitat details for PSCIS crossing 199663.

Site	Location	Length Surveyed (m)	Average Channel Width (m)	Average Wetted Width (m)	Average Pool Depth (m)	Average Gradient (%)	Total Cover	Habitat Value
199663	Downstream	80	2.6	2.8	–	–	abundant	–
199663	Upstream	600	3.0	2.1	0.2	2.8	abundant	medium

Conclusion



Figure 5.20: Left: Typical habitat downstream of PSCIS crossing 199663. Right: Typical habitat upstream of PSCIS crossing 199663.

Tributary To Nation River - 203597 - Appendix

Site Location

PSCIS crossing 203597 is located on Tributary To Nation River, approximately 65km west of McKenzie, BC, as the crow flies, and is in the Nation River watershed group (Figure 5.21). The crossing is located 1.9m upstream of the stream's confluence with the Nation River on Thut 15000 road and is situated under Sekani Forest Products Ltd. road tenure R07214 8.




Figure 5.21: Map of Tributary To Nation River

Background

At PSCIS crossing 203597, Tributary To Nation River is a second order stream and drains a watershed of approximately 8.2km². The watershed ranges in elevation from a maximum of 1065m to 900m near the crossing (Table 5.16).

This was the first assessment of PSCIS crossing 203597, in part because the site is situated on a road identified by BC Timber Sales as a potential deactivation candidate. The stream exhibited high-quality habitat, so a habitat confirmation was completed. Bull trout are documented in the Nation River near the confluence with this stream (Norris [2018] 2024; MoE 2024a).

A summary of habitat modelling outputs for the crossing are presented in Table [5.17](#).

Table 5.16: Summary of derived upstream watershed statistics for PSCIS crossing 203597.

Site	Area Km	Elev Site	Elev Min	Elev Max	Elev Median	Elev P60	Aspect
203597	8.2	900	907	1065	980	971	S

* Elev P60 = Elevation at which 60% of the watershed area is above

Table 5.17: Summary of fish habitat modelling for PSCIS crossing 203597.

Habitat	Potential	Remediation Gain	Remediation Gain (%)
BT Rearing (km)	5.6	5.6	100
BT Spawning (km)	0.7	0.7	100
BT Network (km)	16.8	9.5	57
BT Stream (km)	12.2	5.2	43
BT Lake Reservoir (ha)	1.0	1.0	100
BT Wetland (ha)	47.1	46.2	98
BT Slopeclass03 (km)	7.9	2.5	32
BT Slopeclass05 (km)	4.0	2.5	62
BT Slopeclass08 (km)	0.5	0.5	100
BT Slopeclass15 (km)	0.0	0.0	–

* Model data is preliminary and subject to adjustments.

Stream Characteristics at Crossing 203597

At the time of the 2025 assessment, PSCIS crossing 203597 on Thut 15000 was un-embedded, non-backwatered and ranked as barrier to upstream fish passage according to the provincial protocol (MoE 2011) (Table [5.18](#)).

Stream Characteristics at Crossing 20...

The water temperature was 6.5°C, pH was 7.5 and conductivity was 188 uS/cm.

Table 5.18: Summary of fish passage assessment for PSCIS crossing 203597.

Location and Stream Data		Crossing Characteristics	
Date	2025-09-22	Crossing Sub Type	Round Culvert
PSCIS ID	203597	Diameter (m)	0.8
External ID	15201563	Length (m)	19
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	425959	Resemble Channel	–
Northing	6122703	Backwatered	No
Stream	Tributary To Nation River	Percent Backwatered	–
Road	Thut 15000	Fill Depth (m)	1.5
Road Tenure	Sekani R07214 8	Outlet Drop (m)	0.25
Channel Width (m)	2.5	Outlet Pool Depth (m)	0.5
Stream Slope (%)	2.5	Inlet Drop	Yes
Beaver Activity	No	Slope (%)	1
Habitat Value	High	Valley Fill	Deep Fill
Final score	29	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: High-quality stream with abundant gravels and cover, directly connected to the Nation River. Bull trout noted near the confluence. eDNA samples collected upstream and downstream of the crossing. This crossing is located on a road identified by BCTS as a candidate for potential deactivation.			
Photos: From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			

Stream Characteristics at Crossing 20...



Stream Characteristics Downstream of Crossing 203597

The stream was surveyed downstream from crossing 203597 for 400m (Figure [5.22](#)). The stream had good flow, likely elevated due to significant precipitation the previous night. Abundant gravels were present throughout, and cover was provided by multiple sources including woody debris and undercut banks. No barriers were observed. Total cover amount was rated as abundant with undercut banks dominant. Cover was also present as small woody debris, large woody debris, boulders, deep pools, and overhanging vegetation. The dominant substrate was cobbles with gravels sub-dominant. The average channel width was 2.8m, the average wetted width was 2.6m, and the average gradient was 3%.

Stream Characteristics Upstream of Crossing 203597

The stream was surveyed upstream from crossing 203597 for 350m up to the beginning of the wetlands (Figure [5.22](#)). The habitat was rated as NA value. Total cover amount was rated as moderate with small woody debris dominant. Cover was also present as large woody debris, undercut banks, and overhanging vegetation. The average channel width was 2.5m, the average wetted width was 2.4m, and the average gradient was 3%. The dominant substrate was fines with cobbles sub-dominant. The stream had good flow, although recent rainfall may have influenced discharge. Pockets of gravels were present within the area surveyed. Wetlands adjacent to the channel appeared shallow, suggesting they may freeze to the bottom during winter.

Environmental DNA

eDNA samples were collected both upstream and downstream of crossing 203597 on Thut 15000, with sampling techniques summarized in Table [5.19](#). Samples were submitted to UNBC for ddPCR analysis against Bull Trout, Arctic Grayling, Rainbow Trout, and Kokanee assays. Per-position results from real environmental samples are summarized in Table [5.20](#). Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran.

Field blank at this crossing. Sample 203597_ds_ed1a (UNBC ID M41365) was a field blank — distilled water filtered at the field site to test for protocol contamination, not site eDNA. UNBC's analysis returned a low-level Rainbow Trout signal at this blank (4 positive droplets, ≈ 0.27 copies/ μ L). Per ddPCR convention this represents protocol-side contamination during collection or processing, not species presence at this location. The blank is presented in the Controls layer of the interactive map (off by default) and in the field-blanks table within [Appendix - Environmental DNA Results \(page 145\)](#).

Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#), as well as within the [interactive Peace eDNA map](#).

Table 5.19: Summary of eDNA samples collected at site 203597.

Site	Stream	Habitat Type	Sample Method	Filter Type	Filter Size (um)	Volume filtered (L)	Site Description
203597_ds_ed1b	Tributary To Nation River	riffle	grab	mce	1	1.7	Sample collected 350 - 400m downstream of culvert. Right bank. Accessed via historic road which is now wildlife trail.
203597_us_ed1	Tributary To Nation River	riffle	inatream	mce	1	2.0	Sample collected 5m upstream of the culvert on the left bank

Table 5.20: eDNA detection results at crossing 203597 (real environmental samples).

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
203597_ds_ed1b	M41366	RAIN(173)*	BULT(3)*	GRAY(0), SOCK(0)
203597_us_ed1	M41367	RAIN(213)*	—	BULT(0), GRAY(0), SOCK(0)*

Structure Remediation and Cost Estimate

Since the time of field assessments this crossing was deactivated by BC Timber Sales.

Conclusion

The habitat upstream of PSCIS crossing 203597 on Thut 15000 was documented as medium value for salmonid spawning and rearing, and the crossing was rated as a high for deactivation.

Since the assessment season, the FWCP 2026 proposal for riparian enhancements at this site was accepted. Planned work consists of soil decompaction along the deactivated road prism, riparian planting with native species, and placement of coarse woody debris to advance recovery of the channel margin. Environmental DNA sampling confirmed rainbow trout both upstream and downstream of the crossing; an under-threshold bull trout detection is suspected to reflect real presence given the suitable size of the stream and its proximity to the Nation River, where bull trout detections are recorded in provincial records near the subject stream's confluence with the Nation.

Table 5.21: Summary of habitat details for PSCIS crossing 203597.

Site	Location	Length Surveyed (m)	Average Channel Width (m)	Average Wetted Width (m)	Average Pool Depth (m)	Average Gradient (%)	Total Cover	Habitat Value
203597	Downstream	400	2.8	2.6	0.3	3	abundant	high
203597	Upstream	350	2.5	2.4	0.3	3	moderate	–

Conclusion



Figure 5.22: Left: Typical habitat downstream of PSCIS crossing 203597. Right: Wetland habitat located 350m upstream of PSCIS crossing 203597.

Tributary To Williston Reservoir - 203605 - Appendix

Site Location

PSCIS crossing 203605 is located on Tributary To Williston Reservoir, on the western side of the reservoir, approximately 10km southeast of MacKenzie, BC, in the Parsnip Arm watershed group (Figure 5.23). The crossing is located 1.3m upstream of the stream's confluence with the Williston Reservoir on Finlay FSR road which is the responsibility of the Ministry of Forests.




Figure 5.23: Map of Tributary To Williston Reservoir

Background

At PSCIS crossing 203605, Tributary To Williston Reservoir is a fourth order stream and drains a watershed of approximately 55.6km². The watershed ranges in elevation from a maximum of 1644m to 681m near the crossing (Table 5.22).

This was the first assessment of PSCIS crossing 203605, selected because it is located very close to Williston Reservoir, which supports documented bull trout. Approximately 35km of bull trout rearing habitat is modelled upstream with no additional modelled barriers, and rainbow trout have previously been recorded in the system (Norris [2018] 2024; MoE 2024a). The stream exhibited high-quality habitat, therefore a habitat confirmation was completed.

Approximately 215m upstream of Finlay FSR, the stream crosses Modeste FSR at a bridge, which was also assessed for the first time in 2025 (PSCIS crossing 203611). Modeste FSR provides access to the lower half of the watershed, and is identified by BC Timber Sales as a potential deactivation candidate.

A summary of habitat modelling outputs for the crossing are presented in Table [5.23](#).

Table 5.22: Summary of derived upstream watershed statistics for PSCIS crossing 203605.

Site	Area Km	Elev Site	Elev Min	Elev Max	Elev Median	Elev P60	Aspect
203605	55.6	681	671	1644	980	937	SE

* Elev P60 = Elevation at which 60% of the watershed area is above

Table 5.23: Summary of fish habitat modelling for PSCIS crossing 203605.

Habitat	Potential	Remediation Gain	Remediation Gain (%)
BT Rearing (km)	35.9	30.8	86
BT Spawning (km)	22.0	19.5	89
BT Network (km)	100.5	76.4	76
BT Stream (km)	86.7	64.1	74
BT Lake Reservoir (ha)	51.1	42.4	83
BT Wetland (ha)	51.9	50.3	97
BT Slopeclass03 (km)	16.3	14.4	88
BT Slopeclass05 (km)	21.8	10.4	48
BT Slopeclass08 (km)	17.6	12.2	69
BT Slopeclass15 (km)	28.2	24.4	87

* Model data is preliminary and subject to adjustments.

Stream Characteristics at Crossing 203605

At the time of the 2025 assessment, PSCIS crossing 203605 on Finlay FSR was un-embedded, non-backwatered and ranked as barrier to upstream fish passage according to the provincial protocol (MoE 2011) (Table [5.24](#)). The culvert had a 0.95m outlet drop and a 0.5m deep outlet pool.

The water temperature was 6.8°C, pH was 7.6 and conductivity was 251 uS/cm.

Table 5.24: Summary of fish passage assessment for PSCIS crossing 203605.

Location and Stream Data		Crossing Characteristics	
Date	2025-09-23	Crossing Sub Type	Round Culvert
PSCIS ID	203605	Diameter (m)	3.9
External ID	16401519	Length (m)	46
Crew	AI	Embedded	No
UTM Zone	10	Depth Embedded (m)	–
Easting	486504	Resemble Channel	–
Northing	6124990	Backwatered	No
Stream	Tributary To Williston Reservoir	Percent Backwatered	–
Road	Finlay FSR	Fill Depth (m)	2.5
Road Tenure	MoF	Outlet Drop (m)	0.95
Channel Width (m)	6	Outlet Pool Depth (m)	0.5
Stream Slope (%)	2	Inlet Drop	No
Beaver Activity	No	Slope (%)	2
Habitat Value	High	Valley Fill	Deep Fill
Final score	37	Barrier Result	Barrier
Fix type	Replace with New Open Bottom Structure	Fix Span / Diameter	15
Comments: High value habitat on very large system with pipe that has large outlet drop near 1m. Habitat confirmation and eDNA sampling conducted upstream and downstream of the crossing.			
Photos: From top left clockwise: Road/Site Card, Barrel, Outlet, Downstream, Upstream, Inlet.			

Stream Characteristics at Crossing 20...



Stream Characteristics Downstream of Crossing 203605

The stream was surveyed downstream from crossing 203605 for 250m to the mapped reservoir location, where the channel retained stream characteristics at the time of assessment (Figure 5.24). Abundant gravels and cover were present throughout, with numerous run sections providing suitable rearing habitat for juvenile westslope cutthroat trout and rainbow trout. No barriers to fish passage were observed in the area surveyed and the habitat was rated as high value for salmonid spawning and rearing. Total cover amount was rated as moderate with dominant. Cover was also present as large woody debris and overhanging vegetation. The dominant substrate was gravels with fines sub-dominant. The average channel width was 8.5m, the average wetted width was 5.6m, and the average gradient was 1.8%.

Stream Characteristics Upstream of Crossing 203605

The stream was surveyed upstream from crossing 203605 for 650m (Figure 5.24). The habitat was rated as high value. The average channel width was 6.7m, the average wetted width was 4.6m, and the average gradient was 2.9%. The dominant substrate was gravels with cobbles sub-dominant. Total cover amount was rated as moderate with undercut banks dominant. Cover was also present as small woody debris, large woody debris, boulders, deep pools, and overhanging vegetation. The stream was large with complex habitat and diverse cover including deep pools, boulders, large and small woody debris, and undercut banks. A canyon section ~100m long occurred ~100m upstream of the Modeste FSR bridge, containing multiple cascade steps up to 1m high with deep downstream pools. At the upstream end of the site, the channel deflected against steep gully walls. Abundant gravels were present, suitable for adfluvial and resident rainbow trout as well as bull trout.

Environmental DNA

eDNA samples were collected both upstream and downstream of crossing 203605 on Finlay FSR, with sampling techniques summarized in Table 5.25. Samples were submitted to UNBC for ddPCR analysis against Bull Trout, Arctic Grayling, Rainbow Trout, and Kokanee assays. Per-position results are summarized in Table 5.26. Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran. Sub-threshold signals (1-3 droplets) are tracked separately as they are not interpretable as species presence per ddPCR convention.

Table 5.25: Summary of eDNA samples collected at site 203605.

Site	Stream	Habitat Type	Sample Method	Filter Type	Filter Size (um)	Volume filtered (L)	Site Description
203605_ds_ed1	Tributary To Williston Reservoir	riffle	grab	mce	1	—	Sample collected 15m downstream of the culvert outlet on the right bank

Site	Stream	Habitat Type	Sample Method	Filter Type	Filter Size (um)	Volume filtered (L)	Site Description
203605_us_ed1	Tributary To Williston Reservoir	rifle	instream	mce	1	2	Approximately 45 m upstream of culvert inlet on the left bank

Table 5.26: eDNA detection results at crossing 203605.

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
203605_ds_ed1	M41368	RAIN(455)	BULT(3)*	GRAY(0), SOCK(0)
203605_us_ed1	M41369	RAIN(429)	BULT(1)	GRAY(0), SOCK(0)

Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#). The [interactive Peace eDNA map](#) shows toggleable per-species layers.

Structure Remediation and Cost Estimate

Should restoration/maintenance activities proceed, replacement of the Finlay FSR crossing (203605) with a bridge (15m span) is recommended. At the time of reporting in 2026, the cost of the work is estimated at \$450,000.

Conclusion

The habitat upstream of PSCIS crossing 203605 on Finlay FSR was documented as high value for salmonid spawning and rearing, and the crossing is rated as a high priority for replacement due to the large outlet drop. There is a significant amount of bull trout rearing habitat modelled upstream, and since the road that provides access to lower half of the may be deactivated, this crossing could be a good candidate for replacement.

Table 5.27: Summary of habitat details for PSCIS crossing 203605.

Site	Location	Length Surveyed (m)	Average Channel Width (m)	Average Wetted Width (m)	Average Pool Depth (m)	Average Gradient (%)	Total Cover	Habitat Value
203605	Downstream	250	8.5	5.6	0.2	1.8	moderate	high
203605	Upstream	650	6.7	4.6	3.7	2.9	moderate	high

Conclusion



Figure 5.24: Left: Typical habitat downstream of PSCIS crossing 203605. Right: Typical habitat upstream of PSCIS crossing 203605.

Appendix - Environmental DNA Results

This appendix contains site-level detail for the 2025 environmental DNA (eDNA) sampling program in the Peace project area. Headline per-species detection counts are presented in the main Results chapter; the tables below show which sites detected which species. Field blanks are tracked separately as quality-assurance samples — they test for protocol contamination, not site eDNA. Office blanks (samples filtered at the field accommodation rather than at site) are excluded from these tables because their UTM coordinates are inherited from related sites and do not represent real sampling locations.

The interactive map of these results is linked from the main Results chapter and at the end of this appendix.

Per-site detection results

Format: CODE(max_droplets), with * flagging samples that UNBC reran. Detected = ≥ 4 positive droplets per the standard ddPCR call threshold; Sub-threshold = 1-3 droplets, not interpretable as species presence per ddPCR convention; Not detected = 0 droplets across all runs.

Table 5.28: Site-level eDNA detection results across real environmental samples in the Peace project area. UNBC ID is the lab sample identifier; Pos. control = TRUE indicates the sample was collected at a site where the target species was also confirmed by electrofishing.

Site ID	Stream	UNBC ID	Pos. control	Detected	Sub-threshold	Not detected
125000_ds_ed1	Tributary To Parsnip River	M41347		RAIN(980)	—	BULT(0), GRAY(0), SOCK(0)
125000_us_ed1	Tributary To Parsnip River	M41348	—	RAIN(143)	—	BULT(0), GRAY(0), SOCK(0)
125179_ds_ed1a	Tributary To Missinka River	M41349	—	RAIN(309)	BULT(1)*	GRAY(0), SOCK(0)
125179_us_ed1a	Tributary To Missinka River	M41351	yes	RAIN(383)*	—	BULT(0), GRAY(0), SOCK(0)
125180_ds_ed1	Tributary To Missinka River	M41353		BULT(4), RAIN(330)	—	GRAY(0), SOCK(0)
125180_us_ed1	Tributary To Missinka River	M41354		RAIN(293)	BULT(3)	GRAY(0), SOCK(0)
125194_ds_ed1a	Tributary To Missinka River	M21242		—	BULT(1), RAIN(3)*	—
125231_ds_ed1c	Tributary To Table River	M41356	—	RAIN(205)	—	BULT(0), GRAY(0), SOCK(0)
125231_us_ed1	Tributary To Table River	M41357	—	RAIN(5)	—	BULT(0), GRAY(0), SOCK(0)

Appendix - Environmental DNA Results

Site ID	Stream	UNBC ID	Pos. control	Detected	Sub-threshold	Not detected
125261_ds_ed1	Fern Creek	M41358	—	RAIN(258)	—	BULT(0), GRAY(0), SOCK(0)
125261_us_ed1a	Fern Creek	M41359	—	RAIN(533)	—	BULT(0), GRAY(0), SOCK(0)
125261_us_ed1b	Fern Creek	M41360	—	RAIN(567)	—	BULT(0), GRAY(0), SOCK(0)
198692_ds_ed1b	Tributary To Kerry Lake	M41362	—	RAIN(119)	BULT(1)	GRAY(0), SOCK(0)
198692_ds_ed2	Tributary To Kerry Lake	M41363	—	RAIN(630)	—	BULT(0), GRAY(0), SOCK(0)
198692_us_ed1	Tributary To Kerry Lake	M41364	—	RAIN(235)	—	BULT(0), GRAY(0), SOCK(0)
199663_ds_ed1	Tributary To Parsnip River	M10139	—	BURB(195), RAIN(112)	—	BULT(0), GRAY(0)
199663_us_ed1	Tributary To Parsnip River	M10140	—	RAIN(65)	—	BULT(0), BURB(0), GRAY(0)
203597_ds_ed1b	Tributary To Nation River	M41366		RAIN(173)*	BULT(3)*	GRAY(0), <i>SOCK(0)</i>
203597_us_ed1	Tributary To Nation River	M41367		RAIN(213)*	—	BULT(0), <i>GRAY(0)</i> , SOCK(0)*
203605_ds_ed1	Tributary To Williston Reservoir	M41368	—	RAIN(455)	BULT(3)*	GRAY(0), SOCK(0)
203605_us_ed1	Tributary To Williston Reservoir	M41369	—	RAIN(429)	BULT(1)	GRAY(0), SOCK(0)
24718557_ed1	Missinka River	M41370	—	RAIN(4)	BULT(2), <i>GRAY(1)</i>	SOCK(0)
57690_ds_ed1	Tributary To Wichcika Creek	M41371	—	RAIN(10)	—	BULT(0), GRAY(0), SOCK(0)
57690_us_ed1	Tributary To Wichcika Creek	M41372	—	—	—	BULT(0), GRAY(0), RAIN(0), SOCK(0)
57695_ds_ed1	Tributary To Wichcika Creek	M41373	—	—	—	BULT(0), GRAY(0), RAIN(0), SOCK(0)
57695_us_ed1	Tributary To Wichcika Creek	M21244	—	—	RAIN(1)	BULT(0)
57696_ds_ed1	Tributary To Wichcika Creek	M21245		—	—	BULT(0), RAIN(0)
57696_us_ed1	Tributary To Wichcika Creek	M21246		—	—	BULT(0), RAIN(0)

Field blanks

Field blanks filter distilled water at the field site to test for protocol contamination during collection and processing. **Detections in blanks indicate potential cross-contamination, not species presence at the location** — these rows are presented for QA transparency only, not as eDNA detections.

Retests

Table 5.29: Field blanks collected during the 2025 eDNA program in the Peace project area. Detections in blanks reflect protocol-side contamination, not site eDNA.

Site ID	Stream	UNBC ID	Detected	Sub-threshold	Clean (0 droplets)
125231_ds_ed1a	Tributary To Table River	M41355	—	—	BULT(0), GRAY(0), RAIN(0), SOCK(0)
198692_ds_ed1a	Tributary To Kerry Lake	M41361	RAIN(5)*	—	BULT(0), GRAY(0), SOCK(0)*
203597_ds_ed1a	Tributary To Nation River	M41365	RAIN(4)*	—	BULT(0), GRAY(0), SOCK(0)*

Retests

UNBC reran a subset of samples to confirm borderline results. The table below lists every (site x species) combination that was rerun, including reruns from both real environmental samples and field blanks.

Table 5.30: UNBC retests for the 2025 Peace eDNA samples. Runs is the total number of times the sample x species combination was assayed; Max droplets is the highest positive-droplet count observed across all runs.

Site ID	Stream	Species	Code	Sample type	Runs	Max droplets	Result
125179_ds_ed1a	Tributary To Missinka River	Bull Trout	BULT	environmental	4	1	Sub-threshold (1-3 droplets, NOT a detection)
125179_us_ed1a	Tributary To Missinka River	Rainbow Trout	RAIN	environmental	4	383	Detected (≥ 4 droplets)
125194_ds_ed1a	Tributary To Missinka River	Rainbow Trout	RAIN	environmental	4	3	Sub-threshold (1-3 droplets, NOT a detection)
198692_ds_ed1a	Tributary To Kerry Lake	Bull Trout	BULT	field blank	4	0	No detection (0 droplets)
198692_ds_ed1a	Tributary To Kerry Lake	Arctic Grayling	GRAY	field blank	4	0	No detection (0 droplets)
198692_ds_ed1a	Tributary To Kerry Lake	Rainbow Trout	RAIN	field blank	6	5	Detected (≥ 4 droplets)
198692_ds_ed1a	Tributary To Kerry Lake	Kokanee	SOCK	field blank	6	0	No detection (0 droplets)
203597_ds_ed1a	Tributary To Nation River	Bull Trout	BULT	field blank	4	0	No detection (0 droplets)
203597_ds_ed1a	Tributary To Nation River	Arctic Grayling	GRAY	field blank	4	0	No detection (0 droplets)
203597_ds_ed1a	Tributary To Nation River	Rainbow Trout	RAIN	field blank	4	4	Detected (≥ 4 droplets)

Appendix - Environmental DNA Results

Site ID	Stream	Species	Code	Sample type	Runs	Max droplets	Result
203597_ds_ed1a	Tributary To Nation River	Kokanee	SOCK	field blank	4	0	No detection (0 droplets)
203597_ds_ed1b	Tributary To Nation River	Bull Trout	BULT	environmental	4	3	Sub-threshold (1-3 droplets, NOT a detection)
203597_ds_ed1b	Tributary To Nation River	Arctic Grayling	GRAY	environmental	4	0	No detection (0 droplets)
203597_ds_ed1b	Tributary To Nation River	Rainbow Trout	RAIN	environmental	4	173	Detected (≥ 4 droplets)
203597_ds_ed1b	Tributary To Nation River	Kokanee	SOCK	environmental	4	0	No detection (0 droplets)
203597_us_ed1	Tributary To Nation River	Bull Trout	BULT	environmental	4	0	No detection (0 droplets)
203597_us_ed1	Tributary To Nation River	Arctic Grayling	GRAY	environmental	4	0	No detection (0 droplets)
203597_us_ed1	Tributary To Nation River	Rainbow Trout	RAIN	environmental	4	213	Detected (≥ 4 droplets)
203597_us_ed1	Tributary To Nation River	Kokanee	SOCK	environmental	4	0	No detection (0 droplets)
203605_ds_ed1	Tributary To Williston Reservoir	Bull Trout	BULT	environmental	4	3	Sub-threshold (1-3 droplets, NOT a detection)
24718557_ed1	Missinka River	Bull Trout	BULT	environmental	4	2	Sub-threshold (1-3 droplets, NOT a detection)
24718557_ed1	Missinka River	Arctic Grayling	GRAY	environmental	4	1	Sub-threshold (1-3 droplets, NOT a detection)

Interactive map

The interactive Peace eDNA map shows site-level detection results with toggleable per-species layers, an opt-in sub-threshold-signals layer (off by default), and an opt-in Controls layer for field blanks. See the [interactive Peace eDNA map](#).

Tributary to Missinka River - 125179 - Monitoring Appendix

Effectiveness monitoring was conducted in 2025 at PSCIS crossing 125179 on the Chuchinka-Missinka FSR. Site location, background, prior-year assessment and replacement history, and earlier effectiveness monitoring results are documented in Irvine (2020), Irvine and Winterscheidt (2023), and Irvine and Schick (2025); the 2024 site appendix is available at https://www.newgraphenvironment.com/fish_passage_peace_2024_reporting/trib-to-missinka.html.

Monitoring Form

2025 effectiveness monitoring form results are summarized in Table 5.31.

Table 5.31: Summary of effectiveness monitoring form results for site 125179 in 2025.

Variable	Value
Pscis Crossing Id	125179
Stream Name	Tributary to Missinka River
Road Name	Chuchinka-Missinka FSR
Crossing Subtype	BRIDGE
Span	16
Width	4.8
Assessment Comment	Routine effectiveness evaluation. Structure was replaced several years ago and monitoring is ongoing. Electrofishing with fish over 65mm in length PIT tagged, as well as EDNA sampling was conducted in 2025.
Dewatering Notes	Stream is flowing nicely through the construction footprint with no evidence of aggregation, or any sign of dewatering
Velocity Notes	Velocity within project footprint is slightly faster than naturalized areas upstream and downstream because of the construction from riprap however, overall they do not appear to prevent migration of juveniles larger than approximately 130mm. During higher flow times of the year velocity may be quite higher, however.
Constriction Notes	Channel is constricted within the 4 - 5m upstream and downstream of the bridge with channel widths ranging from 2.5 - 3.4m.
Substrate Notes	Substrate is filling in nicely with gravels and cobble/boulder stone lines with a composition comparable to upstream and downstream within natural areas of the stream.
Riparian Notes	Overall, the riparian footprint disturbed is quite small and shrub growth from the edges of the footprint is beginning to fill in the upstream and downstream edges.
Flow Depth Notes	Flow depth within the construction footprint are comparable to those upstream and downstream in natural areas with some deeper pockets scoured adjacent to stone lines and placed boulders within the channel.
Stability Notes	Structure appears stable with no indication of any issues.

Variable	Value
Revegetation Notes	Adjacent to the bridge there's only a small amount of herbaceous regrowth in between heavily rip wrapped areas. It is expected that as time passes fine substrates will infiltrate the riprap more during high flow events, and that those areas will colonize more with natural herb and shrub species over time.
Cover Notes	Placed riprap boulders and developed stone lines have created areas of bubble curtain and shallow pool cover throughout the construction footprint. Overhead cover from vegetation is extremely minimal still however some willow and other shrub regrowth indicates it will improve overtime.
Maintenance Notes	No maintenance required

Fish Sampling

Electrofishing was conducted downstream and upstream of PSCIS crossing 125179 in 2025. Per-site capture results are in Table [5.32](#); densities are in Table [5.33](#) and Figure [5.25](#).

Table 5.32: Fish sampling site summary for 125179.

site	passes	ef_length_m	ef_width_m	area_m2	enclosure
125179_us_ef1	1	45	2.2	99.0	partial enclosure
125179_ds_ef1	1	35	2.7	94.5	partial enclosure

Table 5.33: Fish sampling density results summary for 125179.

local_name	species_code	life_stage	catch	density_100m2	nfc_pass
125179_ds_ef1	Rainbow Trout	adult	2	2.1	FALSE
125179_ds_ef1	Rainbow Trout	fry	6	6.3	FALSE
125179_ds_ef1	Rainbow Trout	juvenile	2	2.1	FALSE
125179_ds_ef1	Rainbow Trout	parr	3	3.2	FALSE
125179_us_ef1	Rainbow Trout	adult	2	2.0	FALSE
125179_us_ef1	Rainbow Trout	fry	3	3.0	FALSE
125179_us_ef1	Rainbow Trout	juvenile	2	2.0	FALSE
125179_us_ef1	Rainbow Trout	parr	4	4.0	FALSE

* nfc_pass FALSE means fish were captured in final pass indicating more fish of this species/lifestage may have remained in site. Mark-recaptured required to reduce uncertainties.




Figure 5.25: Densities of fish (fish/100m2) captured upstream and downstream of PSCIS crossing 125179 in 2025.

Environmental DNA

eDNA samples were collected upstream and downstream of PSCIS crossing 125179 in 2025 as part of the program-wide eDNA sampling effort. The reference site at PSCIS crossing 125180, located on the same road as documented in Irvine and Schick (2025), was also sampled for eDNA in 2025. Per-position detection results for both sites are presented in Table 5.34. Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran. Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#), and the [interactive Peace eDNA map](#) provides a toggleable per-species view.

Table 5.34: eDNA detection results at crossing 125179.

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
125179_ds_ed1a	M41349	RAIN(309)	BULT(1)*	GRAY(0), SOCK(0)

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
125179_us_ed1a	M41351	RAIN(383)*	—	BULT(0), GRAY(0), SOCK(0)
125180_ds_ed1	M41353	BULT(4), RAIN(330)	—	GRAY(0), SOCK(0)
125180_us_ed1	M41354	RAIN(293)	BULT(3)	GRAY(0), SOCK(0)

Conclusion

At the time of reporting, the structure at PSCIS crossing 125179 appears stable and the site is showing positive signals of recovery. Continued effectiveness monitoring at this site is recommended in future years to extend the multi-year time series of structure performance and fish use.

Tributary to the Table River - 125231 - Monitoring Appendix

Effectiveness monitoring was conducted in 2025 at PSCIS crossing 125231 on the Chuchinka-Table FSR. Site location, background, and the 2023 replacement of the crossing with a clear-span bridge by Canfor (with engineering oversight by DWB Consulting Services Ltd. and 50% funding from FWCP via SERNbc coordination) are documented in Irvine and Winterscheidt (2024); the 2023 site appendix is available at https://www.newgraphenvironment.com/fish_passage_peace_2023_reporting/tributary-to-the-table-river—125231—appendix.html.

Monitoring Form

2025 effectiveness monitoring form results are summarized in Table [5.35](#).

Table 5.35: Summary of effectiveness monitoring form results for site 125231 in 2025.

Variable	Value
Pscis Crossing Id	125231
Stream Name	Tributary to Table River
Road Name	Table FSR
Crossing Subtype	BRIDGE
Span	—
Width	—
Assessment Comment	Overall, the site looks very similar to last year, in good condition with no evidence of any structural issues and the instream portion doing quite well. As there is a large amount of rock rip wrap adjacent to the stream that was not planted the riparian vegetation within the footprint is basically nonexistent. Electrofishing was conducted at sites downstream and upstream of the bridge with fish over 65mm PIT tagged. eDNA samples were collected as well.
Dewatering Notes	No degrading of the channel noted. Good flow throughout construction footprint.
Velocity Notes	Velocities in the construction footprint are similar to upstream and downstream with maturing stone lines and placed boulders, stepping out the gradient to natural conditions
Constriction Notes	Width of channel appears similar to upstream and downstream with room for channel migration.
Substrate Notes	Substrate is very similar to upstream and downstream with well placed riprap and boulders providing scour and creating pool habitat.
Riparian Notes	Basically no riparian regrowth within the construction footprint due to large amount of rip wrap placed.
Flow Depth Notes	Flow depths are acceptable with deep sections adjacent to stone lines and placed boulders.

Variable	Value
Stability Notes	Structure appears stable with no evidence of any movement
Revegetation Notes	Revegetation of the site is generally poor due to the large amount of rock placed in the construction footprint and a lack of planting and seeding.
Cover Notes	Cover is available adjacent to stone lines and placed boulders were scour and some shallow pools have been created. Fish were captured in these areas within the construction footprint, indicating that they are providing suitable habitat for local fish species.
Maintenance Notes	No maintenance required

Fish Sampling

Electrofishing was conducted downstream and upstream of PSCIS crossing 125231 in 2025. Per-site capture results are in Table [5.36](#); densities are in Table [5.37](#) and Figure [5.26](#).

Table 5.36: Fish sampling site summary for 125231.

site	passes	ef_length_m	ef_width_m	area_m2	enclosure
125231_ds_ef1	1	50	2.8	140	partial enclosure
125231_us_ef1	1	50	1.6	80	partial enclosure

Table 5.37: Fish sampling density results summary for 125231.

local_name	species_code	life_stage	catch	density_100m2	nfc_pass
125231_ds_ef1	Rainbow Trout	adult	1	0.7	FALSE
125231_ds_ef1	Rainbow Trout	fry	7	5.0	FALSE
125231_ds_ef1	Rainbow Trout	juvenile	1	0.7	FALSE
125231_ds_ef1	Rainbow Trout	parr	5	3.6	FALSE
125231_us_ef1	Rainbow Trout	adult	1	1.2	FALSE
125231_us_ef1	Rainbow Trout	fry	5	6.2	FALSE
125231_us_ef1	Rainbow Trout	juvenile	3	3.8	FALSE
125231_us_ef1	Rainbow Trout	parr	2	2.5	FALSE
125231_us_ef1	Sucker (General)	adult	1	1.2	FALSE

* nfc_pass FALSE means fish were captured in final pass indicating more fish of this species/lifestage may have remained in site. Mark-recaptured required to reduce uncertainties.




Figure 5.26: Densities of fish (fish/100m2) captured upstream and downstream of PSCIS crossing 125231 in 2025.

Environmental DNA

eDNA samples were collected upstream and downstream of PSCIS crossing 125231 in 2025 as part of the program-wide eDNA sampling effort. Per-position detection results are presented in Table 5.38. Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran. Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#), and the [interactive Peace eDNA map](#) provides a toggleable per-species view.

Table 5.38: eDNA detection results at crossing 125231.

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
125231_ds_ed1c	M41356	RAIN(205)	—	BULT(0), GRAY(0), SOCK(0)
125231_us_ed1	M41357	RAIN(5)	—	BULT(0), GRAY(0), SOCK(0)

Conclusion

At the time of reporting, the structure at PSCIS crossing 125231 appears stable and the site is showing positive signals of recovery. Continued effectiveness monitoring at this site is recommended in future years to extend the multi-year time series of structure performance and fish use.

Tributary to Kerry Lake - 198692 - Monitoring Appendix

Baseline monitoring was conducted in 2025 at PSCIS crossing 198692 ahead of planned remediation, to inform future effectiveness monitoring once the structure is upgraded. Site location, background, and the 2024 habitat confirmation assessment are documented in Irvine and Schick (2025); the 2024 site appendix is available at https://www.newgraphenvironment.com/fish_passage_peace_2024_reporting/trib-to-kerry.html.

Fish Sampling

Electrofishing was conducted downstream and upstream of PSCIS crossing 198692 in 2025 to establish a pre-remediation baseline. Per-site capture results are in Table 5.39; densities are in Table 5.40 and Figure 5.27.

Table 5.39: Fish sampling site summary for 198692.

site	passes	ef_length_m	ef_width_m	area_m2	enclosure
198692_us_ef1	1	16	–	–	partial enclosure
198692_us_ef2	1	12	–	–	open
198692_ds_ef1	1	34	–	–	partial enclosure
198692_ds_ef2	1	4	–	–	open

Table 5.40: Fish sampling density results summary for 198692.

local_name	species_code	life_stage	catch	density_100m2	nfc_pass
198692_ds_ef1	Rainbow Trout	fry	6	–	FALSE
198692_ds_ef1	Rainbow Trout	parr	4	–	FALSE
198692_ds_ef2	Rainbow Trout	fry	1	–	FALSE
198692_ds_ef2	Rainbow Trout	juvenile	4	–	FALSE
198692_ds_ef2	Rainbow Trout	parr	9	–	FALSE
198692_us_ef1	Rainbow Trout	fry	1	–	FALSE
198692_us_ef2	Rainbow Trout	fry	3	–	FALSE
198692_us_ef2	Rainbow Trout	parr	7	–	FALSE

* nfc_pass FALSE means fish were captured in final pass indicating more fish of this species/lifestage may have remained in site. Mark-recaptured required to reduce uncertainties.

Density plot not shown — wetted-area data unavailable for this site.

Figure 5.27: Densities of fish (fish/100m2) captured upstream and downstream of PSCIS crossing 198692 in 2025. Densities are shown where wetted-area data are available; some surveys at this site had no recorded width.

Environmental DNA

eDNA samples were collected upstream and downstream of PSCIS crossing 198692 in 2025 as part of the program-wide eDNA sampling effort. Per-position detection results are presented in Table 5.41. Detections are based on the standard ddPCR call threshold of ≥ 4 positive droplets; format is CODE(max_droplets), with * flagging samples UNBC reran. Site-level detail across all sampled crossings is in [Appendix - Environmental DNA Results \(page 145\)](#), and the [interactive Peace eDNA map](#) provides a toggleable per-species view.

Table 5.41: eDNA detection results at crossing 198692.

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
198692_ds_ed1b	M41362	RAIN(119)	BULT(1)	GRAY(0), SOCK(0)
198692_ds_ed2	M41363	RAIN(630)	—	BULT(0), GRAY(0), SOCK(0)

Conclusion

Site ID	UNBC ID	Detected	Sub-threshold	Not detected
198692_us_ed1	M41364	RAIN(235)	—	BULT(0), GRAY(0), SOCK(0)

Conclusion

Baseline fish and eDNA data were collected at PSCIS crossing 198692 in 2025 to inform future effectiveness monitoring once remediation completes. Resampling at the site once the structure is upgraded will allow direct comparison of fish abundance and habitat conditions before and after the intervention.

Appendix - UAV Imagery

This appendix lists the UAV (drone) imagery products generated for the Peace project area since 2020 and stored as Cloud Optimized GeoTIFFs on AWS. Each row links to a tile-service viewer for in-browser inspection alongside a direct download link to the source raster.

Appendix - Collaborative GIS Layers

This appendix lists background spatial layers and tables loaded to the collaborative GIS project (sern_peace_fwcp_2023) at the time of writing. The project serves as the spatial integration layer for fieldwork — viewable in QGIS in the office and synced to mobile devices for navigation and standardized data collection in the field via Mergin Maps.

References

- Amazon Web Services. 2025. *Aws/Aws-Cli: Universal Command Line Interface for Amazon Web Services*. V. 2.18.15. Released. <https://github.com/aws/aws-cli>.
- BC Species & Ecosystem Explorer. 2020. "Salvelinus Confluentus (Bull Trout)." <https://a100.gov.bc.ca/pub/eswp/reports.do?elcode=AFCHA05020>.
- Beaudry, P. 2014. *Assessment and Assignment of Sensitivity Ratings to Sub-Basins of the Table Watershed in Parsnip Drainage – Ominieca Region*. Contract Number: GS15823011. https://a100.gov.bc.ca/pub/acat/documents/r51634/3_Beaudry_2014_Table_FSW_51634_1480368942206_0367832493.pdf.
- Beaudry, Pierre G. 2013a. *Assessment and Assignment of Sensitivity Ratings to Sub-Basins of the Anzac Watershed in Parsnip Drainage – Ominieca Region*. Contract Number: GS14823018. Report. <http://a100.gov.bc.ca/pub/acat/public/viewReport.do?reportId=51630>.
- Beaudry, Pierre G. 2013b. *Assessment and Assignment of Sensitivity Ratings to Sub-Basins of the Missinka Watershed in Parsnip Drainage – Ominieca Region*. Contract Number: GS14FWH-006. Report. P. Beaudry and Associates Ltd. <http://a100.gov.bc.ca/pub/acat/public/viewReport.do?reportId=51631>.
- Beaudry, Pierre G. 2014. *Assessment and Assignment of Sensitivity Ratings to Sub-Basins of the Hominka Watershed in Parsnip Drainage – Ominieca Region*. Contract Number: GS15823011. Report. <http://a100.gov.bc.ca/pub/acat/public/viewReport.do?reportId=51633>.
- Bell, M. C. 1991. *Fisheries Handbook of Engineering Requirements and Biological Criteria*. https://www.fs.fed.us/biology/nsaec/fishxing/fplibrary/Bell_1991_Fisheries_handbook_of_engineering_requirements_and.pdf.
- Bourne, Christina, Dan Kehler, Yolanda Wiersma, and David Cote. 2011. "Barriers to Fish Passage and Barriers to Fish Passage Assessments: The Impact of Assessment Methods and Assumptions on Barrier Identification and Quantification of Watershed Connectivity." *Aquatic Ecology* 45: 389–403. <https://doi.org/10.1007/s10452-011-9362-z>.
- Bramblett, Robert G., Mason D. Bryant, Brenda E. Wright, and Robert G. White. 2002. "Seasonal Use of Small Tributary and Main-Stem Habitats by Juvenile Steelhead, Coho Salmon, and Dolly Varden in a Southeastern Alaska Drainage Basin." *Transactions of the American Fisheries Society* 131 (3): 498–506. [https://doi.org/10.1577/1548-8659\(2002\)131<0498:SUOSTA>2.0.CO;2](https://doi.org/10.1577/1548-8659(2002)131<0498:SUOSTA>2.0.CO;2).
- Busch, D. Shalin, Mindi Sheer, Kelly Burnett, Paul McElhany, and Tom Cooney. 2011. "Landscape-Level Model to Predict Spawning Habitat For Lower Columbia River Fall Chinook Salmon (*Oncorhynchus Tshawytscha*): Intrinsic Potential Model for Spawning Fall Chinook Salmon." *River Research and Applications* 29 (3): 297–312. <https://doi.org/10.1002/rra.1597>.
- Cayan, Daniel R., Michael D. Dettinger, Susan A. Kammerdiener, Joseph M. Caprio, and David H. Peterson. 2001. "Changes in the Onset of Spring in the Western United States." *Bulletin of the American Meteorological Society* 82 (3): 399–415. [https://doi.org/10.1175/1520-0477\(2001\)082<0399:citoos>2.3.co;2](https://doi.org/10.1175/1520-0477(2001)082<0399:citoos>2.3.co;2).
- Clarkin, K, A Connor, M Furniss, et al. 2005. *National Inventory and Assessment Procedure For Identifying Barriers to Aquatic Organism Passage at Road-Stream Crossings*. United States Department of Agriculture, Forest Service, National Technology; Development Program. <https://www.fs.fed.us/biology/nsaec/fishxing/publications/PDFs/NIAP.pdf>.
- cogeotiff. (2018) 2025. *Cogeotiff/Rio-Cogeo*. March 9; Cogeotiff, released March 19. <https://github.com/cogeotiff/rio-cogeo>.
- Cooney, Thomas, and Damon Holzer. 2006. *Appendix C: Interior Columbia Basin Stream Type Chinook Salmon and Steelhead Populations: Habitat Intrinsic Potential Analysis*. 21.

References

- COSEWIC. 2012. *COSEWIC Assessment and Status Report on the Bull Trout Salvelinus Confluentus in Canada*. Report. Committee on the Status of Endangered Wildlife in Canada. https://www.sararegistry.gc.ca/virtual_sara/files/cosewic/sr_omble_tete_plat_bull_trout_1113_e.pdf.
- Cote, David, P Frampton, M Langdon, and R Collier. 2005. *Fish Passage and Stream Habitat Restoration in Terra Nova National Park Highway Culverts*.
- Development Seed. (2019) 2025. *Developmentseed/Titiler*. V. 0.21.1. June 28; Development Seed, released. <https://github.com/developmentseed/titiler>.
- Diebel, M. W., M. Fedora, S. Cogswell, and J. R. O'Hanley. 2015. "Effects of Road Crossings on Habitat Connectivity for Stream-Resident Fish: STREAM-RESIDENT FISH HABITAT CONNECTIVITY." *River Research and Applications* 31 (10): 1251–61. <https://doi.org/10.1002/rra.2822>.
- Fernandes, I. M., Y. F. Bastos, D. S. Barreto, L. S. Lourenço, and J. M. Penha. 2017. "The Efficacy of Clove Oil as an Anaesthetic and in Euthanasia Procedure for Small-Sized Tropical Fishes." *Brazilian Journal of Biology = Revista Brasileira De Biologia* 77 (3): 444–50. <https://doi.org/10.1590/1519-6984.15015>.
- Finnegan, Noah J., Gerard Roe, David R. Montgomery, and Bernard Hallet. 2005. "Controls on the Channel Width of Rivers: Implications for Modeling Fluvial Incision of Bedrock." *Geology* 33 (3): 229–32. <https://doi.org/10.1130/G21171.1>.
- Fish and Wildlife Compensation Program. 2020. *Peace Region Rivers, Lakes, & Reservoirs Action Plan*. <https://fwcp.ca/app/uploads/2020/08/Action-Plan-Peace-Region-Rivers-Lakes-Reservoirs-Aug-11-2020.pdf>.
- Fish Passage Technical Working Group. 2011. *A Checklist for Fish Habitat Confirmation Prior to the Rehabilitation Fo a Stream Crossing*. <https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/land-based-investment/forests-for-tomorrow/checklist-for-fish-habitat-confirmation-201112.pdf>.
- FLNRORD. 2020a. "Digital Road Atlas (DRA) - Master Partially-Attributed Roads - Data Catalogue." Published by Ministry of Forests, Lands, Natural Resource Operations and Rural Development. <https://catalogue.data.gov.bc.ca/dataset/digital-road-atlas-dra-master-partially-attributed-roads>.
- FLNRORD. 2020b. "Forest Tenure Road Section Lines - Data Catalogue." Published by Ministry of Forests, Lands, Natural Resource Operations and Rural Development (FLNRORD). <https://catalogue.data.gov.bc.ca/dataset/forest-tenure-road-section-lines>.
- Forest and Range Practices Act. 2018. *Order - Fisheries Sensitive Watershed Prince George Forest District*. <http://www.env.gov.bc.ca/wld/frpa/fsw/approved.html>.
- Forest Investment Account. 2003. *Guidelines for in-Stream and Off-Channel Routine Effectiveness Evaluation*. British Columbia Ministry of Forests. http://env.gov.bc.ca/wld/documents/fia_docs/ree_guidelines.pdf.
- Gilbert, D. E., J. E. Morris, A. R. Kaveney, and J. D. Dery. 2022. "Sub-Hourly Water Temperature Data Collected Across the Nechako Watershed, 2019-2021 | Elsevier Enhanced Reader." <https://doi.org/10.1016/j.dib.2022.108425>.
- Hagen, John, Ray Pillipow, and Nikolaus Gantner. 2018. *Trend in Abundance of Arctic Grayling (Thymallus Arcticus) in Index Sites of the Parsnip River Watershed, 1995-2018*. 37.
- Hagen, John, Ian Spendlow, and Ray Pillipow. 2020. *Critical Spawning Habitats and Abundance of Bull Trout in the Williston Reservoir Watershed, 2019*. 57.
- Hagen, John, and Mike Stamford. 2021. *Arctic Grayling (Thymallus Arcticus) Abundance and Trend in the Parsnip River Watershed, 1995-2020*. https://a100.gov.bc.ca/pub/acat/documents/r59157/PEA_F21_F_3203_1622500728936_967641590A.pdf.

Hagen, John, and Mike Stamford. 2023. *Parsnip Arctic Grayling Abundance and Critical Habitats 2018-2022 Final Report*.

Hagen, John, and Susanne Weber. 2019. *Limiting Factors, Enhancement Potential, Critical Habitats, and Conservation Status for Bull Trout of the Williston Reservoir Watershed: Information Synthesis and Recommended Monitoring Framework*. Report. <http://fwcp.ca/app/uploads/2019/08/Bull-Trout-Info-Synthesis-Monitoring-Framework- FINAL29Aug2019.pdf>.

Hagen, John, Susanne Williamson, Mike Stamford, and Ray Phillipow. 2015. *Critical Habitats for Bull Trout and Arctic Grayling Within the Parsnip River and Pack River Watersheds*.

Hall, Jason E., Diane M. Holzer, and Timothy J. Beechie. 2007. "Predicting River Floodplain and Lateral Channel Migration for Salmon Habitat Conservation." *Journal of the American Water Resources Association* 43 (3): 786–97. <https://doi.org/10.1111/j.1752-1688.2007.00063.x>.

Hansen, James, Makiko Sato, and Reto Ruedy. 2012. "Perception of Climate Change." *Proceedings of the National Academy of Sciences* 109 (37). <https://doi.org/10.1073/pnas.1205276109>.

Hill, N. H., J. L. Thorley, and A. Irvine. 2024. *Spatial Stream Network Analysis of Nechako Watershed Stream Temperatures 2022b*. Poisson Consulting Analysis Appendix. <https://www.poissonconsulting.ca/f/1295467017>.

Hirst, S. M. 1991. *Impacts of the Operations of Existing Hydroelectric Developments on Fishery Resources in British Columbia*. <https://waves-vagues.dfo-mpo.gc.ca/Library/124832main.pdf>.

"History Who We Are." 2023. McLeod Lake Indian Band. <https://www.mlib.ca/about/History/>.

Hobbs, Jared, Caren C. Helbing, and Caren Goldberg. 2025. *Environmental DNA Protocol for Freshwater Aquatic Ecosystems: Version 3.0*. BC Ministry of Environment, Ecosystems Branch.

Irvine. (2021) 2022. *Bulkley River and Morice River Watershed Groups Fish Passage Restoration Planning 2021*. https://github.com/NewGraphEnvironment/fish_passage_skeena_2021_reporting.

Irvine. 2022. *PEA-F22-F-3577-DCA Restoring Fish Passage in the Peace Region*. https://newgraphenvironment.github.io/fish_passage_parsnip_2021_reporting/.

Irvine. (2025) 2025. *NewGraphEnvironment/Stac_uav_bc*. January 29, released. https://github.com/NewGraphEnvironment/stac_uav_bc.

Irvine, Allan. 2020. *Parsnip River Watershed – Fish Habitat Confirmations (PEA-F20-F-2967)*. https://newgraphenvironment.github.io/Parsnip_Fish_Passage/.

Irvine, and L Schick. 2025. *Restoring Fish Passage in the Peace Region - 2024*. https://newgraphenvironment.github.io/fish_passage_peace_2024_reporting/.

Irvine, and M Winterscheidt. 2023. *Restoring Fish Passage in the Peace Region - 2022 - PEA-F23-F-3761-DCA*. https://newgraphenvironment.github.io/fish_passage_peace_2022_reporting/.

Irvine, and M Winterscheidt. 2024. *Restoring Fish Passage in the Peace Region - 2023*. https://newgraphenvironment.github.io/fish_passage_peace_2023_reporting/.

Kang, Do Hyuk, Huilin Gao, Xiaogang Shi, Siraj ul Islam, and Stephen J. Déry. 2016. "Impacts of a Rapidly Declining Mountain Snowpack on Streamflow Timing in Canada's Fraser River Basin." *Scientific Reports* 6 (1). <https://doi.org/10.1038/srep19299>.

Karl, Thomas R., Richard W. Knight, Kevin P. Gallo, et al. 1993. "A New Perspective on Recent Global Warming: Asymmetric Trends of Daily Maximum and Minimum Temperature." *Bulletin of the American Meteorological Society* 74 (6): 1007–23. [https://doi.org/10.1175/1520-0477\(1993\)074<1007:ANPORG>2.0.CO;2](https://doi.org/10.1175/1520-0477(1993)074<1007:ANPORG>2.0.CO;2).

Kemp, P. S., and J. R. O'Hanley. 2010. "Procedures for Evaluating and Prioritising the Removal of Fish Passage Barriers: A Synthesis: EVALUATION OF FISH PASSAGE BARRIERS." *Fisheries Management and Ecology*, no–. <https://doi.org/10.1111/j.1365-2400.2010.00751.x>.

Kirsch, J M, Joseph D Buckwalter, and Daniel J Reed. 2014. *Fish Inventory and Anadromous Cataloging in the Susitna River, Matanuska River, and Knik River Basins, 2003 and 2011*. 149.

References

- Knowles, Noah, Michael D. Dettinger, and Daniel R. Cayan. 2006. "Trends in Snowfall Versus Rainfall in the Western United States." *Journal of Climate* 19 (18): 4545–59. <https://doi.org/10.1175/jcli3850.1>.
- Martins, E, B O'Connor, J Bottoms, et al. 2022. *Spatial Ecology of Arctic Grayling in the Parsnip Core Area*. https://a100.gov.bc.ca/pub/acat/documents/r59889/PEA_F22_F_3388_1664380292568_8FCE11DCCD.pdf.
- MoE. 2011. *Field Assessment for Determining Fish Passage Status of Closed Bottom Structures*. BC Ministry of Environment (MoE). <https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/land-based-investment/forests-for-tomorrow/field-assessment-for-determining-fish-passage-status-of-cbs.pdf>.
- MoE. 2024a. "Known BC Fish Observations and BC Fish Distributions." <https://catalogue.data.gov.bc.ca/dataset/known-bc-fish-observations-and-bc-fish-distributions>.
- MoE. 2024b. "Provincial Obstacles to Fish Passage - Data Catalogue." <https://catalogue.data.gov.bc.ca/dataset/provincial-obstacles-to-fish-passage>.
- Mote, Philip W., Alan F. Hamlet, Martyn P. Clark, and Dennis P. Lettenmaier. 2005. "DECLINING MOUNTAIN SNOWPACK IN WESTERN NORTH AMERICA*." *Bulletin of the American Meteorological Society* 86 (1): 39–50. <https://doi.org/10.1175/bams-86-1-39>.
- Mote, Philip W., Sihan Li, Dennis P. Lettenmaier, Mu Xiao, and Ruth Engel. 2018. "Dramatic Declines in Snowpack in the Western US." *Npj Climate and Atmospheric Science* 1 (1). <https://doi.org/10.1038/s41612-018-0012-1>.
- Muñoz Sabater, Joaquín. 2019. *ERA5-Land Hourly Data from 1950 to Present*. Copernicus Climate Change Service (C3S) Climate Data Store (CDS). <https://doi.org/10.24381/cds.e2161bac>.
- Murray, Brent W., and Barry Booth. 2023. *Report on the 2023 eDNA Survey of Study Creeks in Nechako Watershed from Prince George to the Upper Fraser*. University of Northern British Columbia, Department of Ecosystem Science and Management. <https://www.newssociety.org/uploads/images/News%20report%20Images/e-DNA%20summary-%20UNBC-NEWSS%202023.pdf>.
- Nagel, David E., John M. Buffington, Sharon L. Parkes, Seth Wenger, and John R. Goode. 2014. *A Landscape Scale Valley Confinement Algorithm: Delineating Unconfined Valley Bottoms for Geomorphic, Aquatic, and Riparian Applications*. Gen. Tech. Rep. RMRS-GTR-321. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. <https://www.fs.usda.gov/rmrs/publications/landscape-scale-valley-confinement-algorithm-delineating-unconfined-valley-bottoms>.
- Norris, Simon. (2018) 2024. *Smnorris/Bcfishobs*. May 11, released. <https://github.com/smnorris/bcfishobs>.
- Norris, Simon. (2020) 2024. *Smnorris/Bcfishpass*. November 7, released January 12. <https://github.com/smnorris/bcfishpass>.
- OpenDroneMap Authors. (2014) 2025. *OpenDroneMap/ODM*. V. 3.5.4. September 25; OpenDroneMap, released. <https://github.com/OpenDroneMap/ODM>.
- Pacific Climate Impacts Consortium. n.d. "Gridded Hydrologic Model Output." Accessed March 26, 2025. https://data.pacificclimate.org/portal/hydro_model_out/map/.
- Pearce, T, S Lieske, J Morgan, and A Salonas. 2019. *First Nations Information Gathering on Kokanee, Bull Trout and Arctic Grayling: : Report for the Tse'khene Nation (McLeod Lake Indian Band)*. https://a100.gov.bc.ca/pub/acat/documents/r57248/PEA-F19-F-2866-DCA-5-Final_Report-Arctic_North-Ko_1600902816644_0895426641.pdf.
- Pepin, N., R. S. Bradley, H. F. Diaz, et al. 2015. "Elevation-Dependent Warming in Mountain Regions of the World." *Nature Climate Change* 5 (5): 424–30. <https://doi.org/10.1038/nclimate2563>.

Porter, Marc, Darcy Pickard, Katherine Wieckowski, and Katy Bryan. 2008. *Developing Fish Habitat Models for Broad-Scale Forest Planning in the Southern Interior of B.C.* ESSA Technologies Ltd. and B.C. Ministry of the Environment (MOE) for B.C. Forest Science Program. https://www.for.gov.bc.ca/hfd/library/FIA/2008/FSP_Y081231.pdf.

PySTAC Authors. (2017) 2025. *Stac-Utills/PyStac*. V. 1.12.2. November 8; Stac-utills, released. <https://github.com/stac-utills/pystac>.

Radianteearth. 2024. "STAC: SpatioTemporal Asset Catalogs." <https://stacspect.org/en/>.

Resource Inventory Standards Committee. 2001. *Reconnaissance (1:20 000) Fish and Fish Habitat Inventory Standards and Procedures*. <https://www2.gov.bc.ca/assets/gov/environment/natural-resource-stewardship/nr-laws-policy/risc/recce2c.pdf>.

Roberge, M, J M B Hume, C K Minns, and T Slaney. 2002. *Life History Characteristics of Freshwater Fishes Occurring in British Columbia and the Yukon, with Major Emphasis on Stream Habitat Characteristics*. 262.

Rosenfeld, Jordan, Marc Porter, and Eric Parkinson. 2000. *Habitat Factors Affecting the Abundance and Distribution of Juvenile Cutthroat Trout (Oncorhynchus Clarki) and Coho Salmon (Oncorhynchus Kisutch)*. 57: 9.

Shrimpton, J. M., S. L. Roberts, and A. D. Clarke. 2012. *Genetic Analysis of Arctic Grayling Population Structure in the Williston Watershed*. http://a100.gov.bc.ca/appsd/aca/documents/r38246/Report_311_GR_genetics_2012_1379090176074_c3e946726980918f9466d47b59978ad28ded2b969041dd7fe53b83cfe3011493.pdf.

Simoes, Rolf, Felipe Souza, Matheus Zaglia, Gilberto Ribeiro de Queiroz, Rafael dos Santos, and Karine Ferreira. 2021. "Rstac: An R Package to Access Spatiotemporal Asset Catalog Satellite Imagery." *2021 IEEE International Geoscience and Remote Sensing Symposium (IGARSS)*, 7674–77. <https://doi.org/10.1109/IGARSS47720.2021.9553518>.

Slaney, P A, and D Zaldokas. 1997. *Fish Habitat Rehabilitation Procedures*. https://www.for.gov.bc.ca/hfd/library/FFIP/Slaney_PA1997_A.pdf.

Sloat, Matthew R., Gordon H. Reeves, and Kelly R. Christiansen. 2017. "Stream Network Geomorphology Mediates Predicted Vulnerability of Anadromous Fish Habitat to Hydrologic Change in Southeast Alaska." *Global Change Biology* 23 (2): 604–20. <https://doi.org/10.1111/gcb.13466>.

Species Registry Canada. 2020. "Bull Trout (Salvelinus Confluentus), Western Arctic Populations - Species Search - Species at Risk Registry." <https://species-registry.canada.ca/index-en.html#/species/1202-869>.

stac-utills. 2025. *Stac-Utills/Stac-Fastapi-Pgstac: PostgreSQL Backend for Stac-Fastapi Using Pgstac* (<https://github.com/stac-utills/pgstac>). V. 4.0.2. Released. <https://github.com/stac-utills/stac-fastapi-pgstac>.

Stamford, Mike, John Hagen, and Susanne Williamson. 2017. *FWCP Arctic Grayling Synthesis Report*. 148. http://fwcp.ca/app/uploads/2017/07/FWCP_Grayling_Synthesis_Final.pdf.

Stewart, D B, N J Mochnacz, J D Reist, T J Carmichael, and C D Sawatzky. 2007. "Fish Life History and Habitat Use in the Northwest Territories: Arctic Grayling (Thymallus Arcticus)." *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 2797, 64. https://www.researchgate.net/publication/255580930_Fish_life_history_and_habitat_use_in_the_Northwest_Territories_Arctic_grayling_Thymallus_arcticus.

Stewart, Iris T., Daniel R. Cayan, and Michael D. Dettinger. 2005. "Changes Toward Earlier Streamflow Timing Across Western North America." *Journal of Climate* 18 (8): 1136–55. <https://doi.org/10.1175/jcli3321.1>.

Swales, S., and C. D. Levings. 1989. "Role of Off-Channel Ponds in the Life Cycle of Coho Salmon (*Oncorhynchus Kisutch*) and Other Juvenile Salmonids in the Coldwater River, British

References

- Columbia.” *Canadian Journal of Fisheries and Aquatic Sciences* 46 (2): 232–42. <https://doi.org/10.1139/f89-032>.
- Thompson, Richard. 2013. *Assessing Fish Passage at Culverts – the Method, Its Metrics and Preliminary Findings from over 4,000 Assessments*. https://www2.gov.bc.ca/assets/gov/environment/plants-animals-and-ecosystems/fish-fish-habitat/fish-passage/assessing_fish_passage_at_culverts.pdf.
- Thorley, J. L., S. Norris, and A. Irvine. 2021. *Channel Width 2021b*. <https://www.poissonconsulting.ca/f/859859031>.
- Thorley, J, and A Irvine. 2021. “Channel Width 2021.” <https://www.poissonconsulting.ca/f/1792764180>.
- Toffolon, Marco, and Sebastiano Piccolroaz. 2015. “A Hybrid Model for River Water Temperature as a Function of Air Temperature and Discharge.” *Environmental Research Letters* 10 (11): 114011. <https://doi.org/10.1088/1748-9326/10/11/114011>.
- Washington Department of Fish & Wildlife. 2009. *Fish Passage Barrier and Surface Water Diversion Screening Assessment and Prioritization Manual*. Washington Department of Fish; Wildlife. Olympia, Washington. <https://wdfw.wa.gov/sites/default/files/publications/00061/wdfw00061.pdf>.
- Williamson, S. A, and J. T. Zimmerman. 2005. *Region 7a, Omineca Arctic Grayling (Thymallus Arcticus): Data Consolidation Review and Gap Analysis*. B.C. Ministry of Water, Land, and Air Protection.
- Woll, Christine, David Albert, and Diane Whited. 2017. *Salmon Ecological Systems*. The Nature Conservancy. https://www.conservationgateway.org/ConservationByGeography/NorthAmerica/UnitedStates/alaska/sw/cpa/Documents/TNC_A_Salmon_Ecological_Systems_Model_Nushagak_Kvichak.pdf.

Note that this reference was not actually cited in the body of the report but rather in the index.Rmd file yam! headerj under the no_cite key.

Session Info

Information about the computing environment is important for reproducibility. A summary of the computing environment is saved to `session_info.csv`, which can be viewed and downloaded from [here](#).

– Session info

```
## setting value
## version R version 4.5.2 (2025-10-31)
## os      macOS Tahoe 26.2
## system  aarch64, darwin20
## ui      X11
## language (EN)
## collate en_US.UTF-8
## ctype   en_US.UTF-8
## tz      America/Vancouver
## date    2026-05-14
## pandoc  3.9.0.2 @ /opt/homebrew/bin/ (via rmarkdown)
## quarto  1.4.553 @ /usr/local/bin/quarto
```

– Packages

## package	* version	date (UTC)	lib	source
## arrow	22.0.0.1	2025-12-23	[2]	CRAN (R 4.5.2)
## askpass	1.2.1	2024-10-04	[1]	CRAN (R 4.5.0)
## assertthat	0.2.1	2019-03-21	[2]	CRAN (R 4.5.0)
## base64enc	0.1-6	2026-02-02	[1]	CRAN (R 4.5.2)
## bit	4.6.0	2025-03-06	[1]	CRAN (R 4.5.0)
## bit64	4.8.0	2026-04-21	[1]	CRAN (R 4.5.2)
## blob	1.3.0	2026-01-14	[1]	CRAN (R 4.5.2)
## bookdown	* 0.46	2025-12-05	[2]	CRAN (R 4.5.2)
## boot	1.3-32	2025-08-29	[2]	CRAN (R 4.5.2)
## bslib	0.10.0	2026-01-26	[1]	CRAN (R 4.5.2)
## cachem	1.1.0	2024-05-16	[1]	CRAN (R 4.5.0)
## cd	* 0.3.0	2026-05-12	[1]	Github
(NewGraphEnvironment/cd@a3e4733)				
## chk	0.10.0	2025-01-24	[1]	CRAN (R 4.5.0)
## class	7.3-23	2025-01-01	[2]	CRAN (R 4.5.2)
## classInt	0.4-11	2025-01-08	[1]	CRAN (R 4.5.0)
## cli	3.6.6	2026-04-09	[1]	CRAN (R 4.5.2)
## codetools	0.2-20	2024-03-31	[2]	CRAN (R 4.5.2)
## crayon	1.5.3	2024-06-20	[1]	CRAN (R 4.5.0)
## crosstalk	1.2.2	2025-08-26	[1]	CRAN (R 4.5.0)
## curl	7.1.0	2026-04-22	[1]	CRAN (R 4.5.2)
## data.table	1.18.2.1	2026-01-27	[1]	CRAN (R 4.5.2)

Session Info

```
## DBI 1.3.0 2026-02-25 [1] CRAN (R 4.5.2)
## dbplyr 2.5.2 2026-02-13 [1] CRAN (R 4.5.2)
## desc 1.4.3 2023-12-10 [2] CRAN (R 4.5.0)
## devtools 2.4.6 2025-10-03 [2] CRAN (R 4.5.0)
## digest 0.6.39 2025-11-19 [1] CRAN (R 4.5.2)
## doParallel 1.0.17 2022-02-07 [2] CRAN (R 4.5.0)
## dplyr * 1.2.1 2026-04-03 [1] CRAN (R 4.5.2)
## DT 0.34.0 2025-09-02 [2] CRAN (R 4.5.0)
## e1071 1.7-17 2025-12-18 [1] CRAN (R 4.5.2)
## elevatr 0.99.1 2025-09-10 [2] CRAN (R 4.5.0)
## ellipsis 0.3.2 2021-04-29 [2] CRAN (R 4.5.0)
## english 1.2-6 2021-08-21 [2] CRAN (R 4.5.0)
## evaluate 1.0.5 2025-08-27 [1] CRAN (R 4.5.0)
## farver 2.1.2 2024-05-13 [1] CRAN (R 4.5.0)
## fasstr 0.5.3 2024-09-27 [2] CRAN (R 4.5.0)
## fastmap 1.2.0 2024-05-15 [1] CRAN (R 4.5.0)
## fishbc * 0.2.1.9000 2026-01-15 [2] Github (lucy-
schick/fishbc@0607ffd)
## forcats * 1.0.1 2025-09-25 [2] CRAN (R 4.5.0)
## foreach 1.5.2 2022-02-02 [2] CRAN (R 4.5.0)
## fpr * 1.2.0 2026-02-03 [2] Github
(NewGraphEnvironment/fpr@2817533)
## fs 2.1.0 2026-04-18 [1] CRAN (R 4.5.2)
## generics 0.1.4 2025-05-09 [1] CRAN (R 4.5.0)
## ggplot2 * 4.0.3 2026-04-22 [1] CRAN (R 4.5.2)
## ggrepel 0.9.7 2026-02-25 [2] CRAN (R 4.5.2)
## gh 1.5.0 2025-05-26 [2] CRAN (R 4.5.0)
## glue 1.8.1 2026-04-17 [1] CRAN (R 4.5.2)
## gtable 0.3.6 2024-10-25 [1] CRAN (R 4.5.0)
## here 1.0.2 2025-09-15 [2] CRAN (R 4.5.0)
## hms 1.1.4 2025-10-17 [1] CRAN (R 4.5.0)
## htmltools 0.5.9 2025-12-04 [1] CRAN (R 4.5.2)
## htmlwidgets 1.6.4 2023-12-06 [1] CRAN (R 4.5.0)
## http 1.4.8 2026-02-13 [1] CRAN (R 4.5.2)
## iterators 1.0.14 2022-02-05 [2] CRAN (R 4.5.0)
## janitor 2.2.1 2024-12-22 [1] CRAN (R 4.5.0)
## jquerylib 0.1.4 2021-04-26 [1] CRAN (R 4.5.0)
## jsonlite 2.0.0 2025-03-27 [1] CRAN (R 4.5.0)
## kableExtra * 1.4.0 2024-01-24 [2] CRAN (R 4.5.0)
## Kendall 2.2.2 2025-12-22 [2] CRAN (R 4.5.2)
## KernSmooth 2.23-26 2025-01-01 [2] CRAN (R 4.5.2)
## knitr * 1.51 2025-12-20 [1] CRAN (R 4.5.2)
## labeling 0.4.3 2023-08-29 [1] CRAN (R 4.5.0)
## lattice 0.22-9 2026-02-09 [2] CRAN (R 4.5.2)
## leafem * 0.2.5 2025-08-28 [2] CRAN (R 4.5.0)
## leaflet * 2.2.3 2025-09-04 [1] CRAN (R 4.5.0)
## lifecycle 1.0.5 2026-01-08 [1] CRAN (R 4.5.2)
```

```

## lubridate      * 1.9.5      2026-02-04 [1] CRAN (R 4.5.2)
## magick         2.9.0       2025-09-08 [2] CRAN (R 4.5.0)
## magrittr       2.0.5       2026-04-04 [1] CRAN (R 4.5.2)
## Matrix         1.7-4       2025-08-28 [2] CRAN (R 4.5.2)
## memoise        2.0.1       2021-11-26 [1] CRAN (R 4.5.0)
## mgcv           1.9-3       2025-04-04 [2] CRAN (R 4.5.2)
## ngr            * 0.0.1      2026-01-22 [2] Github
(newgraphenvironment/ngr@22c9714)
## nlme           3.1-168     2025-03-31 [2] CRAN (R 4.5.2)
## otel           0.2.0       2025-08-29 [1] CRAN (R 4.5.0)
## pagedown       * 0.23      2025-08-20 [2] CRAN (R 4.5.0)
## pak            0.9.5       2026-04-27 [1] CRAN (R 4.5.2)
## pdftools       * 3.7.0      2026-01-30 [2] CRAN (R 4.5.2)
## pillar         1.11.1      2025-09-17 [1] CRAN (R 4.5.0)
## pkgbuild       1.4.8       2025-05-26 [2] CRAN (R 4.5.0)
## pkgconfig      2.0.3       2019-09-22 [1] CRAN (R 4.5.0)
## pkgload        1.5.0       2026-02-03 [2] CRAN (R 4.5.2)
## plyr           1.8.9       2023-10-02 [1] CRAN (R 4.5.0)
## png            0.1-9       2026-03-15 [1] CRAN (R 4.5.2)
## prettyunits    1.2.0       2023-09-24 [1] CRAN (R 4.5.0)
## processx       3.9.0       2026-04-22 [1] CRAN (R 4.5.2)
## progress       1.2.3       2023-12-06 [2] CRAN (R 4.5.0)
## progressr      0.18.0      2025-11-06 [2] CRAN (R 4.5.0)
## proxy          0.4-29      2025-12-29 [1] CRAN (R 4.5.2)
## purrr          * 1.2.2      2026-04-10 [1] CRAN (R 4.5.2)
## qpdf           1.4.1       2025-07-02 [2] CRAN (R 4.5.0)
## R6             2.6.1       2025-02-15 [1] CRAN (R 4.5.0)
## rappdirs       0.3.4       2026-01-17 [1] CRAN (R 4.5.2)
## raster         3.6-32      2025-03-28 [1] CRAN (R 4.5.0)
## rayshader      0.37.3      2024-02-21 [2] CRAN (R 4.5.0)
## rbbt           * 0.0.0.9000 2026-01-15 [2] Github
(paleolimbot/rbbt@212680c)
## RColorBrewer   1.1-3       2022-04-03 [1] CRAN (R 4.5.0)
## Rcpp           1.1.1-1.1   2026-04-24 [1] CRAN (R 4.5.2)
## RcppRoll       0.3.1       2024-07-07 [2] CRAN (R 4.5.0)
## readr          * 2.2.0      2026-02-19 [1] CRAN (R 4.5.2)
## readwritesqlite * 0.2.0.9021 2026-01-15 [2] Github
(poissonconsulting/readwritesqlite@ccdb297)
## remotes        2.5.0       2024-03-17 [2] CRAN (R 4.5.0)
## rgl            1.3.31      2025-11-14 [2] CRAN (R 4.5.2)
## rlang          1.2.0       2026-04-06 [1] CRAN (R 4.5.2)
## rmarkdown      * 2.31       2026-03-26 [1] CRAN (R 4.5.2)
## RPostgres      * 1.4.10     2026-02-16 [1] CRAN (R 4.5.2)
## rprojroot      2.1.1       2025-08-26 [2] CRAN (R 4.5.0)
## RSQLite        3.52.0      2026-05-10 [1] CRAN (R 4.5.2)
## rstudioapi     0.18.0      2026-01-16 [2] CRAN (R 4.5.2)
## rvest          1.0.5       2025-08-29 [2] CRAN (R 4.5.0)

```

Session Info

```
## s2 1.1.9 2025-05-23 [1] CRAN (R 4.5.0)
## S7 0.2.2 2026-04-22 [1] CRAN (R 4.5.2)
## sass 0.4.10 2025-04-11 [1] CRAN (R 4.5.0)
## scales 1.4.0 2025-04-24 [1] CRAN (R 4.5.0)
## sessioninfo 1.2.3 2025-02-05 [2] CRAN (R 4.5.0)
## sf * 1.1-0 2026-02-24 [1] CRAN (R 4.5.2)
## snakecase 0.11.1 2023-08-27 [1] CRAN (R 4.5.0)
## sp 2.2-1 2026-02-13 [1] CRAN (R 4.5.2)
## staticimports * 0.0.0.9001 2026-01-15 [2] Github
(newgraphenvironment/staticimports@19cd73c)
## stringi 1.8.7 2025-03-27 [1] CRAN (R 4.5.0)
## stringr * 1.6.0 2025-11-04 [1] CRAN (R 4.5.0)
## svglite 2.2.2 2025-10-21 [2] CRAN (R 4.5.0)
## systemfonts 1.3.1 2025-10-01 [2] CRAN (R 4.5.0)
## terra * 1.9-11 2026-03-26 [1] CRAN (R 4.5.2)
## textshaping 1.0.4 2025-10-10 [2] CRAN (R 4.5.0)
## tibble * 3.3.1 2026-01-11 [1] CRAN (R 4.5.2)
## tidyhydat 0.7.2 2025-10-23 [2] CRAN (R 4.5.0)
## tidyr * 1.3.2 2025-12-19 [1] CRAN (R 4.5.2)
## tidyselect 1.2.1 2024-03-11 [1] CRAN (R 4.5.0)
## tidyterra * 1.1.0 2026-03-11 [2] CRAN (R 4.5.2)
## tidyverse * 2.0.0 2023-02-22 [2] CRAN (R 4.5.0)
## tidycl 1.0.10 2026-01-15 [2] Github
(nacnudus/tidycl@7e2f7be7)
## timechange 0.4.0 2026-01-29 [1] CRAN (R 4.5.2)
## tzdb 0.5.0 2025-03-15 [1] CRAN (R 4.5.0)
## units 1.0-1 2026-03-11 [1] CRAN (R 4.5.2)
## usethis 3.2.1 2025-09-06 [2] CRAN (R 4.5.0)
## vctrs 0.7.3 2026-04-11 [1] CRAN (R 4.5.2)
## viridisLite 0.4.3 2026-02-04 [1] CRAN (R 4.5.2)
## vroom 1.7.1 2026-03-31 [1] CRAN (R 4.5.2)
## withr 3.0.2 2024-10-28 [1] CRAN (R 4.5.0)
## wk 0.9.5 2025-12-18 [1] CRAN (R 4.5.2)
## xfun 0.57 2026-03-20 [1] CRAN (R 4.5.2)
## xml2 1.5.2 2026-01-17 [1] CRAN (R 4.5.2)
## yaml 2.3.12 2025-12-10 [1] CRAN (R 4.5.2)
## zyp 0.11-1 2023-03-22 [2] CRAN (R 4.5.0)
##
## [1] /Users/airvine/Library/R/arm64/4.5/library
## [2] /Library/Frameworks/R.framework/Versions/4.5-
arm64/Resources/library
## * — Packages attached to the search path.
##
##
```

Attachment - Phase 1 Data and Photos

Data and photos for all Phase 1 - fish passage assessments are provided online at https://www.newgraphenvironment.com/fish_passage_peace_2024_reporting/appendix---phase-1-fish-passage-assessment-data-and-photos.html - with a pdf version available at https://github.com/NewGraphEnvironment/fish_passage_peace_2024_reporting/raw/main/docs/Appendix_1.pdf

Attachment - Habitat Assessment and Fish Sampling Data

All field data collected is available [here](#).

Habitat assessment data (including fish sampling and PIT tagging information) is available for download here https://github.com/NewGraphEnvironment/fish_passage_peace_2025_reporting/blob/main/data/habitat_confirmations.xls.

Raw fish data is available for download here https://github.com/NewGraphEnvironment/fish_passage_peace_2025_reporting/blob/main/data/fish_data_tags_joined.csv.