

Chapter 5 Smoothness and convexity

Siheng Zhang
zhangsiheng@cvte.com

2021 年 11 月 1 日

This part corresponds to **Chapter 1,3,4 of PRML, Chapter of UML**. It mainly introduces some important properties regarding with functions: convexity, smoothness, strong convexity and Lipschitzness, which are basis for the next chapters.

目录

1	Convexity	2
2	Strong convexity	3
3	Lipschitzness	4
4	Smoothness	4

1 Convexity

Definition 5.1 (convex set) A set C in a vector space is convex if for any two vectors \mathbf{u}, \mathbf{v} in C , the line segment between them is in C , that is, for any $\alpha \in [0, 1]$, $\alpha\mathbf{u} + (1 - \alpha)\mathbf{v} \in C$.

Definition 5.2 (convex function) Let C be a convex set. A function $f : C \rightarrow \mathcal{R}$ is convex if for all $\mathbf{u}, \mathbf{v} \in C$ and $\alpha \in [0, 1]$,

$$f(\alpha\mathbf{u} + (1 - \alpha)\mathbf{v}) \leq \alpha f(\mathbf{u}) + (1 - \alpha)f(\mathbf{v})$$

Claim 5.1 local minimum of convex function is global minimum

Proof Let $B(\mathbf{u}, r) = \{\mathbf{v} : \|\mathbf{v} - \mathbf{u}\| \leq r\}$, we say that $f(\mathbf{u})$ is a local minimum if there exists some $r > 0$ such that $\forall \mathbf{v} \in B(\mathbf{u}, r), f(\mathbf{v}) \geq f(\mathbf{u})$. Then for any \mathbf{v} (not necessarily in B), there exists a small enough $\alpha > 0$ such that $\mathbf{u} + \alpha(\mathbf{v} - \mathbf{u}) \in B(\mathbf{u}, r)$, and therefore $f(\mathbf{u}) \leq f(\mathbf{u} + \alpha(\mathbf{v} - \mathbf{u}))$. If f is convex, we also have $f(\mathbf{u} + \alpha(\mathbf{v} - \mathbf{u})) \leq (1 - \alpha)f(\mathbf{u}) + \alpha f(\mathbf{v})$, which leads to $f(\mathbf{u}) \leq f(\mathbf{v})$ and hence is a global minimum.

Claim 5.2 For a convex function f , we can construct a tangent at any point that lies below the function everywhere. If f is differential, then we have

$$\forall \mathbf{u}, f(\mathbf{u}) \geq f(\mathbf{v}) + \nabla f(\mathbf{v})^\top (\mathbf{u} - \mathbf{v})$$

To generalize this inequality to non-differential functions, we should study for sub-gradient.

Definition 5.3 (sub-gradient) Define the sub-gradient of a function f at \mathbf{x} to be a vector $\mathbf{g} \in \mathcal{R}^d$ which satisfies that $f(\mathbf{y}) \geq f(\mathbf{x}) + \mathbf{g}^\top (\mathbf{y} - \mathbf{x}), \forall \mathbf{y} \in \mathcal{R}^d$.

Theorem 5.1 Let f be a differential function, then f is convex iff. ∇f is monotonically non-decreasing, and iff. $\nabla^2 f$ is non-negative.

Ex1 $f(\mathbf{x}) = \mathbf{x}^\top \mathbf{x}$ is convex. To see this, note that $\nabla f(\mathbf{x}) = 2\mathbf{x}$ and $\nabla^2 f(\mathbf{x}) = 2$.

Ex2 Consider scalar function $f(x) = \log(1 + \exp(x))$. It is convex since $f'(x) = \frac{1}{1 + \exp(-x)}$, which is the sigmoid function, and $f''(x) = \frac{\exp(-x)}{(1 + \exp(-x))^2} > 0$.

Claim 5.1 (linear transformation preserves convexity) Assume that $g : \mathcal{R}^d \rightarrow \mathcal{R}$ can be written as $g(\mathbf{w}) = f(\mathbf{w}^\top \mathbf{x} + b)$, for some $\mathbf{x} \in \mathcal{R}^d, b \in \mathcal{R}$. Then, if f is convex, g is convex too.

Proof For $\mathbf{w}_1, \mathbf{w}_2 \in \mathcal{R}^d$, and $\alpha \in (0, 1)$

$$\begin{aligned} g(\alpha\mathbf{w}_1 + (1 - \alpha)\mathbf{w}_2) &= f(\alpha\mathbf{w}_1^\top \mathbf{x} + (1 - \alpha)\mathbf{w}_2^\top \mathbf{x} + b) \\ &= f(\alpha(\mathbf{w}_1^\top \mathbf{x} + b) + (1 - \alpha)(\mathbf{w}_2^\top \mathbf{x} + b)) \\ &\leq \alpha f(\mathbf{w}_1^\top \mathbf{x} + b) + (1 - \alpha)f(\mathbf{w}_2^\top \mathbf{x} + b) = \alpha g(\mathbf{x}_1) + (1 - \alpha)g(\mathbf{x}_2) \end{aligned}$$

Following Claim 5.1,

Ex3 $f(\mathbf{w}) = (\mathbf{w}^\top \mathbf{x} - b)^2$ is convex.

Ex4 $f(\mathbf{w}) = \log(1 + \exp(-y\mathbf{w}^\top \mathbf{x}))$ is convex.

Claim 5.2 (Pointwise supremum of convex functions is convex) For $i = 1, \dots, r$, let $f_i : \mathcal{R}^d \rightarrow \mathcal{R}$ be a convex function, then $g(\mathbf{x}) = \max_i f_i(\mathbf{x})$ is convex.

Proof For $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{R}^d$, and $\alpha \in (0, 1)$

$$\begin{aligned} g(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2) &= \max_i f_i(\alpha\mathbf{x}_1 + (1 - \alpha)\mathbf{x}_2) \leq \max_i [\alpha f_i(\mathbf{x}_1) + (1 - \alpha)f_i(\mathbf{x}_2)] \\ &\leq \alpha \max_i f_i(\mathbf{x}_1) + (1 - \alpha) \max_i f_i(\mathbf{x}_2) = \alpha g(\mathbf{x}_1) + (1 - \alpha)g(\mathbf{x}_2) \end{aligned}$$

Following Claim 5.2,

Ex5 $g(\mathbf{x}) = |\mathbf{x}|$ is convex. Since $f_1(\mathbf{x}) = \mathbf{x}$, $f_2(\mathbf{x}) = -\mathbf{x}$ are both convex, and $g(\mathbf{x}) = \max\{f_1(\mathbf{x}), f_2(\mathbf{x})\}$.

Claim 5.3 (linear combination with non-negative weights preserves convexity) For $i = 1, \dots, r$, let $f_i : \mathcal{R}^d \rightarrow \mathcal{R}$ be a convex function, then $\forall w_i \geq 0, i \in [1, r], g(\mathbf{x}) = \sum_i w_i f_i(\mathbf{x})$ is convex.

Proof For $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{R}^d$, and $\alpha \in (0, 1)$

$$\begin{aligned} g(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) &= \sum_i w_i f_i(\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2) \leq \sum_i w_i (\alpha f_i(\mathbf{x}_1) + (1 - \alpha) f_i(\mathbf{x}_2)) \\ &= \alpha \sum_i w_i f_i(\mathbf{x}_1) + (1 - \alpha) \sum_i w_i f_i(\mathbf{x}_2) = \alpha g(\mathbf{x}_1) + (1 - \alpha) g(\mathbf{x}_2) \end{aligned}$$

Theorem 5.2 Jensen's Inequality Let $f : \mathcal{R}^d \rightarrow \mathcal{R}$ be a measurable convex function and $x \in \mathcal{R}^d$, $\mathbb{E}[\xi]$ exists. Then

$$\mathbb{E}[f(\mathbf{x})] \geq f(\mathbb{E}[\mathbf{x}])$$

Ex6 Consider function $f(x) = |x|$, then the sub-differential set $\partial f(x)$ is

$$\partial f(x) = \begin{cases} \{1\}, & x > 0 \\ [-1, 1], & x = 0 \\ \{-1\}, & x < 0 \end{cases}$$

Ex7 *Sub-gradient of Hinge loss.* Consider the loss $l(x) = \max(1 - \mathbf{z}^\top \mathbf{x}, 0)$, then the sub-differential set is

$$\partial f(x) = \begin{cases} \{0\}, & 1 - \mathbf{z}^\top \mathbf{x} < 0 \\ \{-\alpha \mathbf{z}, \alpha \in [0, 1]\}, & 1 - \mathbf{z}^\top \mathbf{x} = 0 \\ \{-\mathbf{z}\}, & 1 - \mathbf{z}^\top \mathbf{x} > 0 \end{cases}$$

2 Strong convexity

Definition 5.5 (strongly convex function) A function f is λ -strongly convex if for all $\mathbf{u}, \mathbf{v} \in C$ and $\alpha \in [0, 1]$,

$$f(\alpha \mathbf{u} + (1 - \alpha) \mathbf{v}) \leq \alpha f(\mathbf{u}) + (1 - \alpha) f(\mathbf{v}) - \frac{\lambda}{2} \alpha (1 - \alpha) \|\mathbf{u} - \mathbf{v}\|_2^2$$

Theorem 5.3 If f is λ -strongly convex then for every \mathbf{u}, \mathbf{v} and $\mathbf{g} \in \partial f(\mathbf{v})$ we have

$$f(\mathbf{v}) \geq f(\mathbf{u}) + \mathbf{g}^\top (\mathbf{u} - \mathbf{v}) + \frac{\lambda}{2} \|\mathbf{u} - \mathbf{v}\|_2^2$$

Proof According to the definition,

$$\begin{aligned} f(\mathbf{u}) &\geq \frac{f(\alpha \mathbf{u} + (1 - \alpha) \mathbf{v})}{\alpha} - \frac{1 - \alpha}{\alpha} f(\mathbf{v}) + \frac{\lambda}{2} (1 - \alpha) \|\mathbf{u} - \mathbf{v}\|_2^2 \\ &\geq \frac{f(\mathbf{v}) + \mathbf{g}^\top (\alpha \mathbf{u} + (1 - \alpha) \mathbf{v}) - \mathbf{v}}{\alpha} - \frac{1 - \alpha}{\alpha} f(\mathbf{v}) + \frac{\lambda}{2} (1 - \alpha) \|\mathbf{u} - \mathbf{v}\|_2^2 \quad (\text{Claim 5.2}) \\ &= \frac{f(\mathbf{v}) + \alpha \mathbf{g}^\top (\mathbf{u} - \mathbf{v})}{\alpha} - \frac{1 - \alpha}{\alpha} f(\mathbf{v}) + \frac{\lambda}{2} (1 - \alpha) \|\mathbf{u} - \mathbf{v}\|_2^2 \\ &= f(\mathbf{v}) + \mathbf{g}^\top (\mathbf{u} - \mathbf{v}) + \frac{\lambda}{2} (1 - \alpha) \|\mathbf{u} - \mathbf{v}\|_2^2 \end{aligned}$$

Note that it holds for all $\alpha \in [0, 1]$, setting $\alpha = 0$ leads to the conclusion.

Corollary 5.1 Following this theorem, if f is λ -strongly convex and \mathbf{v} is a minimizer of f , then for any \mathbf{u} , $f(\mathbf{v}) \geq f(\mathbf{u}) + \frac{\lambda}{2} \|\mathbf{u} - \mathbf{v}\|_2^2$.

Ex8 $f(\mathbf{x}) = \|\mathbf{x}\|_2^2$ is 2-strongly convex, $f(\mathbf{x}) = \lambda\|\mathbf{x}\|_2^2$ is 2λ -strongly convex. Note that $f(\mathbf{x}') = \lambda\|\mathbf{x}' - \mathbf{x} + \mathbf{x}\|_2^2 = \lambda\|\mathbf{x}\|_2^2 + 2\lambda\mathbf{x}^\top(\mathbf{x}' - \mathbf{x}) + \lambda\|\mathbf{x}' - \mathbf{x}\|_2^2$, in which $2\lambda\mathbf{x}$ is the gradient of $f(\mathbf{x})$.

Claim 5.4 If f is λ -strongly convex and g is convex, then $f + g$ is λ -strongly convex.

Following **Claim 5.4**, if we will to optimize a convex loss function, then with l_2 -norm regularization, the loss function becomes strongly convex. This speeds up the convergence of learning procedure, which we will see in the next chapter.

3 Lipschitzness

Definition 5.6 (Lipschitzness) A function $f : \mathcal{R}^d \rightarrow \mathcal{R}^k$ is ρ -Lipschitz over set C if for every $\mathbf{w}_1, \mathbf{w}_2 \in C$, $\|f(\mathbf{w}_1) - f(\mathbf{w}_2)\| \leq \rho\|\mathbf{w}_1 - \mathbf{w}_2\|$. (Note that it can be any norm, not only for Euclidean norm.)

With regard to l_2 -norm, according to the mean value theorem, we have $f(\mathbf{w}_1) - f(\mathbf{w}_2) = g^\top(\mathbf{w}_1 - \mathbf{w}_2)$, where $g \in \partial f(\mathbf{u})$, \mathbf{u} is some point between \mathbf{w}_1 and \mathbf{w}_2 . So Lipschitzness is equivalent to bound the l_2 norm of sub-gradients by ρ everywhere.

Theorem 5.4 If f is Lipschitz w.r.t l_2 -norm, then for all $g \in \partial f(\mathbf{x})$, we have $\|g\|_2 \leq \rho$.

Ex9 The function $f(x) = |x|$ is 1-Lipschitz over \mathcal{R} . Since $|x_1| - |x_2| = |x_1 - x_2 + x_2| - |x_2| \leq |x_1 - x_2| + |x_2| - |x_2| = |x_1 - x_2|$.

Claim 5.5 (Lipschitzness preserves on compound functions)

4 Smoothness

Definition 5.7 (dual norm)

Definition 5.8 (smooth function)

remark1. 1

4.4 According to definition, the dual norm of $\|\cdot\|_p (p \geq 1)$ is $\|\theta\|_* = \max_{\mathbf{x}: \|\mathbf{x}\|_p \leq 1} \langle \theta, \mathbf{x} \rangle$.

Using the Hölder inequality

$$\|\langle \mathbf{z}, \mathbf{x} \rangle\|_1 \leq \|\mathbf{x}\|_q \|\mathbf{z}\|_p$$

where $\frac{1}{p} + \frac{1}{q} = 1$, we have that:

$$\|\theta\|_* \leq \|\mathbf{x}\|_q \|\mathbf{z}\|_p \leq \|\mathbf{x}\|_q$$

So the dual norm is $\|\cdot\|_q$, where $\frac{1}{p} + \frac{1}{q} = 1$.

4.5 Start from strong convexity, we have that

$$\sum_{t=1}^T l_t(\mathbf{x}_t) - \sum_{t=1}^T l_t(\mathbf{u}) \leq \sum_{t=1}^T \left(\frac{1}{2\eta_t} \|\mathbf{x}_t - \mathbf{u}\|_2^2 - \frac{1}{2\eta_t} \|\mathbf{x}_{t+1} - \mathbf{u}\|_2^2 - \frac{\mu_t}{2} \|\mathbf{x}_t - \mathbf{u}\|_2^2 + \frac{\eta_t}{2} \|\mathbf{g}_t\|_2^2 \right)$$

Let all $\mu_t \leq \mu$, choose $\eta_t = 1/\mu t$. Given that the loss functions are Lipschitz, we can further bound the regret to be

$$\sum_{t=1}^T l_t(\mathbf{x}_t) - \sum_{t=1}^T l_t(\mathbf{u}) \leq \sum_{t=1}^T \frac{\eta_t}{2} \|\mathbf{g}_t\|_2^2 = \frac{L^2}{2\mu} \sum_{t=1}^T \frac{1}{t}$$

Since that the loss functions are smooth, t can be rewrite as $t \geq \sum_{i=1}^t \frac{L \|\mathbf{x} - \mathbf{u}\|_2^2}{\|l_i(\mathbf{x}) - l_i(\mathbf{u})\|_2^2} \geq \frac{L}{M} \sum_{i=1}^t \frac{\|\mathbf{g}_i\|_2^2}{\|l_i(\mathbf{x}) - l_i(\mathbf{u})\|_2^2}$, so $\frac{1}{t} \leq \frac{M}{L} \frac{1}{\sum_{i=1}^t \frac{\|\mathbf{g}_i\|_2^2}{\|l_i(\mathbf{x}) - l_i(\mathbf{u})\|_2^2}}$.

Now using the fact that $\sum_{t=1}^T \frac{1}{t} \leq 1 + \ln T$, we can also make integral by $\|l_i(\mathbf{x}) - l_i(\mathbf{u})\|_2^2$ and get that

$$\sum_{t=1}^T l_t(\mathbf{x}_t) - \sum_{t=1}^T l_t(\mathbf{u}) \leq O(\ln(1 + [\sum_{t=1}^T l_t(\mathbf{x}_t) - \sum_{t=1}^T l_t(\mathbf{u})] \sum_{t=1}^T \sum_{t=1}^T l_t(\mathbf{u})))$$

Using the fact that $x \leq a \ln(1 + bx) + c$ leads to $x \leq a \ln(2ab \ln(ab) + 2bc + 2) + c$, it leads to

$$\sum_{t=1}^T l_t(\mathbf{x}_t) - \sum_{t=1}^T l_t(\mathbf{u}) \leq O(1 + \ln \sum_{t=1}^T l_t(\mathbf{u}))$$

which is the $O(1 + \ln L^*)$ bound.

4.6 With regard to $\|\cdot\|_2$, we need to prove that

$$\|l''(\mathbf{x}_1)\|_2 \leq \frac{1}{4}$$

Note that

$$l'(\mathbf{x}) = \frac{-y\mathbf{z}e^{-y\langle \mathbf{z}, \mathbf{x} \rangle}}{1 + e^{-y\langle \mathbf{z}, \mathbf{x} \rangle}} = -y\mathbf{z}(1 - \exp(-l(\mathbf{x})))$$

$$l''(\mathbf{x}) = y\mathbf{z} \exp(-l(\mathbf{x}))(1 - \exp(-l(\mathbf{x})))$$

Since $y \in \{-1, 1\}$ and $\|\mathbf{z}\|_2 \leq 1$, we can bound it by

$$\|l''(\mathbf{x}_1)\|_2 \leq |\exp(-l(\mathbf{x}))(1 - \exp(-l(\mathbf{x})))|$$

Note that $l(\mathbf{x}) > 0$, so $0 < \exp(-l(\mathbf{x})) < 1$, so the upper bound is $1/4$.