Discussion

<u>Help</u>

shengtatng >

★ Course / UNIT 3 / Lecture 9 - Sampling and Standard Error

<u>Dates</u>

()



<u>Calendar</u>

<u>Notes</u>

Exercise 5

<u>Course</u>


□ Bookmark this page

<u>Progress</u>

Exercise 5

3/3 points (graded)

You are given two data files. Each file contains 1800 data points measuring the heart rate (in beats per minute, every 0.5 seconds) of a subject prforming comparable activities for the duration of 15 minutes: https://doi.org/10.15/ and https://doi.org/10.15/ and https:



Using a sample size of 250, decide whether the following methods of drawing samples will yield samples where the examples are independent of each other.

| 1 | Usina | random.sample |
|----|--------|----------------|
| ι. | OSILIG | random. Sampte |

| ✓ | Examples | are inde | pendent i | n hr1 | sample |
|---|----------|----------|-----------|-------|--------|

| ✓ | Examples are independent in hr2 sample |
|----------|--|



2. Getting a random number between 1 and 1800, 250 times.

| Examples are independent in hr1 sample. |
|---|
|---|

| sample. |
|---------|
| |




3. Starting at the first example and going until the 500th example.

| Examples are independent in hr1 sample. | |
|---|--|
| Examples are independent in hr2 sample. | |

| | Neither h1 nor h2 give independent examples. |
|----------|--|
| Y | Neither III nor itz give independent examples. |



Submit





edX

About

Affiliates

edX for Business

Open edX

Careers

News

Legal

Terms of Service & Honor Code

Privacy Policy

Accessibility Policy

Trademark Policy

<u>Sitemap</u>

Connect

<u>Blog</u>

Contact Us

Help Center

<u>Security</u>

Media Kit















© 2022 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>