

Process	Biochemical reaction	Formulation	Description
Gross primary production (GPP)	$106\text{CO}_2 + 16\text{HNO}_3 + \text{H}_3\text{PO}_4 + 122\text{H}_2\text{O} + (\text{sunlight}) \rightarrow (\text{CH}_2\text{O})_{106}(\text{NH}_3)_{16}\text{H}_3\text{PO}_4 + 138\text{O}_2$ <p>Nutrient limitation for phytoplankton growth</p> $\text{nlim} = \frac{\text{NO}_3 + \text{NH}_4}{\text{NO}_3 + \text{NH}_4 + K_N} \times \frac{\text{PO}_4}{\text{PO}_4 + K_{\text{PO}_4}} \times \frac{DSi}{DSi + K_{Si}}$	$\text{GPP} = P_{max}^B(T) \times \text{nlim} \times \text{phytoplankton} \times \int_H^0 1 - \exp\left(-\frac{\alpha}{P_{max}^B(T)} \times I(0) \times \exp(-K_D \cdot H)\right) dz$ <p>P_{max}^B: photosynthesis rate nlim: Nutrient limitation for phytoplankton growth</p>	<p>GPP refers to the total rate of organic carbon production by phytoplankton based on the rate of photosynthesis</p> <p>$I(0)$: solar radiation K_D: Light extinction coefficient H: water depth</p>
Net primary production (NPP)	<p>NPP is the rate of phytoplankton produces biomass which already subtract the respiration of primary producers, including:</p> <p>k_{excr} phytoplankton excretion k_{maint} phytoplanktonic maintenance</p>	$\text{NPP} = \frac{\text{GPP}}{H} \times (1 - k_{\text{excr}}) \times (1 - k_{\text{growth}}) - k_{\text{maint}}(T) \times \text{phytoplankton}$ <p>k_{growth} growth constants of phytoplankton</p>	<p>NPP is GPP minus the autotrophs' respiration rate (i.e., only by the primary producers).</p>
Nitrification	$\text{NH}_4^+ + 2\text{O}_2 \rightarrow \text{NO}_3^- + \text{H}_2\text{O} + 2\text{H}^+$	$N = k_{\text{nit}}(T) \times \frac{\text{NH}_4}{\text{NH}_4 + K_{\text{NH}_4}} \times \frac{\text{O}_2}{\text{O}_2 + K_{\text{O}_2,\text{nit}}}$ <p>$k_{\text{nit}}(T)$ maximum rate constant</p>	<p>Under aerobic conditions, ammonia is oxidized to nitrite and nitrate via nitrification</p> <p>K_{NH_4} half-saturation constants</p>
Denitrification	$94.4\text{HNO}_3 + (\text{CH}_2\text{O})_{106}(\text{NH}_3)_{16}(\text{H}_3\text{PO}_4) \rightarrow 106\text{CO}_2 + 55.2\text{N}_2 + \text{H}_3\text{PO}_4 + 177.2\text{H}_2\text{O}$	$D = k_{\text{denit}}(T) \times \frac{\text{TOC}}{\text{TOC} + K_{\text{TOC}}} \times \frac{\text{NO}_3}{\text{NO}_3 + K_{\text{NO}_3}} \times \frac{K_{\text{in},\text{O}_2}}{\text{O}_2 + K_{\text{in},\text{O}_2}}$	<p>Under anaerobic conditions, nitrate is reduced to gas forms as N_2, N_2O while organic P is degraded to inorganic PO_4^{3-}</p>
Aerobic degradation (respiration)	$138\text{O}_2 + (\text{CH}_2\text{O})_{106}(\text{NH}_3)_{16}(\text{H}_3\text{PO}_4) \rightarrow 106\text{CO}_2 + 16\text{HNO}_3 + \text{H}_3\text{PO}_4 + 122\text{H}_2\text{O}$	$R = k_{\text{ox}}(T) \cdot \frac{\text{TOC}}{\text{TOC} + K_{\text{TOC}}} \cdot \frac{\text{O}_2}{\text{O}_2 + K_{\text{O}_2,\text{ox}}}$	<p>Degradation of organic carbon in the aerobic condition that converts into inorganic matters</p>
Phytoplankton = NPP - Phytoplankton mortality		TOC = Phytoplankton mortality – Aerobic degradation – Denitrification	
DSi = NPP of Diatom = NPP x Redfield ratio for silica (15/106)		O_2 = Oxygen air exchange + NPP – Aerobic degradation – Nitrification	
NH_4^+ = Aerobic degradation – Nitrification – NPP_{NH_4}		PO_4^{3-} = Aerobic degradation + Denitrification – NPP_{PO_4} – PO_4 adsorption	
NO_3^- = Nitrification - Denitrification – NPP_{NO_3}			