

Renal Unit Clinical Protocol RU-07

Management of Anaemia in Chronic Kidney Disease (ICD-10 D63.1) with Epoetin alfa

Applies to adult haemodialysis patients with haemoglobin persistently below 100 g/L after iron repletion. Epoetin alfa is given by subcutaneous injection using prefilled 4000 IU syringes. Standard aseptic technique: one alcohol swab per injection and one pair of sterile gloves per administration.

Phase 1 - Correction (28-day cycle, 3 cycles)

Epoetin alfa 4000 IU subcutaneously three times per week (12 injections per 28-day cycle). Review haemoglobin at the end of each cycle.

Item	Dose / use	Per 28-day cycle
Epoetin alfa 4000IU prefilled syringe	1 syringe per injection, 3x weekly	12 syringes
Alcohol swab	1 per injection	12 swabs
Sterile gloves	1 pair per administration	12 pairs

Phase 2 - Maintenance (28-day cycle, ongoing)

Once haemoglobin reaches 100-115 g/L, reduce to Epoetin alfa 4000 IU once weekly (4 injections per 28-day cycle). Continue for as long as the patient remains on dialysis.

Item	Dose / use	Per 28-day cycle
Epoetin alfa 4000IU prefilled syringe	1 syringe per injection, weekly	4 syringes
Alcohol swab	1 per injection	4 swabs
Sterile gloves	1 pair per administration	4 pairs