


Product datasheet

Anti-Tubulin antibody [YOL1/34] - Microtubule Marker ab6161

★★★★★ 17 Abreviews 48 References 10 Images

Overview

Product name	Anti-Tubulin antibody [YOL1/34] - Microtubule Marker
Description	Rat monoclonal [YOL1/34] to Tubulin - Microtubule Marker
Tested applications	Suitable for: WB, ELISA, IHC-FoFr, RIA, Flow Cyt, ICC/IF
Species reactivity	Reacts with: Mouse, Rat, Dog, Human, Saccharomyces cerevisiae, Schizosaccharomyces pombe, Alligator Predicted to work with: Plants, a wide range of other species, Mammal 
Immunogen	Full length native protein (purified) (S. cerevisiae).
Epitope	ab6161 binds to an epitope between amino acids 414 and 422 of alpha tubulin.
Positive control	WB: HeLa and NIH3T3 whole cell lysates and rat brain tissue lysate. Flow Cyt: methanol fixed/tween permeabilised HeLa cells.
General notes	<p>We can conjugate this antibody to FITC for you (please see ab150252 for details). This antibody can be used as a Western blotting loading control (Kops et al.) and as a Microtubule Marker.</p> <p>Has been used for the selection of specific recombinant antibodies engineered to incorporate its epitope. It is also useful for studying the function of microtubules.</p> <p>Alternative versions available:</p> <p>Anti-Tubulin antibody - Microtubule Marker (Alexa Fluor® 488) [YOL1/34] (ab195883)</p> <p>Anti-Tubulin antibody - Microtubule Marker (Alexa Fluor® 647) [YOL1/34] (ab195884)</p> <p>Anti-Tubulin antibody - Microtubule Marker (HRP) [YOL1/34] (ab196583)</p> <p>Anti-Tubulin antibody - BSA and Azide free [YOL1/34] (ab174643)</p> <p>Anti-Tubulin antibody (FITC) [YOL1/34] (ab150252)</p>

Properties

Form	Liquid
Storage instructions	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
Storage buffer	<p>pH: 7.40</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituent: PBS</p> <p>batches contain 0.4M Arginine</p>

Purity	IgG fraction
Primary antibody notes	Has been used for the selection of specific recombinant antibodies engineered to incorporate its epitope. It is also useful for studying the function of microtubules.
Clonality	Monoclonal
Clone number	YOL1/34
Isotype	IgG2a

Applications

Our [Abpromise guarantee](#) covers the use of **ab6161** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

Application	Abreviews	Notes
WB	★★★★★	Use a concentration of 1 µg/ml.
ELISA		1/4000.
IHC-FoFr		1/600. PubMed: 15831501 Suggested working dilution of 1/600.
RIA		Use at an assay dependent concentration.
Flow Cyt		Use 1µg for 10 ⁶ cells. ab18450 - Rat monoclonal IgG2a, is suitable for use as an isotype control with this antibody.
ICC/IF	★★★★★	Use a concentration of 5 µg/ml.

Target

Function	Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha-chain.
Sequence similarities	Belongs to the tubulin family.
Post-translational modifications	<p>Undergoes a tyrosination/detyrosination cycle, the cyclic removal and re-addition of a C-terminal tyrosine residue by the enzymes tubulin tyrosine carboxypeptidase (TTCP) and tubulin tyrosine ligase (TTL), respectively.</p> <p>Some glutamate residues at the C-terminus are polyglutamylated. This modification occurs exclusively on glutamate residues and results in polyglutamate chains on the gamma-carboxyl group. Also monoglycylated but not polyglycylated due to the absence of functional TTL10 in human. Monoglycylation is mainly limited to tubulin incorporated into axonemes (cilia and flagella) whereas glutamylation is prevalent in neuronal cells, centrioles, axonemes, and the mitotic spindle. Both modifications can coexist on the same protein on adjacent residues, and lowering glycylation levels increases polyglutamylated, and reciprocally. The precise function of such modifications is still unclear but they regulate the assembly and dynamics of axonemal microtubules.</p> <p>Acetylation of alpha-tubulins at Lys-40 stabilizes microtubules and affects affinity and processivity of microtubule motors. This modification has a role in multiple cellular functions, ranging from cell motility, cell cycle progression or cell differentiation to intracellular trafficking and signaling.</p>
Cellular localization	Cytoplasm > cytoskeleton.

Anti-Tubulin antibody [YOL1/34] - Microtubule Marker images




Western blot - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

Western blot against tubulin with ab6161 at 1/3000. Secondary Rabbit anti-Rat IgG HRP (ab6734) was used at 1/2000. Exposure time: 2mins.

Lane 1: 20µg/lane HeLa (Human) whole cell lysates (ab7898).

Lane 2: 20µg/lane 3T3 (Mouse) whole cell lysate (ab7901).


Lane 3: 20µg/lane Rat brain tissue lysate (ab7942).



Immunocytochemistry/ Immunofluorescence - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)


Confocal image of 21 day in vitro rat hippocampal neurons, stained with rat monoclonal antibody to Tubulin - Microtubule Marker (ab6161) in green at 1/500 and Microtubule Associated protein 2 in blue.

This picture was kindly supplied as part of the review submitted by Dr Jonathon Burman.




Flow Cytometry - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

Overlay histogram showing HeLa cells stained with ab6161 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab6161, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rat IgG (H+L) ([ab98386](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rat IgG2a [aRTK2758] ([ab18450](#), 1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.



Immunocytochemistry/ Immunofluorescence - Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

ICC/IF image of ab6161 stained HeLa cells. The cells were 100% methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab6161, 1µg/ml) overnight at +4°C. The secondary antibody (green) was DyLight® 488 goat anti-rat- H&L, pre-adsorbed ([ab98420](#)) used at a 1/250 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.




Immunocytochemistry/ Immunofluorescence - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

Cultured human macrophages were used with ab6161 at 1/1000 for immunofluorescence. Cells were fixed with cold 2% formaldehyde for 20mins.

Green staining is Alexa 568, Blue staining is DAPI stain.

This cell represents a young macrophage, the staining patterns varied as the cells aged in culture.



All lanes : Anti-Tubulin antibody [YOL1/34] -
Microtubule Marker (ab6161) at 1/2000
dilution

Lane 1 : Yeast whole cell extract prepared by
bead-beating

Lane 2 : Yeast whole cell extract prepared by
bead-beating

Lane 3 : Yeast whole cell extract prepared by
bead-beating

Lane 4 : Yeast whole cell extract prepared by
bead-beating

Lane 5 : Yeast whole cell extract prepared by
bead-beating

Lane 6 : Yeast whole cell extract prepared by
bead-beating

Lane 7 : Yeast whole cell extract prepared by
bead-beating

Lane 8 : Yeast whole cell extract prepared by
bead-beating

Lane 9 : Yeast whole cell extract prepared by
bead-beating

Lane 10 : Yeast whole cell extract prepared
by bead-beating

Lysates/proteins at 5 µg per lane.

Secondary


HRP conjugated goat anti-rat antibody
Developed using the ECL technique

Performed under reducing conditions.

Observed band size : 50 kDa


Exposure time : 30 seconds

*This image is courtesy of an anonymous
Abreview*



Immunocytochemistry/ Immunofluorescence - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

ICC/IF image of ab6161 stained human HepG2 cells. The cells were 4% PFA fixed (10 min), permeabilised in 0.1% PBS-Tween (20 min) and incubated with the antibody (ab6161, 1 µg/ml) for 1h at room temperature. 1%BSA / 10% normal goat serum / 0.3M glycine was used to block non-specific protein-protein interactions. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rat IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red). DAPI was used to stain the cell nuclei (blue). This antibody also gave a positive IF result in HeLa, HEK 293 and MCF7 cells.



Western blot - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161) at 1 µg/ml + Brain (Rat) Tissue Lysate at 10 µg


Secondary

Rabbit polyclonal to Rat IgG - H&L (HRP) at 1/10000 dilution
Developed using the ECL technique

Performed under reducing conditions.

Observed band size : 54 kDa


Exposure time : 3 minutes



Immunocytochemistry/ Immunofluorescence - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

This image is courtesy of an anonymous Abreview

ab6161 staining mouse NIH 3T3 fibroblast cells by ICC/IF. Cells were PFA fixed and permeabilized in 0.2% Triton X-100 prior to blocking in 5% BSA for 45 minutes at RT. The primary antibody was diluted 1/1000 and incubated with the sample for 1 hour. An Alexa Fluor® 568 conjugated goat anti-rat antibody, diluted 1/3000, was used as the secondary.



Immunocytochemistry/ Immunofluorescence - Anti-Tubulin antibody [YOL1/34] - Microtubule Marker (ab6161)

ab6161 staining tubulin HeLa cells treated with anisomycin ([ab120495](#)), by ICC/IF. Increase in tubulin expression correlates with increased concentration of anisomycin as described in literature.

The cells were incubated at 37°C for 6h in media containing different concentrations of [ab120495](#) (anisomycin) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with ab6161 (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rat polyclonal antibody ([ab98386](#)) at 1/250 dilution was used as the secondary antibody. Nuclei were counterstained with DAPI and are shown in blue.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.com/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors